

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
19 September 2002 (19.09.2002)

PCT

(10) International Publication Number  
**WO 02/072786 A2**

(51) International Patent Classification<sup>7</sup>: **C12N**

(21) International Application Number: **PCT/US02/07903**

(22) International Filing Date: **13 March 2002 (13.03.2002)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:  
**60/275,592 13 March 2001 (13.03.2001) US**

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:  
**US 60/275,592 (CIP)**  
**Filed on 13 March 2001 (13.03.2001)**

(71) Applicant (for all designated States except US): **CORVAS INTERNATIONAL, INC.** [US/US]; 3030 Science Park Road, San Diego, CA 92121 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MADISON, Edwin, L.** [US/US]; 11005 Cedarcrest Way, San Diego, CA 92121 (US). **ONG, Edgar, O.** [CA/US]; 8328 Regents Road, #3C, San Diego, CA 92122 (US).

(74) Agents: **SEIDMAN, Stephanie, L. et al.**; Heller Ehrman White & McAuliffe LLP, 4350 La Jolla Village Drive, San Diego, CA 92122-1246 (US).

(81) Designated States (national): **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,**

**LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PI, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.**

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declaration under Rule 4.17:**

*as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)*

**Published:**

— *without international search report and to be republished upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **NUCLEIC ACID MOLECULES ENCODING A TRANSMEMBRANE SERINE PROTEASE 7, THE ENCODED POLYPEPTIDES AND METHODS BASED THEREON**

(57) Abstract: Provided herein are type II transmembrane serine protease 7 (MTSP7) polypeptides. Zymogen and activated forms of these polypeptides as well as single and two chain forms of the protease domain are also provided. Methods using the polypeptides to identify compounds that modulate the protease activity of an MTSP7 are provided.

WO 02/072786 A2

-1-

**NUCLEIC ACID MOLECULES ENCODING A TRANSMEMBRANE SERINE  
PROTEASE 7, THE ENCODED POLYPEPTIDES AND METHODS BASED THEREON  
RELATED APPLICATIONS**

Benefit of priority is claimed to U.S. provisional application Serial No.

- 5 60/275,592, filed March 13, 2001, to Edwin L. Madison and Edgar O. Ong,  
entitled "NUCLEIC ACID MOLECULES ENCODING A TRANSMEMBRANE SERINE  
PROTEASE, THE ENCODED PROTEINS AND METHODS BASED THEREON."

Where permitted, the subject matter this application is incorporated in its entirety  
by reference thereto.

**10 FIELD OF INVENTION**

Nucleic acid molecules that encode proteases and portions thereof,  
particularly protease domains are provided. Also provided are prognostic,  
diagnostic and therapeutic methods using the proteases and domains thereof and  
the encoding nucleic acid molecules.

**15 BACKGROUND OF THE INVENTION AND OBJECTS THEREOF**

- Cancer, which is a leading cause of death in the United States, is  
characterized by an increase in the number of abnormal neoplastic cells, which  
proliferate to form a tumor mass, the invasion of adjacent tissues by these  
neoplastic tumor cells, and the generation of malignant cells that metastasize via  
20 the blood or lymphatic system to regional lymph nodes and to distant sites.  
Among the hallmarks of cancer is a breakdown in the communication among  
tumor cells and their environment. Normal cells do not divide in the absence of  
stimulatory signals and cease dividing in the presence of inhibitory signals.  
Growth-stimulatory and growth-inhibitory signals, are routinely exchanged  
25 between cells within a tissue. In a cancerous, or neoplastic, state, a cell  
acquires the ability to "override" these signals and to proliferate under conditions  
in which normal cells do not grow.

- In order to proliferate tumor cells acquire a number of distinct aberrant  
traits reflecting genetic alterations. The genomes of certain well-studied tumors  
30 carry several different independently altered genes, including activated  
oncogenes and inactivated tumor suppressor genes. Each of these genetic

-2-

changes appears to be responsible for imparting some of the traits that, in the aggregate, represent the full neoplastic phenotype.

A variety of biochemical factors have been associated with different phases of metastasis. Cell surface receptors for collagen, glycoproteins such as laminin, and proteoglycans, facilitate tumor cell attachment, an important step in invasion and metastases. Attachment triggers the release of degradative enzymes which facilitate the penetration of tumor cells through tissue barriers. Once the tumor cells have entered the target tissue, specific growth factors are required for further proliferation. Tumor invasion and progression involve a complex series of events, in which tumor cells detach from the primary tumor, break down the normal tissue surrounding it, and migrate into a blood or lymphatic vessel to be carried to a distant site. The breaking down of normal tissue barriers is accomplished by the elaboration of specific enzymes that degrade the proteins of the extracellular matrix that make up basement membranes and stromal components of tissues.

A class of extracellular matrix degrading enzymes have been implicated in tumor invasion. Among these are the matrix metalloproteinases (MMP). For example, the production of the matrix metalloproteinase stromelysin is associated with malignant tumors with metastatic potential (see, *e.g.*, McDonnell *et al.* (1990) *Smnrs. in Cancer Biology* 1:107-115; McDonnell *et al.* (1990) *Cancer and Metastasis Reviews* 9:309-319).

The capacity of cancer cells to metastasize and invade tissue is facilitated by degradation of the basement membrane. Several proteinase enzymes, including the MMPs, have been reported to facilitate the process of invasion of tumor cells. MMPs are reported to enhance degradation of the basement membrane, which thereby permits tumorous cells to invade tissues. For example, two major metalloproteinases having molecular weights of about 70 kDa and 92 kDa appear to enhance ability of tumor cells to metastasize.

-3-

**Type II Transmembrane Serine Proteases (TTSPs)**

In addition to the MMPs, serine proteases have been implicated in neoplastic disease progression. Most serine proteases, which are either secreted enzymes or are sequestered in cytoplasmic storage organelles, have roles in blood coagulation, wound healing, digestion, immune responses and tumor invasion and metastasis. A class cell of surface proteins designated type II transmembrane serine proteases, which are membrane-anchored proteins with additional extracellular domains, has been identified. As cell surface proteins, they are positioned to play a role in intracellular signal transduction and in mediating cell surface proteolytic events.

Cell surface proteolysis is a mechanism for the generation of biologically active proteins that mediate a variety of cellular functions. These membrane-anchored proteins, include a disintegrin-like and metalloproteinase (ADAM) and membrane-type matrix metalloproteinase (MT-MMP). In mammals, at least 17 members of the TTSP family are known, including seven in humans (see, Hooper *et al.* (2001) *J. Biol. Chem.* 276:857-860). These include: corin (accession nos. AF133845 and AB013874; see, Yan *et al.* (1999) *J. Biol. Chem.* 274:14926-14938; Tomia *et al.* (1998) *J. Biochem.* 124:784-789; Uan *et al.* (2000) *Proc. Natl. Acad. Sci. U.S.A.* 97:8525-8529); enterpeptidase (also designated enterokinase; accession no. U09860 for the human protein; see, Kitamoto *et al.* (1995) *Biochem.* 27: 4562-4568; Yahagi *et al.* (1996) *Biochem. Biophys. Res. Commun.* 219:806-812; Kitamoto *et al.* (1994) *Proc. Natl. Acad. Sci. U.S.A.* 91:7588-7592; Matsushima *et al.* (1994) *J. Biol. Chem.* 269:19976-19982;); human airway trypsin-like protease (HAT; accession no. AB002134; see Yamaoka *et al.* *J. Biol. Chem.* 273:11894-11901); MTSP1 and matriptase (also called TADG-15; see SEQ ID Nos. 1 and 2; accession nos. AF133086/AF118224, AF04280022; Takeuchi *et al.* (1999) *Proc. Natl. Acad. Sci. U.S.A.* 96:11054-1161; Lin *et al.* (1999) *J. Biol. Chem.* 274:18231-18236; Takeuchi *et al.* (2000) *J. Biol. Chem.* 275:26333-26342; and Kim *et al.* (1999) *Immunogenetics* 49:420-429); hepsin (see, accession nos. M18930, AF030065, X70900; Leytus *et al.* (1988) *Biochem.* 27: 11895-11901; Vu *et al.* (1997) *J. Biol. Chem.* 272:31315-31320; and Farley *et al.* (1993) *Biochem. Biophys. Acta*



-4-

1173:350-352; and see, U.S. Patent No. 5,972,616); TMPRS2 (see, Accession Nos. U75329 and AF113596; Paoloni-Giacobino *et al.* (1997) *Genomics* 44:309-320; and Jacquinet *et al.* (2000) *FEBS Lett.* 468: 93-100); and TMPRSS4 (see, Accession No. NM 016425; Wallrapp *et al.* (2000) *Cancer* 60:2602-2606).

5           Serine proteases, including transmembrane serine proteases, have been implicated in processes involved in neoplastic development and progression. While the precise role of these proteases has not been elaborated, serine proteases and inhibitors thereof are involved in the control of many intra- and extracellular physiological processes, including degradative actions in cancer cell  
10 invasion, metastatic spread, and neovascularization of tumors, that are involved in tumor progression. It is believed that proteases are involved in the degradation of extracellular matrix (ECM) and contribute to tissue remodeling, and are necessary for cancer invasion and metastasis. The activity and/or expression of some proteases have been shown to correlate with tumor  
15 progression and development.

          For example, a membrane-type serine protease MTSP1 (also called matriptase; see SEQ ID Nos. 1 and 2 from U.S. Patent No. 5,972,616; and GenBank Accession No. AF118224; (1999) *J. Biol. Chem.* 274:18231-18236; U.S. Patent No. 5,792,616; see, also Takeuchi (1999) *Proc. Natl. Acad. Sci.*  
20 *U.S.A.* 96:11054-1161) that is expressed in epithelial cancer and normal tissue (Takeuchi *et al.* (1999) *Proc. Natl. Acad. Sci. USA*, 96(20):11054-61) has been identified. Matriptase was originally identified in human breast cancer cells as a major gelatinase (see, U.S. Patent No. 5,482,848), a type of matrix metalloprotease (MMP). It has been proposed that it plays a role in the  
25 metastasis of breast cancer. Matriptase also is expressed in a variety of epithelial tissues with high levels of activity and/or expression in the human gastrointestinal tract and the prostate. Other MTSPs, designated MTSP3, MTSP4 and MTSP6 and protease domains thereof, have been described in International PCT application No. PCT/US01/03471 and copending U.S.  
30 application Serial No. 09/776,191, filed February 2, 2001.

          Prostate-specific antigen (PSA), a kallikrein-like serine protease, degrades extracellular matrix glycoproteins fibronectin and laminin, and, has been

-5-

postulated to facilitate invasion by prostate cancer cells (Webber *et al.* (1995) *Clin. Cancer Res.*, 1(10):1089-94). Blocking PSA proteolytic activity with PSA-specific monoclonal antibodies results in a dose-dependent decrease *in vitro* in the invasion of the reconstituted basement membrane Matrigel by LNCaP  
5 human prostate carcinoma cells which secrete high levels of PSA.

Hepsin, a cell surface serine protease identified in hepatoma cells, is overexpressed in ovarian cancer (Tanimoto *et al.* (1997) *Cancer Res.*, 57(14):2884-7). The hepsin transcript appears to be abundant in carcinoma tissue and is almost never expressed in normal adult tissue, including normal  
10 ovary. It has been suggested that hepsin is frequently overexpressed in ovarian tumors and therefore can be a candidate protease in the invasive process and growth capacity of ovarian tumor cells.

A serine protease-like gene, designated normal epithelial cell-specific 1 (NES1) (Liu *et al.*, *Cancer Res.*, 56(14):3371-9 (1996)) has been identified.  
15 Although expression of the NES1 mRNA is observed in all normal and immortalized nontumorigenic epithelial cell lines, the majority of human breast cancer cell lines show a drastic reduction or a complete lack of its expression. The structural similarity of NES1 to polypeptides known to regulate growth factor activity and a negative correlation of NES1 expression with breast  
20 oncogenesis suggest a direct or indirect role for this protease-like gene product in the suppression of tumorigenesis.

Hence transmembrane serine proteases appear to be involved in the etiology and pathogenesis of tumors. There is a need to further elucidate their role in these processes and to identify additional transmembrane proteases.  
25 Therefore, it is an object herein to provide transmembrane serine protease (MTSP) proteins and nucleic acids encoding such MTSP proteases that are involved in the regulation of or participate in tumorigenesis and/or carcinogenesis. It is also an object herein to provide prognostic, diagnostic and therapeutic screening methods using such proteases and the nucleic acids  
30 encoding such proteases.

**SUMMARY OF THE INVENTION**

- Provided herein are members of the Transmembrane Serine Protease family, particularly the Type II Transmembrane Serine Protease (TTSP) family (also referred to herein as MTSPs), and more particularly TTSP family members
- 5 whose functional activity and/or expression differs in tumor cells from non-tumor cells in the same tissue. The MTSP provided herein is a heretofore unidentified MTSP family member, designated herein as MTSP7. The protease domain and full-length protein, including the zymogen and activated forms, and uses thereof are also provided. Proteins encoded by splice variants are also provided.
- 10 Assays for identifying effectors, such as small molecules and other conditions, that modulate the activation, expression or activity of MTSP7 are also provided herein. In exemplary assays, the affects of test compounds on the ability of a protease domain of MTSP7 to proteolytically cleave a known substrate, typically a fluorescently, chromogenically or otherwise detectably labeled substrate, are
- 15 assessed. Agents, generally compounds, particularly small molecules, that modulate the activity of the protease domain are candidate compounds for modulating the activity of the MTSP7. The protease domains can also be used to produce protease-specific antibodies. The protease domains provided herein include, but are not limited to, the single chain region having an N-terminus at
- 20 the cleavage site for activation of the zymogen, through the C-terminus, or C-terminal truncated portions thereof that exhibit proteolytic activity as a single-chain polypeptide in *vitro* proteolysis assays, of any MTSP family member, including MTSP7, generally from a mammal, including human, that, for example, is expressed in tumor cells at different levels from non-tumor cells.
- 25 Nucleic acid molecules encoding the proteins and protease domains are also provided. The nucleic acid and amino acid sequences of an exemplary full length MTSP7 are set forth in SEQ ID Nos. 15 and 16, and the protease domain is set forth in SEQ ID No. 17 and 18. Nucleic acid molecules that encode a single-chain protease domain or catalytically active portion thereof and also
- 30 those that encode the full-length MTSP7 are provided. Also provided are nucleic acid molecules that hybridize to such MTSP7-encoding nucleic acid along their full length and encode the protease domain or portion thereof are provided.

-7-

Hybridization is generally effected under conditions of at least low, generally at least moderate, and often high stringency.

The isolated nucleic acid fragment is DNA, including genomic or cDNA, or is RNA, or can include other components, such as protein nucleic acid. The  
5 isolated nucleic acid may include additional components, such as heterologous or native promoters, and other transcriptional and translational regulatory sequences, these genes may be linked to other genes, such as reporter genes or other indicator genes or genes that encode indicators.

Also provided is an isolated nucleic acid molecule that includes the  
10 sequence of molecules that is complementary to the nucleotide sequence encoding MTSP7 or the portion thereof.

Also provided are fragments thereof or oligonucleotides that can be used as probes or primers and that contain at least about 10, 14, 16 nucleotides, generally less than 1000 or less than or equal to 100, set forth in SEQ ID No. 15  
15 or 17 (or the complement thereof); or contain at least about 30 nucleotides (or the complement thereof) or contain oligonucleotides that hybridize along their full length (or at least about 70, 80 or 90% thereof) to any such fragments or oligonucleotides. The length of the fragments are a function of the purpose for which they are used and/or the complexity of the genome of interest. Generally  
20 probes and primers contain less than about 500, 150, 100 nucleotides.

Also provided are peptides that are encoded by such nucleic acid molecules. Included among those polypeptides are the MTSP7 protease domain or a polypeptide with amino acid changes such that the specificity and protease activity remains substantially unchanged. In particular, a substantially purified  
25 mammalian MTSP protein is provided that includes a serine protease catalytic domain and may additional include other domains. The MTSP7 can form homodimers and can also form heterodimers with some other protein, such as a membrane-bound protein. Also provided is a substantially purified protein including a sequence of amino acids that has at least 60%, 70%, 80%, 90% or  
30 about 95%, identity to the MTSP7 where the percentage identity is determined using standard algorithms and gap penalties that maximize the percentage identity. A human MTSP7 protein is exemplified, although other mammalian

-8-

MTSP7 proteins are contemplated. Splice variants of the MTSP7, particularly those with a proteolytically active protease domain, are contemplated herein.

In other embodiments, substantially purified polypeptides that include a protease domain of a MTSP7 polypeptide or a catalytically active portion thereof, but that do not include the entire sequence of amino acids set forth in SEQ ID No. 18 are provided. Among these are polypeptides that include a sequence of amino acids that has at least 60%, 70%, 80%, 85%, 90%, 95% or 100% sequence identity to SEQ ID No. 16 or 18.

Muteins in which one or more of the Cys residues, particularly, a residue that is paired in the activated two form, but unpaired in the protease domain alone (*i.e.*, the Cys at residue position Cys313 (see SEQ ID Nos. 15-18) in the protease domain), is/are replaced with any amino acid, typically, although not necessarily, a conservative amino acid residue, such as Ser, are contemplated. Muteins of MTSP7, particularly those in which Cys residues, such as the Cys in the single chain protease domain, is replaced with another amino acid that does not eliminate the activity, are provided.

In certain embodiments, the MTSP7 polypeptide is detectable in a body fluid at a level that differs from its level in body fluids in a subject not having a tumor. In other embodiments, the polypeptide is present in a tumor; and a substrate or cofactor for the polypeptide is expressed at levels that differ from its level of expression in a non-tumor cell in the same type of tissue. In other embodiments, the substantially purified the level of expression and/or activity of the MTSP7 polypeptide in tumor cells differs from its level of expression and/or activity in non-tumor cells.

In other embodiments, the MTSP7 is present in a tumor; and a substrate or cofactor for the MTSP7 is expressed at levels that differ from its level of expression in a non-tumor cell in the same type of tissue.

In a specific embodiment, a nucleic acid that encodes a MTSP, designated MTSP7 is provided. In particular, the nucleic acid includes the sequence of nucleotides set forth in SEQ ID No. 15 or 17 or a portion thereof that encodes a catalytically active polypeptide.

-9-

Also provided are nucleic acid molecules that hybridize under conditions of at least low stringency, generally moderate stringency, more typically high stringency to the SEQ ID No. 5 or degenerates thereof.

In one embodiment, the isolated nucleic acid fragment hybridizes to a  
5 nucleic acid molecule containing the nucleotide sequence set forth in SEQ ID No: 15 or 17 (or degenerates thereof) under high stringency conditions, in one embodiment contains the sequence of nucleotides set forth in any of SEQ ID Nos. 15-18. A full-length MTSP7 is set forth in SEQ ID No. 18 and is encoded by SEQ ID No. 17 or degenerates thereof.

10 Also provided are muteins of the single chain protease domain of MTSP7 particularly muteins in which the Cys residue in the protease domain that is free (*i.e.*, does not form disulfide linkages with any other Cys residue in the protease domain) is substituted with another amino acid substitution, typically, although not necessarily, with a conservative amino acid substitution or a substitution  
15 that does not eliminate the activity, and muteins in which a glycosylation site(s) is eliminated. Muteins in which other conservative or non-conservative amino acid substitutions in which catalytic activity is retained are also contemplated (see, *e.g.*, Table 1, for exemplary amino acid substitutions). Hence, provided herein is a the family of transmembrane serine protease (MTSP) proteins  
20 designated MTSP7, and functional domains, especially protease (or catalytic) domains thereof, muteins and other derivatives and analogs thereof. Also provided herein are nucleic acids encoding the MTSP7s.

Additionally provided herein are antibodies that specifically bind to the MTSP7, cells, combinations, kits and articles of manufacture that contain the  
25 nucleic acid encoding the MTSP7 and/or the MTSP7. Further provided herein are prognostic, diagnostic, therapeutic screening methods using MTSP7 and the nucleic acids encoding MTSP7. Also provided are transgenic non-human animals bearing inactivated genes encoding the MTSP and bearing the genes encoding

-10-

the MTSP7 under non-native promotor control are provided. Such animals are useful in animal models of tumor initiation, growth and/or progression models.

Of interest herein are MTSPs that are expressed or are activated in certain tumor or cancer cells such lung, prostate, colon and breast cancers. In particular, it is shown herein, that MTSP7 is expressed in lung carcinoma, leukemia and cervical carcinoma as well as in certain normal cells and tissues (see *e.g.*, EXAMPLES for tissue-specific expression profile). MTSP7 also can be a marker for breast, prostate and colon cancer. The expression or activation of MTSP7 in a cell in a subject can be a marker for breast, prostate, lung, colon and other cancers.

MTSPs are of interest because they appear to be expressed and/or activated at different levels in tumor cells from normal cells, or have functional activity that is different in tumor cells from normal cells, such as by an alteration in a substrate therefor, or a cofactor. MTSP7 is of interest because it is expressed or is active in tumor cells. Hence the MTSP provided herein can serve as diagnostic markers for certain tumors. The level of activated MTSP7 can be diagnostic of cervical or lung cancer or leukemia.

Also provided herein are methods of modulating the activity of the MTSP7 and screening for compounds that modulate, including inhibit, antagonize, agonize or otherwise alter the activity of the MTSP7. Of particular interest is the extracellular domain of MTSP7 that includes the proteolytic (catalytic) portion of the protein.

MTSP7 proteins, including, but not limited including splice variants thereof, and nucleic acids encoding MTSPs, and domains, derivatives and analogs thereof are provided herein. Single chain protease domains that have an N-terminus generated by activation of the zymogen form of MTSP7 are also provided. The cleavage site for the protease domain is at amino acid I<sub>206</sub> (R↓IVQG). The Cys residues at positions C<sub>188</sub>-C<sub>313</sub>, which links protease domain to another domain, C<sub>233</sub>-C<sub>249</sub>, C<sub>358</sub>-C<sub>374</sub> and C<sub>385</sub>-C<sub>413</sub> form disulfide bonds, so that upon cleavage the resulting polypeptide is a two chain molecule. Hence C<sub>313</sub> is a free Cys in the protease domain, which can also be provided as a two chain

-11-

molecule. It is shown herein, however, that the single chain form is proteolytically active.

Antibodies that specifically bind to the MTSP7, particularly the single chain protease domain, the zymogen and activated form, and cells, combinations, kits and articles of manufacture containing the MTSP7 proteins, domains thereof, or encoding nucleic acids are also provided herein. Transgenic non-human animals bearing inactivated genes encoding MTSP7 and bearing the genes encoding the MTSP7, particularly under a non-native promotor control or on an exogenous element, such as a plasmid or artificial chromosome, are additionally provided herein. Also provided are nucleic acid molecules encoding each of MTSP7 and domains thereof.

Also provided are plasmids containing any of the nucleic acid molecules provided herein. Cells containing the plasmids are also provided. Such cells include, but are not limited to, bacterial cells, yeast cells, fungal cells, plant cells, insect cells and animal cells.

Also provided is a method of producing MTSP7 by growing the above-described cells under conditions whereby the MTSP7 is expressed by the cells, and recovering the expressed MTSP7 protein. Methods for isolating nucleic acid encoding other MTSP7s are also provided.

Also provided are cells, generally eukaryotic cells, such as mammalian cells and yeast cells, in which the MTSP7 protein is expressed on the surface of the cells. Such cells are used in drug screening assays to identify compounds that modulate the activity of the MTSP7 protein. These assays including *in vitro* binding assays, and transcription based assays in which signal transduction mediated directly or indirectly, such as via activation of pro-growth factors, by the MTSP7 is assessed.

Further provided herein are prognostic, diagnostic and therapeutic screening methods using the MTSP7 and the nucleic acids encoding MTSP7. In particular, the prognostic, diagnostic and therapeutic screening methods are used for preventing, treating, or for finding agents useful in preventing or treating, tumors or cancers such as lung carcinoma, colon adenocarcinoma and ovarian carcinoma.



-12-

Also provided are methods for screening for compounds that modulate the activity of MTSP7. The compounds are identified by contacting them with the MTSP7 or protease domain thereof and a substrate for the MTSP7. A change in the amount of substrate cleaved in the presence of the compounds compared to that in the absence of the compound indicates that the compound modulates the activity of the MTSP7. Such compounds are selected for further analyses or for use to modulate the activity of the MTSP7, such as inhibitors or agonists. The compounds can also be identified by contacting the substrates with a cell that expresses the MTSP7 or the extracellular domain or proteolytically active portion thereof.

Also provided herein are modulators of the activity of MTSP7, especially the modulators obtained according to the screening methods provide herein. Such modulators can have use in treating cancerous conditions, and other neoplastic conditions.

Pharmaceutical composition containing the protease domain and/or full-length or other domain of an MTSP7 protein are provided herein in a pharmaceutically acceptable carrier or excipient are provided herein.

Also provided are articles of manufacture that contain MTSP7 protein and protease domains of MTSP7 in single chain forms or activated forms. The articles contain a) packaging material; b) the polypeptide (or encoding nucleic acid), particularly the single chain protease domain thereof; and c) a label indicating that the article is for using ins assays for identifying modulators of the activities of an MTSP7 protein is provided herein.

Conjugates containing a) a MTSP7 protein or protease domain in single chain from; and b) a targeting agent linked to the MTSP directly or via a linker, wherein the agent facilitates: i) affinity isolation or purification of the conjugate; ii) attachment of the conjugate to a surface; iii) detection of the conjugate; or iv) targeted delivery to a selected tissue or cell, is provided herein. The conjugate can contain a plurality of agents linked thereto. The conjugate can be a chemical conjugate; and it can be a fusion protein.

In yet another embodiment, the targeting agent is a protein or peptide fragment. The protein or peptide fragment can include a protein binding

-13-

sequence, a nucleic acid binding sequence, a lipid binding sequence, a polysaccharide binding sequence, or a metal binding sequence.

Methods of diagnosing a disease or disorder characterized by detecting an aberrant level of an MTSP7 in a subject is provided. The method can be  
5 practiced by measuring the level of the DNA, RNA, protein or functional activity of the MTSP7. An increase or decrease in the level of the DNA, RNA, protein or functional activity of the MTSP, relative to the level of the DNA, RNA, protein or functional activity found in an analogous sample not having the disease or disorder (or other suitable control) is indicative of the presence of the disease or  
10 disorder in the subject or other relative any other suitable control.

Combinations are provided herein. The combination can include: a) an inhibitor of the activity of an MTSP7; and b) an anti-cancer treatment or agent. The MTSP inhibitor and the anti-cancer agent can be formulated in a single  
15 pharmaceutical composition or each is formulated in a separate pharmaceutical composition. The MTSP7 inhibitor can be an antibody or a fragment or binding portion thereof made against the MTSP7, such as an antibody that specifically binds to the protease domain, an inhibitor of MTSP7 production, or an inhibitor of MTSP7 membrane-localization or an inhibitor of MTSP7 activation. Other MTSP7 inhibitors include, but are not limited to, an antisense nucleic acid or  
20 double-stranded RNA (dsRNA), such as RNAi, encoding the MTSP7, particularly a portion of the protease domain; a nucleic acid encoding at least a portion of a gene encoding the MTSP7 with a heterologous nucleotide sequence inserted therein such that the heterologous sequence inactivates the biological activity encoded MTSP7 or the gene encoding it. For example, the portion of the gene  
25 encoding the MTSP7 can flank the heterologous sequence to promote homologous recombination with a genomic gene encoding the MTSP7.

Also, provided are methods for treating or preventing a tumor or cancer in a mammal by administering to a mammal an effective amount of an inhibitor of an MTSP7, whereby the tumor or cancer is treated or prevented. The MTSP7  
30 inhibitor used in the treatment or for prophylaxis is administered with a pharmaceutically acceptable carrier or excipient. The mammal treated can be a human. The treatment or prevention method can additionally include

-14-

administering an anti-cancer treatment or agent simultaneously with or subsequently or before administration of the MTSP7 inhibitor.

Also provided is a recombinant non-human animal in which an endogenous gene of an MTSP7 has been deleted or inactivated by homologous recombination or insertional mutagenesis of the animal or an ancestor thereof. A recombinant non-human animal is provided herein, where the gene of an MTSP7 is under control of a promoter that is not the native promoter of the gene or that is not the native promoter of the gene in the non-human animal or where the nucleic acid encoding the MTSP7 is heterologous to the non-human animal and the promoter is the native or a non-native promoter or the MTSP7 is on an extrachromosomal element, such as a plasmid or artificial chromosome.

Also provided are methods of treatments of tumors by administering a prodrug that is activated by MTSP7 that is expressed or active in tumor cells, particularly those in which its functional activity in tumor cells is greater than in non-tumor cells. The prodrug is administered and, upon administration, active MTSP7 expressed on cells cleaves the prodrug and releases active drug in the vicinity of these cells. The active anti-cancer drug accumulates in the vicinity of the tumor. This is particularly useful in instances in which MTSP7 is expressed or active in greater quantity, higher level or predominantly in tumor cells compared to other cells.

Also provided are methods of identifying a compound that binds to the single-chain or two-chain form of MTSP7, by contacting a test compound with a both forms; determining to which form the compound binds; and if it binds to a form of MTSP7, further determining whether the compound has at least one of the following properties:

- (i) inhibits activation of the single-chain zymogen form of MTSP7;
- (ii) inhibits activity of the two-chain or single-chain form; and
- (iii) inhibits dimerization of the protein.

The forms can be full length or the protease domain resulting from cleavage at the RI activation site.

Also provided are methods of diagnosing the presence of a pre-malignant lesion, a malignancy, or other pathologic condition in a subject, by obtaining a

-15-

biological sample from the subject; exposing it to a detectable agent that binds to a two-chain or single-chain form of MTSP7, where the pathological condition is characterized by the presence or absence of the two-chain or single-chain form .

5           Methods of inhibiting tumor invasion or metastasis or treating a malignant or pre-malignant condition by administering an agent that inhibits activation of the zymogen form of MTSP7 or an activity of the activated form are provided. The conditions include, but are not limited to, a condition, such as a tumor, of the breast, cervix, prostate, lung, ovary or colon.

10           Antibodies that specifically bind to the two-chain or single-chain form of MTSP7 are provided. The antibodies include those that specifically bind to the two-chain or single-chain form of the protease domain and/or the full-length protein.

#### **BRIEF DESCRIPTION OF THE DRAWING**

15           The Figure illustrates the domain organization of the MTSP7. MTSP7 has a transmembrane domain, a SEA (sea urchin sperm protein-enterokinase-agrin) domain and a serine protease domain; sequences of the full length protein and encoding nucleic acid are also set forth in the figure (see also, SEQ ID Nos. 15-18).

#### **20 DETAILED DESCRIPTION**

##### **A. DEFINITIONS**

          Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of ordinary skill in the art to which this invention belongs. All patents, applications, published  
25   applications and other publications and sequences from GenBank and other data bases referred to herein are incorporated by reference in their entirety.

          As used herein, the abbreviations for any protective groups, amino acids and other compounds, are, unless indicated otherwise, in accord with their common usage, recognized abbreviations, or the IUPAC-IUB Commission on  
30   Biochemical Nomenclature (see, (1972) Biochem. 11:942-944).

          As used herein, serine protease refers to a diverse family of proteases wherein a serine residue is involved in the hydrolysis of proteins or peptides.

-16-

The serine residue can be part of the catalytic triad mechanism, which includes a serine, a histidine and an aspartic acid in the catalysis, or be part of the hydroxyl/ $\epsilon$ -amine or hydroxyl/ $\alpha$ -amine catalytic dyad mechanism, which involves a serine and a lysine in the catalysis.

5           As used herein, "transmembrane serine protease (MTSP)" refers to a family of transmembrane serine proteases that share common structural features as described herein (see, also Hooper *et al.* (2001) *J. Biol. Chem.* 276:857-860). Thus, reference, for example, to "MTSP" encompasses all proteins encoded by the MTSP gene family, including but are not limited to: MTSP1, MTSP3,  
10 MTSP4, MTSP6, MTSP7 or an equivalent molecule obtained from any other source or that has been prepared synthetically or that exhibits the same activity. Other MTSPs include, but are not limited to, corin, enterpeptidase, human airway trypsin-like protease (HAT), MTSP1, TMPRSS2, and TMPRSS4. The term also encompass MTSPs with amino acid substitutions that do not substantially alter  
15 activity of each member, and also encompasses splice variants thereof. Suitable conservative substitutions of amino acids are known to those of skill in this art and can be made generally without altering the enzymatic activity of the resulting molecule or without eliminating. Of particular interest are MTSPs of mammalian, including human, origin. Those of skill in this art recognize that, in  
20 general, single amino acid substitutions in non-essential regions of a polypeptide do not substantially alter biological activity (see, e.g., Watson et al. *Molecular Biology of the Gene*, 4th Edition, 1987, The Bejacmin/Cummings Pub. co., p.224).

MTSPs of interest include those that are activated and/or expressed in  
25 tumor cells at different levels, typically higher, from non-tumor cells; and those from cells in which substrates therefor differ in tumor cells from non-tumor cells or differ with respect to substrates, co-factors or receptors, or otherwise alter the specificity of the MTSP.

As used herein, a "protease domain of an MTSP" refers to the protease  
30 domain of MTSP that is located within the extracellular domain of a MTSP and exhibits proteolytic activity. It includes at least the smallest fragment thereof that acts catalytically as a single chain form. Hence it is at least the minimal

-17-

portion of the extracellular domain that exhibits proteolytic activity as assessed by standard assays *in vitro* assays. Those of skill in this art recognize that such protease domain is the portion of the protease that is structurally equivalent to the trypsin or chymotrypsin fold. Contemplated herein are such protease domains  
5 and catalytically active portions thereof.

The MTSP7 protein, with the protease domains indicated, is illustrated in Figure 1. Smaller portions thereof that retain protease activity are contemplated.

The protease domains from MTSPs vary in size and constitution, including insertions and deletions in surface loops. They retain conserved structure,  
10 including at least one of the active site triad (see, *e.g.*, the catalytic triad of the MTSP in SEQ ID No. 16 is H<sub>248</sub>, D<sub>293</sub>, S<sub>389</sub>), primary specificity pocket, oxyanion hole and/or other features of serine protease domains of proteases. Thus, for purposes herein, the protease domain is a portion of a MTSP, as defined herein, and is homologous to a domain of other MTSPs, such as corin, enterpeptidase,  
15 human airway trypsin-like protease (HAT), MTSP1, TMPRSS2, and TMPRSS4, which have been previously identified; it was not recognized, however, that an isolated single chain form of the protease domain could function proteolytically in *in vitro* assays. As with the larger class of enzymes of the chymotrypsin (S1) fold (see, *e.g.*, Internet accessible MEROPS data base), the MTSPs protease  
20 domains share a high degree of amino acid sequence identity. The His, Asp and Ser residues necessary for activity are present in conserved motifs. The activation site, which results in the N-terminus of second chain in the two chain forms is has a conserved motif and readily can be identified (see, *e.g.*, amino acids 2-6-208).

25 As used herein, the catalytically active domain of an MTSP refers to the protease domain. Reference to the protease domain of an MTSP includes the single and two-chain forms of any of these proteins. The zymogen form of each protein is single chain form, which can be converted to the active two chain form by cleavage. The protease domain can also be converted to a two chain  
30 form. By active form is meant a form active *in vivo* and/or *in vitro*.

Significantly, at least *in vitro*, the single chain forms of the MTSPs and the catalytic domains or proteolytically active portions thereof (typically C-

-18-

terminal truncations) thereof exhibit protease activity. Hence provided herein are isolated single chain forms of the protease domains of MTSPs and their use in *in vitro* drug screening assays for identification of agents that modulate the activity thereof.

5           As used herein an MTSP7, whenever referenced herein, includes at least one or all of or any combination of:

                  a polypeptide encoded by the sequence of nucleotides set forth in SEQ ID No. 15;

                  a polypeptide encoded by a sequence of nucleotides that  
10       hybridizes under conditions of low, moderate or high stringency to the sequence of nucleotides set forth in SEQ ID No. 15;

                  a polypeptide that comprises the sequence of amino acids set forth as amino acids 206-438 of SEQ ID No. 16;

                  a polypeptide that comprises a sequence of amino acids having at  
15       least about 60%, 70%, 80%, 85%, 90% or 95% sequence identity with the sequence of amino acids set forth in SEQ ID No. 16 or 17; and/or

                  a splice variant of the MTSP7 set forth in SEQ ID No. 15.

                  The MTSP7 can be from any animal, particularly a mammal, and includes but are not limited to, humans, rodents, fowl, ruminants and other animals. The  
20       full length zymogen or two-chain activated form is contemplated or any domain thereof, including the protease domain, which can be a two-chain activated form, or a single chain form.

                  As used herein, a human protein is one encoded by DNA present in the genome of a human, including all allelic variants and conservative variations as  
25       long as they are not variants found in other mammals.

                  As used herein, a "nucleic acid encoding a protease domain or catalytically active portion of a MTSP" shall be construed as referring to a nucleic acid encoding only the recited single chain protease domain or active portion thereof, and not the other contiguous portions of the MTSP as a  
30       continuous sequence.

                  As used herein, catalytic activity refers to the activity of the MTSP as a serine proteases. Function of the MTSP refers to its function in tumor biology,

-19-

including promotion of or involvement in tumorigenesis, metastasis or carcinogenesis, and also roles in signal transduction.

As used herein, a zymogen is an inactive precursor of a proteolytic enzyme. Such precursors are generally larger, although not necessarily larger  
5 than the active form. With reference serine proteases zymogens are converted to active enzymes by specific cleavage, including catalytic and autocatalytic cleavage, or binding of an activating co-factor, which generates the mature active enzyme. A zymogen, thus, is an enzymatically inactive protein that is converted to a proteolytic enzyme by the action of an activator.

10 As used herein, "disease or disorder" refers to a pathological condition in an organism resulting from, *e.g.*, infection or genetic defect, and characterized by identifiable symptoms.

As used herein, neoplasm (neoplasia) refers to abnormal new growth, and thus means the same as *tumor*, which can be benign or malignant. Unlike  
15 *hyperplasia*, neoplastic proliferation persists even in the absence of the original stimulus.

As used herein, neoplastic disease refers to any disorder involving cancer, including tumor development, growth, metastasis and progression.

As used herein, cancer refers to a general term for diseases caused by  
20 any type of malignant tumor.

As used herein, malignant, as applies to tumors, refers to primary tumors that have the capacity of *metastasis* with loss of *growth control* and *positional control*.

As used herein, an anti-cancer agent (used interchangeable with "anti-  
25 tumor or anti-neoplastic agent") refers to any agents used in the anti-cancer treatment. These include any agents, when used alone or in combination with other compounds, that can alleviate, reduce, ameliorate, prevent, or place or maintain in a state of remission of clinical symptoms or diagnostic markers associated with neoplastic disease, tumor and cancer, and can be used in  
30 methods, combinations and compositions provided herein. Non-limiting examples of anti-neoplastic agents include anti-angiogenic agents, alkylating agents, antimetabolite, certain natural products, platinum coordination



-20-

complexes, anthracenediones, substituted ureas, methylhydrazine derivatives, adrenocortical suppressants, certain hormones, antagonists and anti-cancer polysaccharides.

As used herein, a splice variant refers to a variant produced by differential  
5 processing of a primary transcript of genomic DNA that results in more than one type of mRNA. Splice variants of MTSPs are provided herein.

As used herein, angiogenesis is intended to broadly encompass the totality of processes directly or indirectly involved in the establishment and maintenance of new vasculature (neovascularization), including, but not limited  
10 to, neovascularization associated with tumors.

As used herein, anti-angiogenic treatment or agent refers to any therapeutic regimen and compound, when used alone or in combination with other treatment or compounds, that can alleviate, reduce, ameliorate, prevent, or place or maintain in a state of remission of clinical symptoms or diagnostic  
15 markers associated with undesired and/or uncontrolled angiogenesis. Thus, for purposes herein an anti-angiogenic agent refers to an agent that inhibits the establishment or maintenance of vasculature. Such agents include, but are not limited to, anti-tumor agents, and agents for treatments of other disorders associated with undesirable angiogenesis, such as diabetic retinopathies,  
20 restenosis, hyperproliferative disorders and others.

As used herein, non-anti-angiogenic anti-tumor agents refer to anti-tumor agents that do not act primarily by inhibiting angiogenesis.

As used herein, pro-angiogenic agents are agents that promote the establishment or maintenance of the vasculature. Such agents include agents  
25 for treating cardiovascular disorders, including heart attacks and strokes.

As used herein, undesired and/or uncontrolled angiogenesis refers to pathological angiogenesis wherein the influence of angiogenesis stimulators outweighs the influence of angiogenesis inhibitors. As used herein, deficient angiogenesis refers to pathological angiogenesis associated with disorders where  
30 there is a defect in normal angiogenesis resulting in aberrant angiogenesis or an absence or substantial reduction in angiogenesis.

-21-

As used herein, the protease domain of an MTSP protein refers to the protease domain of an MTSP that is located within or is the extracellular domain of an MTSP and exhibits proteolytic activity. Hence it is at least the minimal portion of the extracellular domain that exhibits proteolytic activity as assessed  
5 by standard assays *in vitro*. It refers, herein, to a single chain form heretofore thought to be inactive.

Exemplary protease domains include at least a sufficient portion of sequences of amino acids set forth as amino acids 206-438 in SEQ ID No. 16 (encoded by nucleotides in SEQ ID No. 15; see also the Figure). Also contemplated are  
10 nucleic acid molecules that encode polypeptide that has proteolytic activity in an *in vitro* proteolysis assay and that have at least 60%, 70%, 80%, 85%, 90% or 95% sequence identity with the full length thereof of a protease domain of an MTSP7 protein, or that hybridize along their full length or along at least about 70%, 80% or 90% of the full length, to a nucleic acid molecule that encodes a  
15 protease domain, particularly under conditions of moderate, generally high, stringency.

For the protease domains, residues at the N-terminus can be critical for activity. It is shown herein that the protease domain of the single chain form of the MTSP7 protease is catalytically active. Hence the protease domain will  
20 require the N-terminal amino acids; the c-terminus portion can be truncated. The amount that can be removed can be determined empirically by testing the protein for protease activity in an *in vitro* assays that assesses catalytic cleavage.

Hence smaller portions of the protease domains, particularly the single chain domains, thereof that retain protease activity are contemplated. Such  
25 smaller versions will generally be C-terminal truncated versions of the protease domains. The protease domains vary in size and constitution, including insertions and deletions in surface loops. Such domains exhibit conserved structure, including at least one structural feature, such as the active site triad, primary specificity pocket, oxyanion hole and/or other features of serine protease  
30 domains of proteases. Thus, for purposes herein, the protease domain is a single chain portion of an MTSP7, as defined herein, but is homologous in its structural features and retention of sequence of similarity or homology the

-22-

protease domain of chymotrypsin or trypsin. Most significantly, the polypeptide will exhibit proteolytic activity as a single chain.

As used herein, by homologous means about greater than or equal to 25% nucleic acid sequence identity, typically 25%, 40%, 60%, 70%, 80%, 85%, 90% or 95%; the precise percentage can be specified if necessary. For purposes herein the terms "homology" and "identity" are often used interchangeably. In general, for determination of the percentage identity, sequences are aligned so that the highest order match is obtained (see, e.g.: *Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data, Part I*, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; Carillo *et al.* (1988) *SIAM J Applied Math* 48:1073). By sequence identity, the number of conserved amino acids are determined by standard alignment algorithms programs, and are used with default gap penalties established by each supplier. Substantially homologous nucleic acid molecules would hybridize typically at moderate stringency or at high stringency all along the length of the nucleic acid of interest. Also contemplated are nucleic acid molecules that contain degenerate codons in place of codons in the hybridizing nucleic acid molecule.

Whether any two nucleic acid molecules have nucleotide sequences that are at least 60%, 70%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% "identical" can be determined using known computer algorithms such as the "FAST A" program, using for example, the default parameters as in Pearson *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:2444 (other programs include the GCG program package (Devereux, J., *et al.*, *Nucleic Acids Research* 12(1):387 (1984)), BLASTP, BLASTN, FASTA (Atschul, S.F., *et al.*, *J Molec Biol* 215:403 (1990); Guide to Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and Carillo *et al.* (1988) *SIAM J Applied Math* 48:1073). For example, the BLAST function of the National Center for Biotechnology

-23-

Information database can be used to determine identity. Other commercially or publicly available programs include, DNASTar "MegAlign" program (Madison, WI) and the University of Wisconsin Genetics Computer Group (UWG) "Gap" program (Madison WI)). Percent homology or identity of proteins and/or nucleic acid molecules can be determined, for example, by comparing sequence information using a GAP computer program (*e.g.*, Needleman *et al.* (1970) *J. Mol. Biol.* 48:443, as revised by Smith and Waterman ((1981) *Adv. Appl. Math.* 2:482). Briefly, the GAP program defines similarity as the number of aligned symbols (*i.e.*, nucleotides or amino acids) which are similar, divided by the total number of symbols in the shorter of the two sequences. Default parameters for the GAP program can include: (1) a unary comparison matrix (containing a value of 1 for identities and 0 for non-identities) and the weighted comparison matrix of Gribskov *et al.* (1986) *Nucl. Acids Res.* 14:6745, as described by Schwartz and Dayhoff, eds., *ATLAS OF PROTEIN SEQUENCE AND STRUCTURE*, National Biomedical Research Foundation, pp. 353-358 (1979); (2) a penalty of 3.0 for each gap and an additional 0.10 penalty for each symbol in each gap; and (3) no penalty for end gaps.

Therefore, as used herein, the term "identity" represents a comparison between a test and a reference polypeptide or polynucleotide. As used herein, the term at least "90% identical to" refers to percent identities from 90 to 99.99 relative to the reference polypeptides. Identity at a level of 90% or more is indicative of the fact that, assuming for exemplification purposes a test and reference polynucleotide length of 100 amino acids are compared. No more than 10% (*i.e.*, 10 out of 100) amino acids in the test polypeptide differs from that of the reference polypeptides. Similar comparisons can be made between a test and reference polynucleotides. Such differences can be represented as point mutations randomly distributed over the entire length of an amino acid sequence or they can be clustered in one or more locations of varying length up to the maximum allowable, *e.g.* 10/100 amino acid difference (approximately 90% identity). Differences are defined as nucleic acid or amino acid substitutions, or deletions. At the level of homologies or identities above about 85-90%, the result should be independent of the program and gap parameters

-24-

set; such high levels of identity can be assessed readily, often without relying on software.

As used herein, primer refers to an oligonucleotide containing two or more deoxyribonucleotides or ribonucleotides, generally more than three, from which synthesis of a primer extension product can be initiated. Experimental  
5 conditions conducive to synthesis include the presence of nucleoside triphosphates and an agent for polymerization and extension, such as DNA polymerase, and a suitable buffer, temperature and pH.

As used herein, animals include any animal, such as, but are not limited  
10 to, goats, cows, deer, sheep, rodents, pigs and humans. Non-human animals, exclude humans as the contemplated animal. The MTSPs provided herein are from any source, animal, plant, prokaryotic and fungal, such as MTSP7s are of animal origin, including mammalian origin.

As used herein, genetic therapy involves the transfer of heterologous  
15 DNA to the certain cells, target cells, of a mammal, particularly a human, with a disorder or conditions for which such therapy is sought. The DNA is introduced into the selected target cells in a manner such that the heterologous DNA is expressed and a therapeutic product encoded thereby is produced.

Alternatively, the heterologous DNA can in some manner mediate expression of  
20 DNA that encodes the therapeutic product, or it can encode a product, such as a peptide or RNA that in some manner mediates, directly or indirectly, expression of a therapeutic product. Genetic therapy can also be used to deliver nucleic acid encoding a gene product that replaces a defective gene or supplements a gene product produced by the mammal or the cell in which it is introduced. The  
25 introduced nucleic acid can encode a therapeutic compound, such as a growth factor inhibitor thereof, or a tumor necrosis factor or inhibitor thereof, such as a receptor therefor, that is not normally produced in the mammalian host or that is not produced in therapeutically effective amounts or at a therapeutically useful time. The heterologous DNA encoding the therapeutic product can be modified  
30 prior to introduction into the cells of the afflicted host in order to enhance or otherwise alter the product or expression thereof. Genetic therapy can also

-25-

involve delivery of an inhibitor or repressor or other modulator of gene expression.

As used herein, heterologous DNA is DNA that encodes RNA and proteins that are not normally produced *in vivo* by the cell in which it is expressed or that  
5 mediates or encodes mediators that alter expression of endogenous DNA by affecting transcription, translation, or other regulatable biochemical processes. Heterologous DNA can also be referred to as foreign DNA. Any DNA that one of skill in the art would recognize or consider as heterologous or foreign to the cell in which is expressed is herein encompassed by heterologous DNA. Examples of  
10 heterologous DNA include, but are not limited to, DNA that encodes traceable marker proteins, such as a protein that confers drug resistance, DNA that encodes therapeutically effective substances, such as anti-cancer agents, enzymes and hormones, and DNA that encodes other types of proteins, such as antibodies. Antibodies that are encoded by heterologous DNA can be secreted  
15 or expressed on the surface of the cell in which the heterologous DNA has been introduced.

Hence, herein heterologous DNA or foreign DNA, includes a DNA molecule not present in the exact orientation and position as the counterpart DNA molecule found in the genome. It can also refer to a DNA molecule from  
20 another organism or species (*i.e.*, exogenous).

As used herein, a therapeutically effective product is a product that is encoded by heterologous nucleic acid, typically DNA, that, upon introduction of the nucleic acid into a host, a product is expressed that ameliorates or eliminates the symptoms, manifestations of an inherited or acquired disease or that cures  
25 the disease.

As used herein, recitation that a polypeptide consists essentially of the protease domain means that the only MTSP portion of the polypeptide is a protease domain or a catalytically active portion thereof. The polypeptide can optionally, and generally will, include additional non-MTSP-derived sequences of  
30 amino acids.

As used herein, cancer or tumor treatment or agent refers to any therapeutic regimen and/or compound that, when used alone or in combination

-26-

with other treatments or compounds, can alleviate, reduce, ameliorate, prevent, or place or maintain in a state of remission of clinical symptoms or diagnostic markers associated with deficient angiogenesis.

As used herein, domain refers to a portion of a molecule, e.g., proteins  
5 or the encoding nucleic acids, that is structurally and/or functionally distinct from other portions of the molecule.

As used herein, protease refers to an enzyme catalyzing hydrolysis of proteins or peptides. It includes the zymogen form and activated forms thereof. For clarity reference to protease refers to all forms, and particular forms will be  
10 specifically designated. For purposes herein, the protease domain includes single and two chain forms of the protease domain of MTSP7.

As used herein, catalytic activity refers to the activity of the MTSP as a protease as assessed in *in vitro* proteolytic assays that detect proteolysis of a selected substrate.

As used herein, nucleic acids include DNA, RNA, dsRNA and analog  
15 thereof, including protein nucleic acids (PNA) and mixture thereof. Nucleic acids can be single or double stranded. When referring to probes or primers, optionally labeled, with a detectable label, such as a fluorescent or radiolabel, single-stranded molecules are contemplated. Such molecules are typically of a length  
20 such that target is statistically unique or of low copy number (typically less than 5, generally less than 3) for probing or priming a library. Generally a probe or primer contains at least 14, 16 or 30 contiguous of sequence complementary to or identical a gene of interest. Probes and primers can be 10, 20, 30, 50, 100 or more nucleic acids long.

As used herein, nucleic acid encoding a fragment or portion of an MTSP  
25 refers to a nucleic acid encoding only the recited fragment or portion of MTSP, and not the other contiguous portions of the MTSP.

As used herein, heterologous or foreign DNA and RNA are used interchangeably and refer to DNA or RNA that does not occur naturally as part of  
30 the genome in which it is present or which is found in a location or locations in the genome that differ from that in which it occurs in nature. Heterologous nucleic acid is generally not endogenous to the cell into which it is introduced,

-27-

but has been obtained from another cell or prepared synthetically. Generally, although not necessarily, such nucleic acid encodes RNA and proteins that are not normally produced by the cell in which it is expressed. Any DNA or RNA that one of skill in the art would recognize or consider as heterologous or foreign  
5 to the cell in which it is expressed is herein encompassed by heterologous DNA. Heterologous DNA and RNA can also encode RNA or proteins that mediate or alter expression of endogenous DNA by affecting transcription, translation, or other regulatable biochemical processes.

As used herein, operative linkage of heterologous DNA to regulatory and  
10 effector sequences of nucleotides, such as promoters, enhancers, transcriptional and translational stop sites, and other signal sequences refers to the relationship between such DNA and such sequences of nucleotides. For example, operative linkage of heterologous DNA to a promoter refers to the physical relationship between the DNA and the promoter such that the transcription of such DNA is  
15 initiated from the promoter by an RNA polymerase that specifically recognizes, binds to and transcribes the DNA in reading frame. Thus, as used herein, operatively linked or operationally associated refers to the functional relationship of DNA with regulatory and effector sequences of nucleotides, such as promoters, enhancers, transcriptional and translational stop sites, and other  
20 signal sequences. For example, operative linkage of DNA to a promoter refers to the physical and functional relationship between the DNA and the promoter such that the transcription of such DNA is initiated from the promoter by an RNA polymerase that specifically recognizes, binds to and transcribes the DNA. In order to optimize expression and/or *in vitro* transcription, it can be necessary to  
25 remove, add or alter 5' untranslated portions of the clones to eliminate extra, potential inappropriate alternative translation initiation (*i.e.*, start) codons or other sequences that can interfere with or reduce expression, either at the level of transcription or translation. Alternatively, consensus ribosome binding sites (see, e.g., Kozak *J. Biol. Chem.* 266:19867-19870 (1991)) can be inserted  
30 immediately 5' of the start codon and can enhance expression. The desirability of (or need for) such modification can be empirically determined.



-28-

As used herein, a sequence complementary to at least a portion of an RNA, with reference to antisense oligonucleotides, means a sequence having sufficient complementarity to be able to hybridize with the RNA, generally under moderate or high stringency conditions, forming a stable duplex; in the case of  
5 double-stranded MTSP antisense nucleic acids, a single strand of the duplex DNA can thus be tested, or triplex formation can be assayed. The ability to hybridize depends on the degree of complementarity and the length of the antisense nucleic acid. Generally, the longer the hybridizing nucleic acid, the more base mismatches with a MTSP encoding RNA it can contain and still form  
10 a stable duplex (or triplex, as the case can be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

For purposes herein, amino acid substitutions can be made in any of MTSPs and protease domains thereof provided that the resulting protein exhibits  
15 protease activity. Conservative amino acid substitutions, such as those set forth in Table 1, are those that do not eliminate proteolytic activity. Suitable conservative substitutions of amino acids are known to those of skill in this art and can be made generally without altering the biological activity of the resulting molecule. Those of skill in this art recognize that, in general, single amino acid  
20 substitutions in non-essential regions of a polypeptide do not substantially alter biological activity (see, e.g., Watson et al. *Molecular Biology of the Gene*, 4th Edition, 1987, The Benjamin/Cummings Pub. co., p.224). Also included within the definition, is the catalytically active fragment of an MTSP, particularly a single chain protease portion. Conservative amino acid substitutions are made,  
25 for example, in accordance with those set forth in TABLE 1 as follows:

-29-

TABLE 1

	Original residue	Conservative substitution
	Ala (A)	Gly; Ser, Abu
	Arg (R)	Lys, orn
5	Asn (N)	Gln; His
	Cys (C)	Ser
	Gln (Q)	Asn
	Glu (E)	Asp
	Gly (G)	Ala; Pro
10	His (H)	Asn; Gln
	Ile (I)	Leu; Val; Met; Nle; Nva
	Leu (L)	Ile; Val; Met; Nle; Nv
	Lys (K)	Arg; Gln; Glu
	Met (M)	Leu; Tyr; Ile; NLe Val
15	Ornithine	Lys; Arg
	Phe (F)	Met; Leu; Tyr
	Ser (S)	Thr
	Thr (T)	Ser
	Trp (W)	Tyr
20	Tyr (Y)	Trp; Phe
	Val (V)	Ile; Leu; Met; Nle; Nv

Other substitutions are also permissible and can be determined empirically or in accord with known conservative substitutions.

As used herein, Abu is 2-aminobutyric acid; Orn is ornithine.

25 As used herein, the amino acids, which occur in the various amino acid sequences appearing herein, are identified according to their well-known, three-letter or one-letter abbreviations. The nucleotides, which occur in the various DNA fragments, are designated with the standard single-letter designations used routinely in the art.

30 As used herein, a splice variant refers to a variant produced by differential processing of a primary transcript of genomic DNA that results in more than one type of mRNA.

As used herein, a probe or primer based on a nucleotide sequence disclosed herein, includes at least 10, 14, generally at least 16 or 30 or 100  
 35 contiguous sequence of nucleotides of SEQ ID No. 15, except for a region that includes the sequence that encodes amino acids 117-171 and 185-354 of SEQ ID Nos 15 and 16.

-30-

As used herein, amelioration of the symptoms of a particular disorder by administration of a particular pharmaceutical composition refers to any lessening, whether permanent or temporary, lasting or transient that can be attributed to or associated with administration of the composition.

5       As used herein, antisense polynucleotides refer to synthetic sequences of nucleotide bases complementary to mRNA or the sense strand of double stranded DNA. Admixture of sense and antisense polynucleotides under appropriate conditions leads to the binding of the two molecules, or hybridization. When these polynucleotides bind to (hybridize with) mRNA,  
10 inhibition of protein synthesis (translation) occurs. When these polynucleotides bind to double-stranded DNA, inhibition of RNA synthesis (transcription) occurs. The resulting inhibition of translation and/or transcription leads to an inhibition of the synthesis of the protein encoded by the sense strand. Antisense nucleic acid molecule typically contain a sufficient number of nucleotides to specifically  
15 bind to a target nucleic acid, generally at least 5 contiguous nucleotides, often at least 14 or 16 or 30 contiguous nucleotides or modified nucleotides complementary to the coding portion of a nucleic acid molecule that encodes a gene of interest, for example, nucleic acid encoding a single chain protease domain of an MTSP.

20       As used herein, an array refers to a collection of elements, such as antibodies, containing three or more members. An addressable array is one in which the members of the array are identifiable, typically by position on a solid phase support. Hence, in general the members of the array will be immobilized to discrete identifiable loci on the surface of a solid phase.

25       As used herein, antibody refers to an immunoglobulin, whether natural or partially or wholly synthetically produced, including any derivative thereof that retains the specific binding ability the antibody. Hence antibody includes any protein having a binding domain that is homologous or substantially homologous to an immunoglobulin binding domain. Antibodies include members of any  
30 immunoglobulin claims, including IgG, IgM, IgA, IgD and IgE.

As used herein, antibody fragment refers to any derivative of an antibody that is less than full length, retaining at least a portion of the full-length

-31-

antibody's specific binding ability. Examples of antibody fragments include, but are not limited to, Fab, Fab', F(ab)<sub>2</sub>, single-chain Fvs (scFV), Fv, dsFV diabody and Fd fragments. The fragment can include multiple chains linked together, such as by disulfide bridges. An antibody fragment generally contains at least  
5 about 50 amino acids and typically at least 200 amino acids.

As used herein, an Fv antibody fragment is composed of one variable heavy domain (V<sub>H</sub>) and one variable light domain linked by noncovalent interactions.

As used herein, a dsFV refers to an Fv with an engineered intermolecular  
10 disulfide bond, which stabilizes the V<sub>H</sub>-V<sub>L</sub> pair.

As used herein, an F(ab)<sub>2</sub> fragment is an antibody fragment that results from digestion of an immunoglobulin with pepsin at pH 4.0-4.5; it can be recombinantly produce the equivalent fragment.

As used herein, Fab fragments is an antibody fragment that results from  
15 digestion of an immunoglobulin with papain; it can be recombinantly produce the equivalent fragment.

As used herein, scFVs refer to antibody fragments that contain a variable light chain (V<sub>L</sub>) and variable heavy chain (V<sub>H</sub>) covalently connected by a polypeptide linker in any order. The linker is of a length such that the two  
20 variable domains are bridged without substantial interference. Exemplary linkers include (Gly-Ser)<sub>n</sub> residues with some Glu or Lys residues dispersed throughout to increase solubility.

As used herein, humanized antibodies refer to antibodies that are modified to include human sequences of amino acids so that administration to a  
25 human will not provoke an immune response. Methods for preparation of such antibodies are known. For example, the hybridoma that expresses the monoclonal antibody is altered by recombinant DNA techniques to express an antibody in which the amino acid composition of the non-variable regions is based on human antibodies. Computer programs have been designed to identify  
30 such regions.

As used herein, diabodies are dimeric scFV; diabodies typically have shorter peptide linkers than scFvs, and they typically dimerize.

-32-

As used herein the term assessing is intended to include quantitative and qualitative determination in the sense of obtaining an absolute value for the activity of an MTSP, or a domain thereof, present in the sample, and also of obtaining an index, ratio, percentage, visual or other value indicative of the level of the activity. Assessment can be direct or indirect and the chemical species actually detected need not of course be the proteolysis product itself but can for example be a derivative thereof or some further substance.

As used herein, biological activity refers to the in vivo activities of a compound or physiological responses that result upon in vivo administration of a compound, composition or other mixture. Biological activity, thus, encompasses therapeutic effects and pharmaceutical activity of such compounds, compositions and mixtures. Biological activities can be observed in in vitro systems designed to test or use such activities. Thus, for purposes herein the biological activity of a luciferase is its oxygenase activity whereby, upon oxidation of a substrate, light is produced.

As used herein, a combination refers to any association between two or among more items.

As used herein, a composition refers to a any mixture. It can be a solution, a suspension, liquid, powder, a paste, aqueous, non-aqueous or any combination thereof.

As used herein, a conjugate refers to the compounds provided herein that include one or more MTSPs, including an MTSP7, particularly single chain protease domains thereof, and one or more targeting agents. These conjugates include those produced by recombinant means as fusion proteins, those produced by chemical means, such as by chemical coupling, through, for example, coupling to sulfhydryl groups, and those produced by any other method whereby at least one MTSP, or a domain thereof, is linked, directly or indirectly via linker(s) to a targeting agent.

As used herein, a targeting agent, is any moiety, such as a protein or effective portion thereof, that provides specific binding of the conjugate to a cell surface receptor, which, in certain embodiments internalizes the conjugate or

-33-

MTSP portion thereof. A targeting agent can also be one that promotes or facilitates, for example, affinity isolation or purification of the conjugate; attachment of the conjugate to a surface; or detection of the conjugate or complexes containing the conjugate.

5           As used herein, an antibody conjugate refers to a conjugate in which the targeting agent is an antibody.

          As used herein, derivative or analog of a molecule refers to a portion derived from or a modified version of the molecule.

          As used herein, fluid refers to any composition that can flow. Fluids thus  
10 encompass compositions that are in the form of semi-solids, pastes, solutions, aqueous mixtures, gels, lotions, creams and other such compositions.

          As used herein, an effective amount of a compound for treating a particular disease is an amount that is sufficient to ameliorate, or in some manner reduce the symptoms associated with the disease. Such amount can be  
15 administered as a single dosage or can be administered according to a regimen, whereby it is effective. The amount can cure the disease but, typically, is administered in order to ameliorate the symptoms of the disease. Repeated administration can be required to achieve the desired amelioration of symptoms.

          As used herein equivalent, when referring to two sequences of nucleic  
20 acids means that the two sequences in question encode the same sequence of amino acids or equivalent proteins. When equivalent is used in referring to two proteins or peptides, it means that the two proteins or peptides have substantially the same amino acid sequence with only conservative amino acid substitutions (see, *e.g.*, Table 1, above) that do not substantially alter the  
25 activity or function of the protein or peptide. When equivalent refers to a property, the property does not need to be present to the same extent [*e.g.*, two peptides can exhibit different rates of the same type of enzymatic activity], but the activities are generally substantially the same. Complementary, when referring to two nucleotide sequences, means that the two sequences of  
30 nucleotides are capable of hybridizing, generally with less than 25%, less than 15%, and even less than 5%, including with no mismatches between opposed

-34-

nucleotides. Generally the two molecules will hybridize under conditions of high stringency.

As used herein, an agent that modulates the activity of a protein or expression of a gene or nucleic acid either decreases or increases or otherwise  
5 alters the activity of the protein or, in some manner up- or down-regulates or otherwise alters expression of the nucleic acid in a cell.

As used herein, inhibitor of an activity of an MTSP encompasses any substances that prohibit or decrease production, post-translational  
modification(s), maturation, or membrane localization of the MTSP or any  
10 substances that interfere with or decrease the proteolytic efficacy of thereof, particularly of a single chain form *in vitro* for screening assay.

As used herein, a method for treating or preventing neoplastic disease means that any of the symptoms, such as the tumor, metastasis thereof, the vascularization of the tumors or other parameters by which the disease is  
15 characterized are reduced, ameliorated, prevented, placed in a state of remission, or maintained in a state of remission. It also means that the hallmarks of neoplastic disease and metastasis can be eliminated, reduced or prevented by the treatment. Non-limiting examples of the hallmarks include uncontrolled degradation of the basement membrane and proximal extracellular matrix,  
20 migration, division, and organization of the endothelial cells into new functioning capillaries, and the persistence of such functioning capillaries.

As used herein, pharmaceutically acceptable salts, esters or other derivatives of the conjugates include any salts, esters or derivatives that can be readily prepared by those of skill in this art using known methods for such  
25 derivatization and that produce compounds that can be administered to animals or humans without substantial toxic effects and that either are pharmaceutically active or are prodrugs.

As used herein, a prodrug is a compound that, upon in vivo administration, is metabolized or otherwise converted to the biologically,  
30 pharmaceutically or therapeutically active form of the compound. To produce a prodrug, the pharmaceutically active compound is modified such that the active compound will be regenerated by metabolic processes. The prodrug can be

-35-

designed to alter the metabolic stability or the transport characteristics of a drug, to mask side effects or toxicity, to improve the flavor of a drug or to alter other characteristics or properties of a drug. By virtue of knowledge of pharmacodynamic processes and drug metabolism in vivo, those of skill in this art, once a pharmaceutically active compound is known, can design prodrugs of the compound (see, e.g., Nogrady (1985) Medicinal Chemistry A Biochemical Approach, Oxford University Press, New York, pages 388-392).

As used herein, a drug identified by the screening methods provided herein refers to any compound that is a candidate for use as a therapeutic or as lead compound for design of a therapeutic. Such compounds can be small molecules, including small organic molecules, peptides, peptide mimetics, antisense molecules or dsRNA, such as RNAi, antibodies, fragments of antibodies, recombinant antibodies and other such compound which can serve as drug candidate or lead compound.

As used herein, a peptidomimetic is a compound that mimics the conformation and certain stereochemical features of the biologically active form of a particular peptide. In general, peptidomimetics are designed to mimic certain desirable properties of a compound, but not the undesirable properties, such as flexibility, that lead to a loss of a biologically active conformation and bond breakdown. Peptidomimetics may be prepared from biologically active compounds by replacing certain groups or bonds that contribute to the undesirable properties with bioisosteres. Bioisosteres are known to those of skill in the art. For example the methylene bioisostere  $\text{CH}_2\text{S}$  has been used as an amide replacement in enkephalin analogs (see, e.g., Spatola (1983) pp. 267-357 in Chemistry and Biochemistry of Amino Acids, Peptides, and Proteins, Weistein, Ed. volume 7, Marcel Dekker, New York). Morphine, which can be administered orally, is a compound that is a peptidomimetic of the peptide endorphin. For purposes herein, cyclic peptides are included among peptidomimetics.

As used herein, production by recombinant means by using recombinant DNA methods means the use of the well known methods of molecular biology for expressing proteins encoded by cloned DNA.



-36-

As used herein, a promoter region or promoter element refers to a segment of DNA or RNA that controls transcription of the DNA or RNA to which it is operatively linked. The promoter region includes specific sequences that are sufficient for RNA polymerase recognition, binding and transcription initiation.

- 5 This portion of the promoter region is often referred to as the promoter. In addition, the promoter region includes sequences that modulate this recognition, binding and transcription initiation activity of RNA polymerase. These sequences can be *cis* acting or can be responsive to *trans* acting factors. Promoters, depending upon the nature of the regulation, can be constitutive or regulated.
- 10 Exemplary promoters contemplated for use in prokaryotes include the bacteriophage T7 and T3 promoters.

- As used herein, a receptor refers to a molecule that has an affinity for a given ligand. Receptors can be naturally-occurring or synthetic molecules. Receptors can also be referred to in the art as anti-ligands. As used herein, the
- 15 receptor and anti-ligand are interchangeable. Receptors can be used in their unaltered state or as aggregates with other species. Receptors can be attached, covalently or noncovalently, or in physical contact with, to a binding member, either directly or indirectly via a specific binding substance or linker. Examples of receptors, include, but are not limited to: antibodies, cell membrane receptors
- 20 surface receptors and internalizing receptors, monoclonal antibodies and antisera reactive with specific antigenic determinants (such as on viruses, cells, or other materials), drugs, polynucleotides, nucleic acids, peptides, cofactors, lectins, sugars, polysaccharides, cells, cellular membranes, and organelles.

- Examples of receptors and applications using such receptors, include but
- 25 are not restricted to:

- a) enzymes: specific transport proteins or enzymes essential to survival of microorganisms, which could serve as targets for antibiotic (ligand) selection;
  - b) antibodies: identification of a ligand-binding site on the antibody molecule that combines with the epitope of an antigen of interest can be
- 30 investigated; determination of a sequence that mimics an antigenic epitope can lead to the development of vaccines of which the immunogen is based on one or

-37-

more of such sequences or lead to the development of related diagnostic agents or compounds useful in therapeutic treatments such as for auto-immune diseases

c) nucleic acids: identification of ligand, such as protein or RNA, binding sites;

5 d) catalytic polypeptides: polymers, generally polypeptides, that are capable of promoting a chemical reaction involving the conversion of one or more reactants to one or more products; such polypeptides generally include a binding site specific for at least one reactant or reaction intermediate and an active functionality proximate to the binding site, in which the functionality is  
10 capable of chemically modifying the bound reactant (see, e.g., U.S. Patent No. 5,215,899);

e) hormone receptors: determination of the ligands that bind with high affinity to a receptor is useful in the development of hormone replacement therapies; for example, identification of ligands that bind to such receptors can  
15 lead to the development of drugs to control blood pressure; and

f) opiate receptors: determination of ligands that bind to the opiate receptors in the brain is useful in the development of less-addictive replacements for morphine and related drugs.

As used herein, sample refers to anything which can contain an analyte  
20 for which an analyte assay is desired. The sample can be a biological sample, such as a biological fluid or a biological tissue. Examples of biological fluids include urine, blood, plasma, serum, saliva, semen, stool, sputum, cerebral spinal fluid, tears, mucus, amniotic fluid or the like. Biological tissues are aggregate of cells, usually of a particular kind together with their intercellular substance that  
25 form one of the structural materials of a human, animal, plant, bacterial, fungal or viral structure, including connective, epithelium, muscle and nerve tissues. Examples of biological tissues also include organs, tumors, lymph nodes, arteries and individual cell(s).

As used herein: stringency of hybridization in determining percentage  
30 mismatch is as follows:

1) high stringency: 0.1 x SSPE, 0.1% SDS, 65°C

2) medium stringency: 0.2 x SSPE, 0.1% SDS, 50°C

-38-

3) low stringency: 1.0 x SSPE, 0.1% SDS, 50°C

Those of skill in this art know that the washing step selects for stable hybrids and also know the ingredients of SSPE (see, e.g., Sambrook, E.F. Fritsch, T. Maniatis, in: Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory Press (1989), vol. 3, p. B.13, see, also, numerous catalogs that describe commonly used laboratory solutions). SSPE is pH 7.4 phosphate-buffered 0.18 M NaCl. Further, those of skill in the art recognize that the stability of hybrids is determined by  $T_m$ , which is a function of the sodium ion concentration and temperature ( $T_m = 81.5^\circ \text{C} - 16.6(\log_{10}[\text{Na}^+]) + 0.41(\% \text{G} + \text{C}) - 600/l$ ), so that the only parameters in the wash conditions critical to hybrid stability are sodium ion concentration in the SSPE (or SSC) and temperature.

It is understood that equivalent stringencies can be achieved using alternative buffers, salts and temperatures. By way of example and not limitation, procedures using conditions of low stringency are as follows (see also Shilo and Weinberg, *Proc. Natl. Acad. Sci. USA*, 78:6789-6792 (1981)): Filters containing DNA are pretreated for 6 hours at 40°C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.1% PVP, 0.1% Ficoll, 1% BSA, and 500 µg/ml denatured salmon sperm DNA (10X SSC is 1.5 M sodium chloride, and 0.15 M sodium citrate, adjusted to a pH of 7).

Hybridizations are carried out in the same solution with the following modifications: 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 µg/ml salmon sperm DNA, 10% (wt/vol) dextran sulfate, and  $5-20 \times 10^6$  cpm  $^{32}\text{P}$ -labeled probe is used. Filters are incubated in hybridization mixture for 18-20 hours at 40°C, and then washed for 1.5 hours at 55°C in a solution containing 2X SSC, 25 mM Tris-HCl (pH 7.4), 5 mM EDTA, and 0.1% SDS. The wash solution is replaced with fresh solution and incubated an additional 1.5 hours at 60°C. Filters are blotted dry and exposed for autoradiography. If necessary, filters are washed for a third time at 65-68°C and reexposed to film. Other conditions of low stringency which can be used are well known in the art (*e.g.*, as employed for cross-species hybridizations).

By way of example and not way of limitation, procedures using conditions of moderate stringency is provided. For example, but not limited to,

-39-

procedures using such conditions of moderate stringency are as follows: Filters containing DNA are pretreated for 6 hours at 55°C in a solution containing 6X SSC, 5X Denhart's solution, 0.5% SDS and 100 µg/ml denatured salmon sperm DNA. Hybridizations are carried out in the same solution and 5-20 X 10<sup>6</sup> cpm <sup>32</sup>P-labeled probe is used. Filters are incubated in hybridization mixture for 18-20 hours at 55°C, and then washed twice for 30 minutes at 60°C in a solution containing 1X SSC and 0.1% SDS. Filters are blotted dry and exposed for autoradiography. Other conditions of moderate stringency which can be used are well-known in the art. Washing of filters is done at 37°C for 1 hour in a solution containing 2X SSC, 0.1% SDS.

By way of example and not way of limitation, procedures using conditions of high stringency are as follows: Prehybridization of filters containing DNA is carried out for 8 hours to overnight at 65°C in buffer composed of 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 µg/ml denatured salmon sperm DNA. Filters are hybridized for 48 hours at 65°C in prehybridization mixture containing 100 µg/ml denatured salmon sperm DNA and 5-20 X 10<sup>6</sup> cpm of <sup>32</sup>P-labeled probe. Washing of filters is done at 37°C for 1 hour in a solution containing 2X SSC, 0.01% PVP, 0.01% Ficoll, and 0.01% BSA. This is followed by a wash in 0.1X SSC at 50°C for 45 minutes before autoradiography. Other conditions of high stringency which can be used are well known in the art.

The term substantially identical or homologous or similar varies with the context as understood by those skilled in the relevant art and generally means at least 70%, at least 80%, at least 90%, and generally at least 95% identity. The percentage will be apparent from the context or can be specified.

As used herein, substantially identical to a product means sufficiently similar so that the property of interest is sufficiently unchanged so that the substantially identical product can be used in place of the product.

As used herein, substantially pure means sufficiently homogeneous to appear free of readily detectable impurities as determined by standard methods of analysis, such as thin layer chromatography (TLC), gel electrophoresis and high performance liquid chromatography (HPLC), used by those of skill in the art

-40-

to assess such purity, or sufficiently pure such that further purification would not detectably alter the physical and chemical properties, such as enzymatic and biological activities, of the substance. Methods for purification of the compounds to produce substantially chemically pure compounds are known to those of skill in the art. A substantially chemically pure compound can, however, be a mixture of stereoisomers or isomers. In such instances, further purification might increase the specific activity of the compound.

As used herein, target cell refers to a cell that expresses an MTSP *in vivo*.

As used herein, test substance refers to a chemically defined compound (e.g., organic molecules, inorganic molecules, organic/inorganic molecules, proteins, peptides, nucleic acids, oligonucleotides, lipids, polysaccharides, saccharides, or hybrids among these molecules such as glycoproteins) or mixtures of compounds (e.g., a library of test compounds, natural extracts or culture supernatants) whose effect on an MTSP, particularly a single chain form that includes the protease domain or a sufficient portion thereof for activity, as determined by *in vitro* method, such as the assays provided herein.

As used herein, the terms a therapeutic agent, therapeutic regimen, radioprotectant, chemotherapeutic mean conventional drugs and drug therapies, including vaccines, which are known to those skilled in the art. Radiotherapeutic agents are well known in the art.

As used herein, treatment means any manner in which the symptoms of a conditions, disorder or disease are ameliorated or otherwise beneficially altered. Treatment also encompasses any pharmaceutical use of the compositions herein.

As used herein, vector (or plasmid) refers to discrete elements that are used to introduce heterologous DNA into cells for either expression or replication thereof. The vectors typically remain episomal, but can be designed to effect integration of a gene or portion thereof into a chromosome of the genome. Also contemplated are vectors that are artificial chromosomes, such as yeast artificial chromosomes and mammalian artificial chromosomes. Selection and use of such vehicles are well known to those of skill in the art. An expression vector includes vectors capable of expressing DNA that is operatively linked with regulatory sequences, such as promoter regions, that are capable of effecting

-41-

expression of such DNA fragments. Thus, an expression vector refers to a recombinant DNA or RNA construct, such as a plasmid, a phage, recombinant virus or other vector that, upon introduction into an appropriate host cell, results in expression of the cloned DNA. Appropriate expression vectors are well  
5 known to those of skill in the art and include those that are replicable in eukaryotic cells and/or prokaryotic cells and those that remain episomal or those which integrate into the host cell genome.

As used herein, protein binding sequence refers to a protein or peptide sequence that is capable of specific binding to other protein or peptide  
10 sequences generally, to a set of protein or peptide sequences or to a particular protein or peptide sequence.

As used herein, epitope tag refers to a short stretch of amino acid residues corresponding to an epitope to facilitate subsequent biochemical and immunological analysis of the epitope tagged protein or peptide. Epitope tagging  
15 is achieved by including the sequence of the epitope tag to the protein-encoding sequence in an appropriate expression vector. Epitope tagged proteins can be affinity purified using highly specific antibodies raised against the tags.

As used herein, metal binding sequence refers to a protein or peptide sequence that is capable of specific binding to metal ions generally, to a set of  
20 metal ions or to a particular metal ion.

As used herein, a cellular extract refers to a preparation or fraction which is made from a lysed or disrupted cell.

As used herein, an agent is said to be randomly selected when the agent is chosen randomly without considering the specific sequences involved in the  
25 association of a protein alone or with its associated substrates and binding partners. An example of randomly selected agents is the use a chemical library or a peptide combinatorial library, or a growth broth of an organism or condition medium.

As used herein, an agent is said to rationally selected or designed when  
30 the agent is chosen on a non-random basis which takes into account the sequence of the target site and/or its conformation in connection with the agent's action. As described in the Examples, there are proposed binding sites

-42-

for serine protease and (catalytic) sites in the protein having SEQ ID NO:3 or SEQ ID NO:4. Agents can be rationally selected or rationally designed by utilizing the peptide sequences that make up these sites. For example, a rationally selected peptide agent can be a peptide whose amino acid sequence is  
5 identical to the ATP or calmodulin binding sites or domains.

For clarity of disclosure, and not by way of limitation, the detailed description is divided into the subsections that follow.

**B. MTSP7 PROTEINS, MUTEINS, DERIVATIVES AND ANALOGS THEREOF**  
**MTSPs**

10 The MTSPs are a family of transmembrane serine proteases that are found in mammals and also other species that share a number of common structural features including: a proteolytic extracellular C-terminal domain; a transmembrane domain, with a hydrophobic domain near the N-terminus; a short cytoplasmic domain; and a variable length stem region containing modular  
15 domains. The proteolytic domains share sequence homology including conserved his, asp, and ser residues necessary for catalytic activity that are present in conserved motifs. The MTSPs are synthesized as zymogens, and activated to two chain forms by cleavage. It is shown herein that the single chain proteolytic domain can function *in vitro* and, hence is useful in *in vitro*  
20 assays for identifying agents that modulate the activity of members of this family.

The MTSP family is a target for therapeutic intervention and also some members can serve as diagnostic markers for tumor development, growth and/or progression. As discussed, the members of this family are involved in  
25 proteolytic processes that are implicated in tumor development, growth and/or progression. This implication is based upon their functions as proteolytic enzymes in processes related to ECM degradative pathways. In addition, their levels of expression or level of activation or their apparent activity resulting from substrate levels or alterations in substrates and levels thereof differs in tumor  
30 cells and non-tumor cells in the same tissue. Hence, protocols and treatments that alter their activity, such as their proteolytic activities and roles in signal transduction, and/or their expression, such as by contacting them with a

-43-

compound that modulates their activity and/or expression, could impact tumor development, growth and/or progression. Also, in some instances, the level of activation and/or expression can be altered in tumors, such as lung carcinoma, colon adenocarcinoma and ovarian carcinoma.

- 5 MTSPs are of interest because they appear to be expressed and/or activated at different levels in tumor cells from normal cells, or have functional activity that is different in tumor cells from normal cells, such as by an alteration in a substrate therefor, or a cofactor.

#### MTSP7

- 10 Isolated, substantially pure proteases that include protease domains or a catalytically active portion thereof in single chain form of MTSPs also are provided. Provided is the family member designated MTSP7. The protease domains can be included in a longer protein, and such longer protein is optionally the MTSP7 zymogen. MTSP7 is of interest because it is expressed or is active  
15 in tumor cells. Hence the MTSP provided herein can serve as diagnostic markers for certain tumors. The level of activated MTSP7 can be diagnostic of cancers, including cervical or lung cancer or leukemia.

- It is shown herein, that MTSP7s provided herein are expressed and/or activated in certain tumors; hence their activation or expression can serve as a  
20 diagnostic marker for tumor development, growth and/or progression. In other instances the MTSP protein can exhibit altered activity by virtue of a change in activity or expression of a co-factor therefor or a substrate therefor. In addition, in some instances, these MTSPS and/or variants thereof can be shed from cell surfaces. Detection of the shed MTSPS, particularly the extracellular domains, in  
25 body fluids, such as serum, blood, saliva, cerebral spinal fluid, synovial fluid and interstitial fluids, urine, sweat and other such fluids and secretions, can serve as a diagnostic tumor marker. In particular, detection of higher levels of such shed polypeptides in a subject compared to a subject known not to have any neoplastic disease or compared to earlier samples from the same subject, can be  
30 indicative of neoplastic disease in the subject.

The protease domains of an MTSP are single-chain polypeptides, with an N-terminus (such as IV, VV, IL and II) generated at the cleavage site (generally



-44-

having the consensus sequence R↓VVGG, R↓IVGG, R↓IVQ, R↓IVNG, R↓ILGG, R↓VGLL, R↓ILGG or a variation thereof; an N-terminus R↓V or R↓I, where the arrow represents the cleavage point) when the zymogen is activated. The protease domain of MTSP7, produced by cleavage (R↓I) includes the sequence  
5 IVNG, is set forth in SEQ ID Nos. 17 and 18.

The protease domain of the MTSP does not have to result from activation, which produces a two chain activated product, but rather includes single chain polypeptides with the N-terminus include the consensus sequence ↓VVGG, ↓IVGG, ↓VGLL, ↓ILGG, ↓IVQG or ↓IVNG or other such motif at the N-terminus. Such polypeptides, although not the result of activation and not two-chain forms, exhibit proteolytic (catalytic) activity. These protease domain polypeptides are used in assays to screen for agents that modulate the activity of the MTSP7.  
10

The domains, fragments, derivatives or analogs of an MTSP7 that are functionally active are capable of exhibiting one or more functional activities associated with the MTSP7 protein, such as serine protease activity, immunogenicity and antigenicity, are provided.  
15

#### **Polypeptides and muteins**

Provided herein are isolated substantially pure single polypeptides that contain the protease domain of an MTSP7 as a single chain. The protein can also include other non-MTSP sequences of amino acids, but will include the protease domain or a sufficient portion thereof to exhibit catalytic activity in any *in vitro* assay that assess such protease activity, such as any provided herein.  
20

MTSP7 polypeptides provided herein are expressed on or active in or on tumor cells, typically at a level that differs from the level in which they are expressed or active in the non-tumor cell of the same type. Hence, for example, if the MTSP is expressed in an ovarian tumor cell, it is expressed or active at a different level in non-tumor ovarian cells. MTSP protease domains include the single chain protease domains of MTSP7.  
25

Also provided herein are nucleic acid molecules that encode MTSP proteins and the encoded proteins. In particular, nucleic acid molecules encoding MTSP7 from animals, including splice variants thereof are provided.  
30

-45-

The encoded proteins are also provided. Also provided are functional domains thereof.

In specific aspects, the MTSP protease domains, portions thereof, and muteins thereof are from or based on animal MTSPs, including, but are not limited to, rodent, such as mouse and rat; fowl, such as chicken; ruminants, such as goats, cows, deer, sheep; ovine, such as pigs; and humans. In particular, MTSP7 derivatives can be made by altering their sequences by substitutions, additions or deletions. Due to the degeneracy of nucleotide coding sequences, other nucleic sequences which encode substantially the same amino acid sequence as a MTSP7 gene can be used. These include but are not limited to nucleotide sequences comprising all or portions of MTSP7 genes that are altered by the substitution of different codons that encode the amino acid residue within the sequence, thus producing a silent change. Likewise, the MTSP7 derivatives include, but are not limited to, those containing, as a primary amino acid sequence, all or part of the amino acid sequence of MTSP7, including altered sequences in which functionally equivalent amino acid residues are substituted for residues within the sequence resulting in a silent change. For example, one or more amino acid residues within the sequence can be substituted by another amino acid of a similar polarity which acts as a functional equivalent, resulting in a silent alteration. Substitutes for an amino acid within the sequence can be selected from other members of the class to which the amino acid belongs. For example, the nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan and methionine. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine. The positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid (see, *e.g.*, Table 1). Muteins of the MTSP7 or a domain thereof, such as a protease domain, in which up to about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 85%, 90% or 95% of the amino acids are replaced with another amino acid are provided. Generally such muteins retain at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80% or 90% of the protease activity the unmutated protein.

-46-

Also provided are peptides that are encoded by the nucleic acid molecules described herein. Included among those polypeptides are the MTSP7 protease domain or a polypeptide with amino acid changes such that the specificity and protease activity remains substantially unchanged or changed (increased or  
5 decreased) by a specified percentage, such as 10, 20, 30, 40, 50%. In particular, a substantially purified mammalian MTSP protein is provided that has a transmembrane domain and can additionally include a transmembrane (TM) domain, a SEA domain and a serine protease catalytic domain is provided.

Also provided is a substantially purified protein containing a sequence of  
10 amino acids that has at least 60%, at least about 80%, at least about 90% or at least about 95%, identity to the MTSP7 where the percentage identity is determined using standard algorithms and gap penalties that maximize the percentage identity. The human MTSP7 protein is included, although other mammalian MTSP7 proteins are contemplated. The precise percentage of  
15 identity can be specified if needed.

Predicted disulfide bonds pairing in MTSP7 is Cys233 to Cys249; Cys358 to Cys374; Cys385 to Cys413 and Cys186 to Cys313. The Cys313 is in the protease domain and is unpaired in the single chain form of the protease domain.

Mutins of MTSP7, particularly those in which Cys residues, such as the  
20 Cys313 in the single chain protease domain, is replaced with another amino acid, such as Ser, Gly or Ala, that does not eliminate the activity, are provided.

Also provided are substantially purified MTSP7 polypeptides and functional domains thereof, including catalytically active domains and portions, that have at least about 60%, 70%, 80%, 85%, 90% or 95% sequence identity  
25 with a protease domain that includes the sequence of amino acids set forth in SEQ ID No. 16 or a catalytically active portion thereof or with a protease that includes the sequence of amino acids set forth in SEQ ID No. 18 and domains thereof.

In particular, exemplary protease domains include at least a sufficient  
30 portion of sequences of amino acids set forth as amino acids 206-438 in SEQ ID No. 16 (encoded by nucleotides in SEQ ID No. 15 and 17; see also the Figure). Also contemplated are nucleic acid molecules that encode a single chain MTSP

-47-

protease that have proteolytic activity in an *in vitro* proteolysis assay and that have at least 60%, 70%, 80%, 85%, 90% or 95% sequence identity with the full length of a protease domain of an MTSP7 protein, or that hybridize along their full length to a nucleic acids that encode a protease domain, particularly  
5 under conditions of moderate, generally high, stringency. As above, the encoded polypeptides contain the protease as a single chain.

Muteins of the protein are also provided in which amino acids are replaced with other amino acids. Among the muteins are those in which the Cys residues, is/are replaced typically with a conservative amino acid residues, such  
10 as a serine. Such muteins are also provided herein. Muteins in which 10%, 20%, 30%, 35%, 40%, 45%, 50% or more of the amino acids are replaced but the resulting polypeptide retains at least about 10%, 20%, 30%, 35%, 40%, 45%, 50%, 60%, 70%, 80%, 90% or 95% of the catalytic activity as the unmodified form for the same substrate.

15 Muteins can be made by making conservative amino acid substitutions and also non-conservative amino acid substitutions. For example, amino acid substitutions the desirably alter properties of the proteins can be made. In one embodiment, mutations that prevent degradation of the polypeptide can be made. Many proteases cleave after basic residues, such as R and K; to eliminate  
20 such cleavage, the basic residue is replaced with a non-basic residue. Interaction of the protease with an inhibitor can be blocked while retaining catalytic activity by effecting a non-conservative change at the site interaction of the inhibitor with the protease. Receptor binding can be altered without altering catalytic activity.

## 25 Nucleic acids

In one embodiment, the substantially purified MTSP protease is encoded by a nucleic acid that hybridizes to the a nucleic acid molecule containing the protease domain encoded by the nucleotide sequence set forth in any of  
30 SEQ. ID Nos. 15 and 17 under at least moderate, generally high, stringency conditions, such that the protease domain encoding nucleic acid thereof hybridizes along its full length or at least 70%, 80% or 90% of the full length. In certain embodiments the substantially purified MTSP protease is a single

-48-

chain polypeptide that includes substantially the sequence of amino acids set forth in SEQ ID No. 2, 16 or the protease domain portion thereof, or a catalytically active portion thereof. Figure 1 depicts the structural organization of the MTSP7.

- 5           In a specific embodiment, a nucleic acid that encodes a MTSP, designated MTSP7 is provided. In particular, the nucleic acid includes an open reading frame within the following sequence of nucleotides set forth in SEQ ID No. 15.           In particular the protein is encoded by the open reading frame that begins at nucleotide 48 (check and ends at 1360). Also provided are nucleic acid
- 10           molecules that hybridize under conditions of at least low stringency, moderate stringency, and generally high stringency to the following sequence of nucleic acids (SEQ ID No. 15), particularly to the open reading frame encompassed by nucleotides that encode a single protease domain thereof, or any domain of MTSP7.
- 15           Also included are substantially purified MTSP7 zymogen, activated two chain forms, single chain protease domains and two chain protease domains. These are encoded by a nucleic acid that includes sequence encoding a protease domain that exhibits proteolytic activity and that hybridizes to a nucleic acid molecule having a nucleotide sequence set forth in SEQ ID No. 15, typically
- 20           under moderate, generally under high stringency, conditions and generally along the full length (or substantially the full length) of the protease domain. Splice variants are also contemplated herein.

- In certain embodiments, the isolated nucleic acid fragment hybridizes to the nucleic acid having the nucleotide sequence set forth in SEQ ID No: 15 (or
- 25           the molecules in the figure in the FIGURE) under high stringency conditions, and generally contains the sequence of nucleotides set forth in SEQ ID Nos. 15-17; see also the Figure). The protein contains a transmembrane domain (TM), a SEA domain and a serine protease domain. Muteins of the protein are also provided in which amino acids are replaced with conservative amino acids. Among the
- 30           muteins are those in which the Cys residues, is/are replaced with generally conservative amino acid residues, such as a serine. Such muteins are also provided herein. Each of such domains is provided herein as are nucleic acid

-49-

molecules that include sequences of nucleotides encoding such domains. Some MTSPs can additionally include a LDLR domain, a scavenger-receptor cysteine rich (SRCR) domain and other domains.

The isolated nucleic acid fragment is DNA, including genomic or cDNA, or  
5 is RNA, or can include other components, such as protein nucleic acid. The isolated nucleic acid can include additional components, such as heterologous or native promoters, and other transcriptional and translational regulatory sequences, these genes can be linked to other genes, such as reporter genes or other indicator genes or genes that encode indicators.

10 Also provided is an isolated nucleic acid molecule that includes the sequence of molecules that is complementary to the nucleotide sequence encoding MTSP7 or the portion thereof.

Also provided are fragments thereof that can be used as probes or primers and that contain at least about 10 nucleotides, 14 nucleotides, generally  
15 at least about 16 nucleotides, often at least about 30 nucleotides. The length of the probe or primer is a function of the size of the genome probed; the larger the genome, the longer the probe or primer required for specific hybridization to a single site. Those of skill in the art can select appropriately sized probes and primers. Generally probes and primers as described are single-stranded. Double  
20 stranded probes and primers can be used, if they are denatured when used.

Also provided are nucleic acid molecules that hybridize to the above-noted sequences of nucleotides encoding MTSP7 at least at low stringency, moderate stringency, and typically at high stringency, and that encode the protease domain and/or the full length protein or at least 70%, 80% or 90% of  
25 the full length or other domains of an MTSP7 or a splice variant or allelic variant thereof. Generally the molecules hybridize under such conditions along their full length or at least 70%, 80% or 90% of the full length for at least one domain and encode at least one domain, such as the protease or extracellular domain, of the polypeptide. In particular, such nucleic acid molecules include any isolated  
30 nucleic fragment that encodes at least one domain of a membrane serine protease, that (1) contains a sequence of nucleotides that encodes the protease or a domain thereof, and (2) is selected from among:

-50-

- (a) a sequence of nucleotides that encodes the protease or a domain thereof includes a sequence of nucleotides set forth above;
- (b) a sequence of nucleotides that encodes such portion or the full length protease and hybridizes under conditions of high stringency, generally to nucleic acid that is complementary to a mRNA transcript present in a mammalian cell that encodes such protein or fragment thereof;
- (c) a sequence of nucleotides that encodes a transmembrane protease or domain thereof that includes a sequence of amino acids encoded by such portion or the full length open reading frame; and
- (d) a sequence of nucleotides that encodes the transmembrane protease that includes a sequence of amino acids encoded by a sequence of nucleotides that encodes such subunit and hybridizes under conditions of low, moderate or high stringency to DNA that is complementary to the mRNA transcript.

The isolated nucleic acids can include of at least 8 nucleotides of an MTSP7-encoding sequence. In other embodiments, the nucleic acids can contain least 25 (continuous) nucleotides, 50 nucleotides, 100 nucleotides, 150 nucleotides, or 200 nucleotides of an MTSP7-encoding sequence, or a full-length MTSP coding sequence. In another embodiment, the nucleic acids are smaller than 35, 200 or 500 nucleotides in length. Nucleic acids can be single or double stranded. Nucleic acids that hybridizes to or complementary to the foregoing sequences, in particular the inverse complement to nucleic acids that hybridizes to the foregoing sequences (*i.e.*, the inverse complement of a nucleic acid strand has the complementary sequence running in reverse orientation to the strand so that the inverse complement would hybridize without mismatches to the nucleic acid strand; thus, for example, where the coding strand hybridizes to a nucleic acid with no mismatches between the coding strand and the hybridizing strand, then the inverse complement of the hybridizing strand is identical to the coding strand) are also provided.

In specific aspects, nucleic acids are provided that include a sequence complementary to (specifically are the inverse complement of) at least 10, 25,

-51-

50, 100, or 200 nucleotides or the entire coding region of an MTSP7 encoding nucleic acid, particularly the protease domain thereof. For MTSP7 the full-length protein or a domain or active fragment thereof is also provided.

For each of the nucleic acid molecules, the nucleic acid can be DNA or  
5 RNA or PNA or other nucleic acid analogs or can include non-natural nucleotide bases. Also provided are isolated nucleic acid molecules that include a sequence of nucleotides complementary to the nucleotide sequence encoding an MTSP.

Probes and primers derived from the nucleic acid molecules are provided,  
Such probes and primers contain at least 8, 14, 16, 30, 100 or more contiguous  
10 nucleotides with identity to contiguous nucleotides of an MTSP7, generally except for nucleic acids encoding 117-171 and 185-354 of SEQ ID No. 15. The probes and primers are optionally labelled with a detectable label, such as a radiolabel or a fluorescent tag, or can be mass differentiated for detection by mass spectrometry or other means.

#### 15 **Plasmids and Cells**

Plasmids and vectors containing the nucleic acid molecules are also provided. Cells containing the vectors, including cells that express the encoded proteins are provided. The cell can be a bacterial cell, a yeast cell, a fungal cell, a plant cell, an insect cell or an animal cell. Methods for producing an MTSP or  
20 single chain form of the protease domain thereof by, for example, growing the cell under conditions whereby the encoded MTSP is expressed by the cell, and recovering the expressed protein, are provided herein. As noted, for MTSP7, the full-length zymogens and activated proteins and activated (two chain) protease and single chain protease domains are provided.

25 As described herein, the cells are used for expression of the protein, which can be secreted or expressed in the cytoplasm.

The above discussion provides an overview and some details of the exemplified MTSP7s. The following discussion provides additional details (see, also, EXAMPLES).



-52-

**C. Tumor specificity and tissue expression profiles**

Each MTSP has a characteristic tissue expression profile; the MTSPs in particular, although not exclusively expressed or activated in tumors, exhibit characteristic tumor tissue expression or activation profiles. In some instances, MTSPs can have different activity in a tumor cell from a non-tumor cell by virtue of a change in a substrate or cofactor therefor or other factor that would alter the apparent functional activity of the MTSP. Hence each can serve as a diagnostic marker for particular tumors, by virtue of a level of activity and/or expression or function in a subject (i.e. a mammal, particularly a human) with neoplastic disease, compared to a subject or subjects that do not have the neoplastic disease. In addition, detection of activity (and/or expression) in a particular tissue can be indicative of neoplastic disease. Shed MTSPs in body fluids can be indicative of neoplastic disease. Also, by virtue of the activity and/or expression profiles of each, they can serve as therapeutic targets, such as by administration of modulators of the activity thereof, or, as by administration of a prodrug specifically activated by one of the MTSPs.

**Tissue expression profiles****MTSP7**

The MTSP7 transcript was detected in lung carcinoma (A549 cell line), leukemia (K-562 cell line) and cervical carcinoma (HeLaS3 cell line).

**D. Identification and isolation of MTSP protein genes**

The MTSP proteins, or domains thereof, can be obtained by methods well known in the art for protein purification and recombinant protein expression. Any method known to those of skill in the art for identification of nucleic acids that encode desired genes can be used. Any method available in the art can be used to obtain a full length (i.e., encompassing the entire coding region) cDNA or genomic DNA clone encoding an MTSP protein. In particular, the polymerase chain reaction (PCR) can be used to amplify a sequence identified as being differentially expressed in normal and tumor cells or tissues, e.g., nucleic acids encoding an MTSP7 protein (SEQ. NOs: 15-17), in a genomic or cDNA library. Oligonucleotide primers that hybridize to sequences at the 3' and 5' termini of the identified sequences can be used as primers to amplify by PCR sequences

-53-

from a nucleic acid sample (RNA or DNA), generally a cDNA library, from an appropriate source (e.g., tumor or cancer tissue).

PCR can be carried out, e.g., by use of a Perkin-Elmer Cetus thermal cycler and Taq polymerase (Gene Amp<sup>™</sup>). The DNA being amplified can include  
5 mRNA or cDNA or genomic DNA from any eukaryotic species. One can choose to synthesize several different degenerate primers, for use in the PCR reactions. It is also possible to vary the stringency of hybridization conditions used in priming the PCR reactions, to amplify nucleic acid homologs (e.g., to obtain *MTSP protein* sequences from species other than humans or to obtain human  
10 sequences with homology to MTSP7 protein) by allowing for greater or lesser degrees of nucleotide sequence similarity between the known nucleotide sequence and the nucleic acid homolog being isolated. For cross species hybridization, low stringency to moderate conditions are used. For same species hybridization, moderately stringent to highly stringent conditions are used. The  
15 conditions can be empirically determined.

After successful amplification of the nucleic acid containing all or a portion of the identified MTSP protein sequence or of a nucleic acid encoding all or a portion of an MTSP protein homolog, that segment can be molecularly cloned and sequenced, and used as a probe to isolate a complete cDNA or  
20 genomic clone. This, in turn, permits the determination of the gene's complete nucleotide sequence, the analysis of its expression, and the production of its protein product for functional analysis. Once the nucleotide sequence is determined, an open reading frame encoding the MTSP protein gene protein product can be determined by any method well known in the art for determining  
25 open reading frames, for example, using publicly available computer programs for nucleotide sequence analysis. Once an open reading frame is defined, it is routine to determine the amino acid sequence of the protein encoded by the open reading frame. In this way, the nucleotide sequences of the entire MTSP protein genes as well as the amino acid sequences of MTSP protein proteins and  
30 analogs can be identified.

Any eukaryotic cell potentially can serve as the nucleic acid source for the molecular cloning of the MTSP protein gene. The nucleic acids can be

-54-

isolated from vertebrate, mammalian, human, porcine, bovine, feline, avian, equine, canine, as well as additional primate sources, insects, plants and other organisms. The DNA can be obtained by standard procedures known in the art from cloned DNA (*e.g.*, a DNA "library"), by chemical synthesis, by cDNA  
5 cloning, or by the cloning of genomic DNA, or fragments thereof, purified from the desired cell (see, for example, Sambrook et al., 1989, *Molecular Cloning, A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York; Glover, D.M. (ed.), 1985, *DNA Cloning: A Practical Approach*, MRL Press, Ltd., Oxford, U.K. Vol. I, II). Clones derived from  
10 genomic DNA can contain regulatory and intron DNA regions in addition to coding regions; clones derived from cDNA will contain only exon sequences. Whatever the source, the gene should be molecularly cloned into a suitable vector for propagation of the gene.

In the molecular cloning of the gene from genomic DNA, DNA fragments  
15 are generated, some of which will encode the desired gene. The DNA can be cleaved at specific sites using various restriction enzymes. Alternatively, one can use DNase in the presence of manganese to fragment the DNA, or the DNA can be physically sheared, for example, by sonication. The linear DNA fragments then can be separated according to size by standard techniques,  
20 including but not limited to, agarose and polyacrylamide gel electrophoresis and column chromatography.

Once the DNA fragments are generated, identification of the specific DNA fragment containing the desired gene can be accomplished in a number of ways. For example, a portion of the MTSP protein (of any species) gene (*e.g.*, a PCR  
25 amplification product obtained as described above or an oligonucleotide having a sequence of a portion of the known nucleotide sequence) or its specific RNA, or a fragment thereof be purified and labeled, and the generated DNA fragments can be screened by nucleic acid hybridization to the labeled probe (Benton and Davis, *Science* 196:180 (1977); Grunstein and Hogness, *Proc. Natl. Acad. Sci. U.S.A.* 72:3961 (1975)). Those DNA fragments with substantial homology to  
30 the probe will hybridize. It is also possible to identify the appropriate fragment by restriction enzyme digestion(s) and comparison of fragment sizes with those

-55-

expected according to a known restriction map if such is available or by DNA sequence analysis and comparison to the known nucleotide sequence of MTSP protein. Further selection can be carried out on the basis of the properties of the gene. Alternatively, the presence of the gene can be detected by assays based  
5 on the physical, chemical, or immunological properties of its expressed product. For example, cDNA clones, or DNA clones which hybrid-select the proper mRNA, can be selected which produce a protein that, *e.g.*, has similar or identical electrophoretic migration, isoelectric focusing behavior, proteolytic digestion maps, antigenic properties, serine protease activity. If an anti-MTSP  
10 protein antibody is available, the protein can be identified by binding of labeled antibody to the putatively MTSP protein synthesizing clones, in an ELISA (enzyme-linked immunosorbent assay)-type procedure.

Alternatives to isolating the MTSP7 protein genomic DNA include, but are not limited to, chemically synthesizing the gene sequence from a known  
15 sequence or making cDNA to the mRNA that encodes the MTSP protein. For example, RNA for cDNA cloning of the MTSP protein gene can be isolated from cells expressing the protein. The identified and isolated nucleic acids then can be inserted into an appropriate cloning vector. A large number of vector-host systems known in the art can be used. Possible vectors include, but are not  
20 limited to, plasmids or modified viruses, but the vector system must be compatible with the host cell used. Such vectors include, but are not limited to, bacteriophages such as lambda derivatives, or plasmids such as pBR322 or pUC plasmid derivatives or the Bluescript vector (Stratagene, La Jolla, CA). The insertion into a cloning vector can, for example, be accomplished by ligating the  
25 DNA fragment into a cloning vector which has complementary cohesive termini. If the complementary restriction sites used to fragment the DNA are not present in the cloning vector, the ends of the DNA molecules can be enzymatically modified. Alternatively, any site desired can be produced by ligating nucleotide sequences (linkers) onto the DNA termini; these ligated linkers can comprise  
30 specific chemically synthesized oligonucleotides encoding restriction endonuclease recognition sequences. In an alternative method, the cleaved vector and MTSP protein gene can be modified by homopolymeric tailing.

-56-

Recombinant molecules can be introduced into host cells via transformation, transfection, infection, electroporation, calcium precipitation and other methods, so that many copies of the gene sequence are generated.

In an alternative method, the desired gene can be identified and isolated  
5 after insertion into a suitable cloning vector in a "shot gun" approach. Enrichment for the desired gene, for example, by size fractionization, can be done before insertion into the cloning vector.

In specific embodiments, transformation of host cells with recombinant DNA molecules that incorporate the isolated MTSP protein gene, cDNA, or  
10 synthesized DNA sequence enables generation of multiple copies of the gene. Thus, the gene can be obtained in large quantities by growing transformants, isolating the recombinant DNA molecules from the transformants and, when necessary, retrieving the inserted gene from the isolated recombinant DNA.

**E. Vectors, plasmids and cells that contain nucleic acids encoding an MTSP  
15 protein or protease domain thereof and expression of MTSP proteins**

#### **Vectors and cells**

For recombinant expression of one or more of the MTSP proteins, the nucleic acid containing all or a portion of the nucleotide sequence encoding the MTSP protein can be inserted into an appropriate expression vector, *i.e.*, a  
20 vector that contains the necessary elements for the transcription and translation of the inserted protein coding sequence. The necessary transcriptional and translational signals can also be supplied by the native promoter for MTSP genes, and/or their flanking regions.

Also provided are vectors that contain nucleic acid encoding the MTSPs.  
25 Cells containing the vectors are also provided. The cells include eukaryotic and prokaryotic cells, and the vectors are any suitable for use therein.

Prokaryotic and eukaryotic cells, including endothelial cells, containing the vectors are provided. Such cells include bacterial cells, yeast cells, fungal cells, plant cells, insect cells and animal cells. The cells are used to produce an MTSP  
30 protein or protease domain thereof by growing the above-described cells under conditions whereby the encoded MTSP protein or protease domain of the MTSP protein is expressed by the cell, and recovering the expressed protease domain

-57-

protein. In the exemplified embodiments, the protease domain is secreted into the medium.

In one embodiment, the vectors include a sequence of nucleotides that encodes a polypeptide that has protease activity and contains all or a portion of only the protease domain, or multiple copies thereof, of an MTSP protein are provided. Also provided are vectors that comprise a sequence of nucleotides that encodes the protease domain and additional portions of an MTSP protein up to and including a full length MTSP protein, as well as multiple copies thereof, are also provided. The vectors can be selected for expression of the MTSP protein or protease domain thereof in the cell or such that the MTSP protein is expressed as a transmembrane protein. Alternatively, the vectors can include signals necessary for secretion of encoded proteins. When the protease domain is expressed the nucleic acid can be linked to a nucleic acid sequence encoding a secretion signal, such as the *Saccharomyces cerevisiae*  $\alpha$  mating factor signal sequence or a portion thereof sufficient for secretion. Any such signal sequence can be used.

A variety of host-vector systems can be used to express the protein coding sequence. These include but are not limited to mammalian cell systems infected with virus (*e.g.* vaccinia virus, adenovirus, herpes virus, and other virus-derived vectors); insect cell systems infected with virus (*e.g.* baculovirus); microorganisms such as yeast containing yeast vectors; or bacteria transformed with bacteriophage, DNA, plasmid DNA, or cosmid DNA. The expression elements of vectors vary in their strengths and specificities. Depending on the host-vector system used, any one of a number of suitable transcription and translation elements can be used.

Any methods known to those of skill in the art for the insertion of DNA fragments into a vector can be used to construct expression vectors containing a chimeric gene containing of appropriate transcriptional/translational control signals and protein coding sequences. These methods can include *in vitro* recombinant DNA and synthetic techniques and *in vivo* recombinants (genetic recombination). Expression of nucleic acid sequences encoding MTSP protein, or domains, derivatives, fragments or homologs thereof, can be regulated by a

-58-

second nucleic acid sequence so that the genes or fragments thereof are expressed in a host transformed with the recombinant DNA molecule(s). For example, expression of the proteins can be controlled by any promoter/enhancer known in the art. In a specific embodiment, the promoter is not native to the genes for MTSP protein. Promoters which can be used include but are not limited to the SV40 early promoter (Bernoist and Chambon, *Nature* 290:304-310 (1981)), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., *Cell* 22:787-797 (1980)), the herpes thymidine kinase promoter (Wagner et al., *Proc. Natl. Acad. Sci. USA* 78:1441-1445 (1981)), the regulatory sequences of the metallothionein gene (Brinster et al., *Nature* 296:39-42 (1982)); prokaryotic expression vectors such as the  $\beta$ -lactamase promoter (Villa-Kamaroff et al., *Proc. Natl. Acad. Sci. USA* 75:3727-3731 (1978)) or the *tac* promoter (DeBoer et al., *Proc. Natl. Acad. Sci. USA* 80:21-25 (1983)); see also "Useful Proteins from Recombinant Bacteria": in Scientific American 242:79-94 (1980)); plant expression vectors containing the nopaline synthetase promoter (Herrar-Estrella et al., *Nature* 303:209-213 (1984)) or the cauliflower mosaic virus 35S RNA promoter (Gardner et al., *Nucleic Acids Res.* 9:2871 (1981)), and the promoter of the photosynthetic enzyme ribulose biphosphate carboxylase (Herrera-Estrella et al., *Nature* 310:115-120 (1984)); promoter elements from yeast and other fungi such as the Gal4 promoter, the alcohol dehydrogenase promoter, the phosphoglycerol kinase promoter, the alkaline phosphatase promoter, and the following animal transcriptional control regions that exhibit tissue specificity and have been used in transgenic animals: elastase I gene control region which is active in pancreatic acinar cells (Swift et al., *Cell* 38:639-646 (1984); Ornitz et al., *Cold Spring Harbor Symp. Quant. Biol.* 50:399-409 (1986); MacDonald, *Hepatology* 7:425-515 (1987)); insulin gene control region which is active in pancreatic beta cells (Hanahan et al., *Nature* 315:115-122 (1985)), immunoglobulin gene control region which is active in lymphoid cells (Grosschedl et al., *Cell* 38:647-658 (1984); Adams et al., *Nature* 318:533-538 (1985); Alexander et al., *Mol. Cell Biol.* 7:1436-1444 (1987)), mouse mammary tumor virus control region which is active in testicular, breast, lymphoid and mast cells (Leder et al., *Cell* 45:485-495 (1986)), albumin gene

-59-

control region which is active in liver (Pinckert et al., *Genes and Devel.* 1:268-276 (1987)), alpha-fetoprotein gene control region which is active in liver (Krumlauf et al., *Mol. Cell. Biol.* 5:1639-1648 (1985); Hammer et al., *Science* 235:53-58 (1987)), alpha-1 antitrypsin gene control region which is active in liver  
5 (Kelsey et al., *Genes and Devel.* 1:161-171 (1987)), beta globin gene control region which is active in myeloid cells (Mogam et al., *Nature* 315:338-340 (1985); Kollias et al., *Cell* 46:89-94 (1986)), myelin basic protein gene control region which is active in oligodendrocyte cells of the brain (Readhead et al., *Cell* 48:703-712 (1987)), myosin light chain-2 gene control region which is active in  
10 skeletal muscle (Sani, *Nature* 314:283-286 (1985)), and gonadotrophic releasing hormone gene control region which is active in gonadotrophs of the hypothalamus (Mason et al., *Science* 234:1372-1378 (1986)).

In a specific embodiment, a vector is used that contains a promoter operably linked to nucleic acids encoding an MTSP protein, or a domain,  
15 fragment, derivative or homolog, thereof, one or more origins of replication, and optionally, one or more selectable markers (*e.g.*, an antibiotic resistance gene). Expression vectors containing the coding sequences, or portions thereof, of an MTSP protein, is made, for example, by subcloning the coding portions into the EcoRI restriction site of each of the three pGEX vectors (glutathione S-  
20 transferase expression vectors (Smith and Johnson, *Gene* 7:31-40 (1988)). This allows for the expression of products in the correct reading frame. Exemplary vectors and systems for expression of the protease domains of the MTSP proteins include the well-known *Pichia* vectors (available, for example, from Invitrogen, San Diego, CA), particularly those designed for secretion of the  
25 encoded proteins. The protein can also be expressed cytoplasmically, such as in the inclusion bodies. One exemplary vector is described in the EXAMPLES.

Plasmids for transformation of *E. coli* cells, include, for example, the pET expression vectors (see, U.S. patent 4,952,496; available from NOVAGEN, Madison, WI; see, also literature published by Novagen describing the system).  
30 Such plasmids include pET 11a, which contains the T7lac promoter, T7 terminator, the inducible *E. coli* lac operator, and the lac repressor gene; pET 12a-c, which contains the T7 promoter, T7 terminator, and the *E. coli* ompT



-60-

secretion signal; and pET 15b and pET19b (NOVAGEN, Madison, WI), which contain a His-Tag™ leader sequence for use in purification with a His column and a thrombin cleavage site that permits cleavage following purification over the column; the T7-lac promoter region and the T7 terminator.

- 5           The vectors are introduced into host cells, such as *Pichia* cells and bacterial cells, such as *E. coli*, and the proteins expressed therein. Exemplary *Pichia* strains, include, for example, GS115. Exemplary bacterial hosts contain chromosomal copies of DNA encoding T7 RNA polymerase operably linked to an inducible promoter, such as the lacUV promoter (see, U.S. Patent No.
- 10   4,952,496). Such hosts include, but are not limited to, the lysogenic *E. coli* strain BL21(DE3).

#### Expression and production of proteins

- The MTSP domains, derivatives and analogs be produced by various methods known in the art. For example, once a recombinant cell expressing an
- 15   MTSP protein, or a domain, fragment or derivative thereof, is identified, the individual gene product can be isolated and analyzed. This is achieved by assays based on the physical and/or functional properties of the protein, including, but not limited to, radioactive labeling of the product followed by analysis by gel electrophoresis, immunoassay, cross-linking to marker-labeled
- 20   product. The MTSP proteins can be isolated and purified by standard methods known in the art (either from natural sources or recombinant host cells expressing the complexes or proteins), including but not restricted to column chromatography (*e.g.*, ion exchange, affinity, gel exclusion, reversed-phase high pressure and fast protein liquid), differential centrifugation, differential solubility,
- 25   or by any other standard technique used for the purification of proteins. Functional properties can be evaluated using any suitable assay known in the art.

- Alternatively, once an MTSP protein or its domain or derivative is identified, the amino acid sequence of the protein can be deduced from the
- 30   nucleotide sequence of the gene which encodes it. As a result, the protein or its domain or derivative can be synthesized by standard chemical methods known in the art (*e.g.* see Hunkapiller et al, *Nature* 310:105-111 (1984)).

-61-

Manipulations of MTSP protein sequences can be made at the protein level. Also contemplated herein are MTSP protein proteins, domains thereof, derivatives or analogs or fragments thereof, which are differentially modified during or after translation, *e.g.*, by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand. Any of numerous chemical modifications can be carried out by known techniques, including but not limited to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH<sub>4</sub>, acetylation, formylation, oxidation, reduction, metabolic synthesis in the presence of tunicamycin and other such agents.

In addition, domains, analogs and derivatives of an MTSP protein can be chemically synthesized. For example, a peptide corresponding to a portion of an MTSP protein, which includes the desired domain or which mediates the desired activity *in vitro* can be synthesized by use of a peptide synthesizer. Furthermore, if desired, nonclassical amino acids or chemical amino acid analogs can be introduced as a substitution or addition into the MTSP protein sequence. Non-classical amino acids include but are not limited to the D-isomers of the common amino acids,  $\alpha$ -amino isobutyric acid, 4-aminobutyric acid, Abu, 2-aminobutyric acid,  $\epsilon$ -Abu,  $\epsilon$ -Ahx, 6-amino hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, citrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine,  $\beta$ -alanine, fluoro-amino acids, designer amino acids such as  $\beta$ -methyl amino acids, Ca-methyl amino acids, Na-methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

In cases where natural products are suspected of being mutant or are isolated from new species, the amino acid sequence of the MTSP protein isolated from the natural source, as well as those expressed *in vitro*, or from synthesized expression vectors *in vivo* or *in vitro*, can be determined from analysis of the DNA sequence, or alternatively, by direct sequencing of the

-62-

isolated protein. Such analysis can be performed by manual sequencing or through use of an automated amino acid sequenator.

#### Modifications

- A variety of modification of the MTSP proteins and domains are contemplated herein. An MTSP-encoding nucleic acid molecule be modified by any of numerous strategies known in the art (Sambrook et al., 1990, *Molecular Cloning, A Laboratory Manual*, 2d ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York). The sequences can be cleaved at appropriate sites with restriction endonuclease(s), followed by further enzymatic modification if desired, isolated, and ligated *in vitro*. In the production of the gene encoding a domain, derivative or analog of MTSP, care should be taken to ensure that the modified gene retains the original translational reading frame, uninterrupted by translational stop signals, in the gene region where the desired activity is encoded.
- 15        Additionally, the MTSP-encoding nucleic acid molecules can be mutated *in vitro* or *in vivo*, to create and/or destroy translation, initiation, and/or termination sequences, or to create variations in coding regions and/or form new restriction endonuclease sites or destroy pre-existing ones, to facilitate further *in vitro* modification. Also, as described herein muteins with primary sequence alterations, such as replacements of Cys residues and elimination or addition of glycosylation sites are contemplated; the MTSP7 of SEQ ID No. 16 has no potential glycosylation sites. Such mutations can be effected by any technique for mutagenesis known in the art, including, but not limited to, chemical mutagenesis and *in vitro* site-directed mutagenesis (Hutchinson et al., *J. Biol. Chem.* 253:6551-6558 (1978)), use of TAB<sup>®</sup> linkers (Pharmacia). In one embodiment, for example, an MTSP protein or domain thereof is modified to include a fluorescent label. In other specific embodiments, the MTSP protein is modified to have a heterobifunctional reagent, such heterobifunctional reagents can be used to crosslink the members of the complex.
- 25        The MTSP proteins can be isolated and purified by standard methods known in the art (either from natural sources or recombinant host cells expressing the complexes or proteins), including but not restricted to column
- 30

-63-

chromatography (*e.g.*, ion exchange, affinity, gel exclusion, reversed-phase high pressure and fast protein liquid), differential centrifugation, differential solubility, or by any other standard technique used for the purification of proteins.

Functional properties can be evaluated using any suitable assay known in the art.

Alternatively, once a MTSP or its domain or derivative is identified, the amino acid sequence of the protein can be deduced from the nucleotide sequence of the gene which encodes it. As a result, the protein or its domain or derivative can be synthesized by standard chemical methods known in the art (*e.g.*, see Hunkapiller et al, *Nature*, 310:105-111 (1984)).

Manipulations of MTSP sequences can be made at the protein level. MTSP domains, derivatives or analogs or fragments, which are differentially modified during or after translation, *e.g.*, by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule and other cellular ligand, are contemplated herein. Any of numerous chemical modifications can be carried out by known techniques, including but not limited to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH<sub>4</sub>, acetylation, formylation, oxidation, reduction, metabolic synthesis in the presence of tunicamycin.

In addition, domains, analogs and derivatives of a MTSP can be chemically synthesized. For example, a peptide corresponding to a portion of a MTSP, which comprises the desired domain or which mediates the desired activity *in vitro* can be synthesized by use of a peptide synthesizer.

Furthermore, if desired, nonclassical amino acids or chemical amino acid analogs can be introduced as a substitution or addition into the MTSP sequence. Non-classical amino acids include but are not limited to the D-isomers of the common amino acids,  $\alpha$ -amino isobutyric acid, 4-aminobutyric acid, Abu, 2-aminobutyric acid,  $\epsilon$ -Abu, e-Ahx, 6-amino hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, citrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine,  $\beta$ -alanine, fluoro-amino acids, designer amino acids such as  $\beta$ -

-64-

methyl amino acids, Ca-methyl amino acids, Na-methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

#### **F. SCREENING METHODS**

5           The single chain protease domains, as shown herein, can be used in a variety of methods to identify compounds that modulate the activity thereof. For MTSPs that exhibit higher activity or expression in tumor cells, compounds that inhibit the proteolytic activity are of particular interest. For any MTSPs that are active at lower levels in tumor cells, compounds or agents that enhance the  
10 activity are potentially of interest. In all instances the identified compounds will include agents that are candidate cancer treatments.

Several types of assays are exemplified and described herein. It is understood that the protease domains can be used in other assays. It is shown here, however, that the single chain protease domains exhibit catalytic activity.  
15 As such they are ideal for *in vitro* screening assays. They can also be used in binding assays.

The MTSP7 full length zymogens, activated enzymes, single and two chain protease domains are contemplated for use in any screening assay known to those of skill in the art, including those provided herein. Hence the following  
20 description, if directed to proteolytic assays is intended to apply to use of a single chain protease domain or a catalytically active portion thereof of any MTSP, including an MTSP7. Other assays, such as binding assays are provided herein, particularly for use with an MTSP7, including any variants, such as splice variants thereof.

##### **25           1. Catalytic Assays for identification of agents that modulate the protease activity of an MTSP protein**

Methods for identifying a modulator of the catalytic activity of an MTSP, particularly a single chain protease domain or catalytically active portion thereof, are provided herein. The methods can be practiced by: a) contacting the  
30 MTSP7, a full-length zymogen or activated form, and particularly a single-chain domain thereof, with a substrate of the MTSP7 in the presence of a test substance, and detecting the proteolysis of the substrate, whereby the activity

-65-

of the MTSP7 is assessed, and comparing the activity to a control. For example, the control can be the activity of the MTSP7 assessed by contacting an MTSP7, including a full-length zymogen or activated form, and particularly a single-chain domain thereof, particularly a single-chain domain thereof, with a substrate of the MTSP7, and detecting the proteolysis of the substrate, whereby the activity of the MTSP7 is assessed. The results in the presence and absence of the test compounds are compared. A difference in the activity indicates that the test substance modulates the activity of the MTSP7.

In one embodiment a plurality of the test substances are screened simultaneously in the above screening method. In another embodiment, the MTSP7 is isolated from a target cell as a means for then identifying agents that are potentially specific for the target cell.

In still another embodiment, The test substance is a therapeutic compound, and whereby a difference of the MTSP7 activity measured in the presence and in the absence of the test substance indicates that the target cell responds to the therapeutic compound.

One method include the steps of (a) contacting the MTSP7 protein or protease domain thereof with one or a plurality of test compounds under conditions conducive to interaction between the ligand and the compounds; and (b) identifying one or more compounds in the plurality that specifically binds to the ligand.

Another method provided herein includes the steps of a) contacting an MTSP7 protein or protease domain thereof with a substrate of the MTSP7 protein, and detecting the proteolysis of the substrate, whereby the activity of the MTSP7 protein is assessed; b) contacting the MTSP7 protein with a substrate of the MTSP7 protein in the presence of a test substance, and detecting the proteolysis of the substrate, whereby the activity of the MTSP7 protein is assessed; and c) comparing the activity of the MTSP7 protein assessed in steps a) and b), whereby the activity measured in step a) differs from the activity measured in step b) indicates that the test substance modulates the activity of the MTSP7 protein.

-66-

In another embodiment, a plurality of the test substances are screened simultaneously. In comparing the activity of an MTSP7 protein in the presence and absence of a test substance to assess whether the test substance is a modulator of the MTSP7 protein, it is unnecessary to assay the activity in parallel, although such parallel measurement is typically employed. It is possible to measure the activity of the MTSP7 protein at one time point and compare the measured activity to a historical value of the activity of the MTSP7 protein.

For instance, one can measure the activity of the MTSP7 protein in the presence of a test substance and compare with historical value of the activity of the MTSP7 protein measured previously in the absence of the test substance, and *vice versa*. This can be accomplished, for example, by providing the activity of the MTSP7 protein on an insert or pamphlet provided with a kit for conducting the assay.

Methods for selecting substrates for a particular MTSP are described in the EXAMPLES, and particular proteolytic assays are exemplified.

Combinations and kits containing the combinations optionally including instructions for performing the assays are provided. The combinations include an MTSP7 protein and a substrate of the MTSP7 protein to be assayed; and, optionally reagents for detecting proteolysis of the substrate. The substrates, which are typically proteins subject to proteolysis by a particular MTSP7 protein, can be identified empirically by testing the ability of the MTSP7 protein to cleave the test substrate. Substrates that are cleaved most effectively (i.e., at the lowest concentrations and/or fastest rate or under desirable conditions), are identified.

Additionally provided herein is a kit containing the above-described combination. The kit optionally further includes instructions for identifying a modulator of the activity of an MTSP7 protein. Any MTSP7 protein is contemplated as target for identifying modulators of the activity thereof.

-67-

## 2. Binding assays

Also provided herein are methods for identification and isolation of agents, particularly compounds that bind to MTSP7s. The assays are designed to identify agents that bind to the zymogen form, the single chain isolated  
5 protease domain (or a protein, other than an MTSP7 protein, that contains the protease domain of an MTSP7 protein), and to the activated form, including the activated form derived from the full length zymogen or from an extended protease domain. The identified compounds are candidates or leads for identification of compounds for treatments of tumors and other disorders and  
10 diseases involving aberrant angiogenesis. The MTSP7 proteins used in the methods include any MTSP7 protein as defined herein, and generally use MTSP7 single chain domain or proteolytically active portion thereof.

A variety of methods are provided herein. These methods can be performed in solution or in solid phase reactions in which the MTSP7 protein(s)  
15 or protease domain(s) thereof are linked, either directly or indirectly via a linker, to a solid support. Screening assays are described in the Examples, and these assays have been used to identify candidate compounds. For purposes herein, all binding assays described above are provided for MTSP7.

Methods for identifying an agent, such as a compound, that specifically  
20 binds to an MTSP7 single chain protease domain, a zymogen or full-length activated MTSP7 or two chain protease domain thereof are provided herein. The method can be practiced by (a) contacting the MTSP7 with one or a plurality of test agents under conditions conducive to binding between the MTSP7 and an agent; and (b) identifying one or more agents within the plurality that specifically  
25 binds to the MTSP7.

For example, in practicing such methods the MTSP7 polypeptide is mixed with a potential binding partner or an extract or fraction of a cell under conditions that allow the association of potential binding partners with the polypeptide. After mixing, peptides, polypeptides, proteins or other molecules that have become  
30 associated with an MTSP7 are separated from the mixture. The binding partner that bound to the MTSP7 then can be removed and further analyzed. To identify and isolate a binding partner, the entire protein, for instance the entire disclosed



-68-

protein of SEQ ID Nos. 15 or 17 can be used. Alternatively, a fragment of the protein can be used.

A variety of methods can be used to obtain cell extracts. Cells can be disrupted using either physical or chemical disruption methods. Examples of physical disruption methods include, but are not limited to, sonication and mechanical shearing. Examples of chemical lysis methods include, but are not limited to, detergent lysis and enzyme lysis. A skilled artisan can readily adapt methods for preparing cellular extracts in order to obtain extracts for use in the present methods.

Once an extract of a cell is prepared, the extract is mixed with the MTSP7 under conditions in which association of the protein with the binding partner can occur. A variety of conditions can be used. Exemplary conditions are those that closely resemble conditions found in the cytoplasm of a human cell. Features such as osmolarity, pH, temperature, and the concentration of cellular extract used, can be varied to optimize the association of the protein with the binding partner.

After mixing under appropriate conditions, the bound complex is separated from the mixture. A variety of techniques can be used to separate the mixture. For example, antibodies specific to an MTSP7 can be used to immunoprecipitate the binding partner complex. Alternatively, standard chemical separation techniques such as chromatography and density/sediment centrifugation can be used.

After removing the non-associated cellular constituents in the extract, the binding partner can be dissociated from the complex using conventional methods. For example, dissociation can be accomplished by altering the salt concentration or pH of the mixture.

To aid in separating associated binding partner pairs from the mixed extract, the MTSP7 can be immobilized on a solid support. For example, the protein can be attached to a nitrocellulose matrix or acrylic beads. Attachment of the protein or a fragment thereof to a solid support aids in separating peptide/binding partner pairs from other constituents found in the extract. The

-69-

identified binding partners can be either a single protein or a complex made up of two or more proteins.

Alternatively, the nucleic acid molecules encoding the single chain proteases can be used in a yeast two-hybrid system. The yeast two-hybrid  
5 system has been used to identify other protein partner pairs and can readily be adapted to employ the nucleic acid molecules herein described.

Another *in vitro* binding assay, particularly for an MTSP7, uses a mixture of a polypeptide that contains at least the catalytic domain of one of these proteins and one or more candidate binding targets or substrates. After  
10 incubating the mixture under appropriate conditions, the ability of the MTSP7 or a polypeptide fragment thereof containing the catalytic domain to bind to or interact with the candidate substrate is assessed. For cell-free binding assays, one of the components includes or is coupled to a detectable label. The label can provide for direct detection, such as radioactivity, luminescence, optical or  
15 electron density, or indirect detection such as an epitope tag, an enzyme and other such agents. A variety of methods can be employed to detect the label depending on the nature of the label and other assay components. For example, the label can be detected bound to the solid substrate or a portion of the bound complex containing the label can be separated from the solid substrate, and the  
20 label thereafter detected.

### 3. Detection of signal transduction

The cell surface location of the MTSPs suggests a role for some or all of these proteins in signal transduction. Assays for assessing signal transduction are well known to those of skill in the art, and can be adapted for use with the  
25 MTSP7 protein.

Assays for identifying agents that effect or alter signal transduction mediated directly or indirectly, such as via activation of a pro-growth factor, by an MTSP7, particularly the full length or a sufficient portion to anchor the extracellular domain or a function portion thereof of an MTSP on the surface of a  
30 cell are provided. Such assays, include, for example, transcription based assays in which modulation of a transduced signal is assessed by detecting an effect on an expression from a reporter gene (see, *e.g.*, U.S. Patent No. 5,436,128).

-70-

#### 4. Methods for Identifying Agents that Modulate the Expression a Nucleic Acid Encoding an MTSP7

Another embodiment provides methods for identifying agents that modulate the expression of a nucleic acid encoding an MTSP7. Such assays use  
5 any available means of monitoring for changes in the expression level of the nucleic acids encoding an MTSP7.

In one assay format, cell lines that contain reporter gene fusions between the open reading frame of MTSP7 or a domain thereof, particularly the protease domain and any assayable fusion partner can be prepared. Numerous assayable  
10 fusion partners are known and readily available including the firefly luciferase gene and the gene encoding chloramphenicol acetyltransferase (Alam *et al.*, *Anal. Biochem.* 188: 245-54 (1990)). Cell lines containing the reporter gene fusions are then exposed to the agent to be tested under appropriate conditions and time. Differential expression of the reporter gene between samples exposed  
15 to the agent and control samples identifies agents which modulate the expression of a nucleic acid encoding an MTSP7.

Additional assay formats can be used to monitor the ability of the agent to modulate the expression of a nucleic acid encoding an MTSP7. For instance, mRNA expression can be monitored directly by hybridization to the nucleic acids.  
20 Cell lines are exposed to the agent to be tested under appropriate conditions and time and total RNA or mRNA is isolated by standard procedures (see, *e.g.*, Sambrook *et al.* (1989) MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed. Cold Spring Harbor Laboratory Press). Probes to detect differences in RNA expression levels between cells exposed to the agent and control cells can  
25 be prepared from the nucleic acids. Generally, although not necessarily, probes are designed to hybridize only with target nucleic acids under conditions of high stringency. Only highly complementary nucleic acid hybrids form under conditions of high stringency. Stringency should be chosen to maximize the difference in stability between the probe:target hybrid and potential probe:non-  
30 target hybrids.

Probes can be designed from the nucleic acids through methods known in the art. For instance, the G + C content of the probe and the probe length can

-71-

affect probe binding to its target sequence. Methods to optimize probe specificity are commonly available (see, *e.g.*, Sambrook *et al.* (1989) MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed. Cold Spring Harbor Laboratory Press); and Ausubel *et al.* (1995) CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, Greene Publishing Co., NY).

Hybridization conditions are modified using known methods (see, *e.g.*, Sambrook *et al.* (1989) MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed. Cold Spring Harbor Laboratory Press); and Ausubel *et al.* (1995) CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, Greene Publishing Co., NY), as required for each probe. Hybridization of total cellular RNA or RNA enriched for polyA RNA can be accomplished in any available format. For instance, total cellular RNA or RNA enriched for polyA RNA can be affixed to a solid support, and the solid support exposed to at least one probe comprising at least one, or part of one of the nucleic acid molecules under conditions in which the probe will specifically hybridize. Alternatively, nucleic acid fragments comprising at least one, or part of one of the sequences can be affixed to a solid support, such as a porous glass wafer. The glass wafer then can be exposed to total cellular RNA or polyA RNA from a sample under conditions in which the affixed sequences will specifically hybridize. Such glass wafers and hybridization methods are widely available, for example, those disclosed by Beattie (WO 95/11755). By examining for the ability of a given probe to specifically hybridize to an RNA sample from an untreated cell population and from a cell population exposed to the agent, agents which up or down regulate the expression of a nucleic acid encoding the protein having the sequence set forth in any of SEQ ID Nos. 15-18, particularly 18, are identified.

#### **5. Methods for Identifying Agents that Modulate at Least One Activity of an MTSP7**

Methods for identifying agents that modulate at least one activity of an MTSP7 are provided. Such methods or assays can use any means of monitoring or detecting the desired activity. In one format, the relative amounts of a protein between a cell population that has been exposed to the agent to be tested compared to an un-exposed control cell population can be assayed (*e.g.*, a

-72-

prostate cancer cell line, a lung cancer cell line, a colon cancer cell line or a breast cancer cell line). In this format, probes, such as specific antibodies, are used to monitor the differential expression of the protein in the different cell populations. Cell lines or populations are exposed to the agent to be tested  
5 under appropriate conditions and time. Cellular lysates can be prepared from the exposed cell line or population and a control, unexposed cell line or population. The cellular lysates are then analyzed with the probe.

For example, N- and C- terminal fragments of the MTSP7 can be expressed in bacteria and used to search for proteins which bind to these  
10 fragments. Fusion proteins, such as His-tag or GST fusion to the N- or C-terminal regions of the MTSP7 can be prepared for use as a substrate. These fusion proteins can be coupled to, for example, Glutathione-Sepharose beads and then probed with cell lysates. Prior to lysis, the cells can be treated with a candidate agent which can modulate an MTSP7 or proteins that interact with  
15 domains thereon. Lysate proteins binding to the fusion proteins can be resolved by SDS-PAGE, isolated and identified by protein sequencing or mass spectroscopy, as is known in the art.

Antibody probes are prepared by immunizing suitable mammalian hosts in appropriate immunization protocols using the peptides, polypeptides or proteins  
20 if they are of sufficient length (*e.g.*, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, 25, 30, 35, 40 or more consecutive amino acids the MTSP7 protein or if required to enhance immunogenicity, conjugated to suitable carriers. Methods for preparing immunogenic conjugates with carriers, such as bovine serum albumin (BSA), keyhole limpet hemocyanin (KLH), or other carrier proteins are  
25 well known in the art. In some circumstances, direct conjugation using, for example, carbodiimide reagents can be effective; in other instances linking reagents such as those supplied by Pierce Chemical Co., Rockford, IL, can be desirable to provide accessibility to the hapten. Hapten peptides can be extended at either the amino or carboxy terminus with a Cys residue or  
30 interspersed with cysteine residues, for example, to facilitate linking to a carrier. Administration of the immunogens is conducted generally by injection over a suitable time period and with use of suitable adjuvants, as is generally

-73-

understood in the art. During the immunization schedule, titers of antibodies are taken to determine adequacy of antibody formation.

Anti-peptide antibodies can be generated using synthetic peptides corresponding to, for example, the carboxy terminal amino acids of the MTSP7.

5 Synthetic peptides can be as small as 1-3 amino acids in length, but are generally at least 4 or more amino acid residues long. The peptides can be coupled to KLH using standard methods and can be immunized into animals, such as rabbits or ungulate. Polyclonal antibodies then can be purified, for example using Actigel beads containing the covalently bound peptide.

10 While the polyclonal antisera produced in this way can be satisfactory for some applications, for pharmaceutical compositions, generally monoclonal preparations are used. Immortalized cell lines which secrete the desired monoclonal antibodies can be prepared using the standard method of Kohler *et al.*, (*Nature* 256: 495-7 (1975)) or modifications which effect immortalization of  
15 lymphocytes or spleen cells, as is generally known. The immortalized cell lines secreting the desired antibodies are screened by immunoassay in which the antigen is the peptide hapten, polypeptide or protein. When the appropriate immortalized cell culture secreting the desired antibody is identified, the cells can be cultured either *in vitro* or by production *in vivo* via ascites fluid. Of particular  
20 interest, are monoclonal antibodies that recognize the catalytic domain of the an MTSP7.

Additionally, the zymogen or two-chain forms the MTSP7 can be used to make monoclonal antibodies which recognize conformation epitopes. The desired monoclonal antibodies are then recovered from the culture supernatant  
25 or from the ascites supernatant. Fragments of the monoclonals or the polyclonal antisera which contain the immunologically significant portion can be used as antagonists, as well as the intact antibodies. Use of immunologically reactive fragments, such as the Fab, Fab', of F(ab')<sub>2</sub> fragments are often used, especially in a therapeutic context, as these fragments are generally less immunogenic than  
30 the whole immunoglobulin.

-74-

The antibodies or fragments can also be produced. Regions that bind specifically to the desired regions of receptor can also be produced in the context of chimeras with multiple species origin.

Agents that are assayed in the above method can be randomly selected  
5 or rationally selected or designed.

The agents can be, as examples, peptides, small molecules, and carbohydrates. A skilled artisan can readily recognize that there is no limit as to the structural nature of the agents.

The peptide agents can be prepared using standard solid phase (or  
10 solution phase) peptide synthesis methods, as is known in the art. In addition, the DNA encoding these peptides can be synthesized using commercially available oligonucleotide synthesis instrumentation and produced recombinantly using standard recombinant production systems. The production using solid phase peptide synthesis is necessitated if non-gene-encoded amino acids are to  
15 be included.

**G. Assay formats and selection of test substances**

A variety of formats and detection protocols are known for performing screening assays. Any such formats and protocols can be adapted for identifying modulators of MTSP7 protein activities. The following includes a  
20 discussion of exemplary protocols.

**1. High throughput screening assays**

Although the above-described assay can be conducted where a single MTSP7 protein is screened, and/or a single test substance is screened in one assay, the assay is generally conducted in a high throughput screening mode,  
25 *i.e.*, a plurality of the MTSP proteins are screened against and/or a plurality of the test substances are screened simultaneously (*See generally, High Throughput Screening: The Discovery of Bioactive Substances* (Devlin, Ed.) Marcel Dekker, 1997; Sittampalam et al., *Curr. Opin. Chem. Biol.*, 1(3):384-91 (1997); and Silverman et al., *Curr. Opin. Chem. Biol.*, 2(3):397-403 (1998)). For  
30 example, the assay can be conducted in a multi-well (*e.g.*, 24-, 48-, 96-, or 384-well), chip or array format.

-75-

High-throughput screening (HTS) is the process of testing a large number of diverse chemical structures against disease targets to identify "hits" (Sittampalam et al., *Curr. Opin. Chem. Biol.*, 1(3):384-91 (1997)). Current state-of-the-art HTS operations are highly automated and computerized to handle sample preparation, assay procedures and the subsequent processing of large volumes of data.

Detection technologies employed in high-throughput screens depend on the type of biochemical pathway being investigated (Sittampalam et al., *Curr. Opin. Chem. Biol.*, 1(3):384-91 (1997)). These methods include, radiochemical methods, such as the scintillation proximity assays (SPA), which can be adapted to a variety of enzyme assays (Lerner et al., *J. Biomol. Screening*, 1:135-143 (1996); Baker et al., *Anal. Biochem.*, 239:20-24 (1996); Baum et al., *Anal. Biochem.*, 237:129-134 (1996); and Sullivan et al., *J. Biomol. Screening*, 2:19-23 (1997)) and protein-protein interaction assays (Braunwalder et al., *J. Biomol. Screening*, 1:23-26 (1996); Sonatore et al., *Anal. Biochem.*, 240:289-297 (1996); and Chen et al., *J. Biol. Chem.*, 271:25308-25315 (1996)), and non-isotopic detection methods, including but are not limited to, colorimetric and luminescence detection methods, resonance energy transfer (RET) methods, time-resolved fluorescence (HTRF) methods, cell-based fluorescence assays, such as fluorescence resonance energy transfer (FRET) procedures (see, e.g., Gonzalez et al., *Biophys. J.*, 69:1272-1280 (1995)), fluorescence polarization or anisotropy methods (see, e.g., Jameson et al., *Methods Enzymol.*, 246:283-300 (1995); Jolley, *J. Biomol. Screening*, 1:33-38 (1996); Lynch et al., *Anal. Biochem.*, 247:77-82 (1997)), fluorescence correlation spectroscopy (FCS) and other such methods.

## 2. Test Substances

Test compounds, including small molecules, antibodies, proteins, nucleic acids, peptides, and libraries and collections thereof, can be screened in the above-described assays and assays described below to identify compounds that modulate the activity an MTSP7 protein. Rational drug design methodologies that rely on computational chemistry can be used to screen and identify candidate compounds.



-76-

The compounds identified by the screening methods include inhibitors, including antagonists, and can be agonists. Compounds for screening are any compounds and collections of compounds available, known or that can be prepared.

5

a. **Selection of Compounds**

- Compounds can be selected for their potency and selectivity of inhibition of serine proteases, especially an MTSP7 protein. As described herein, and as generally known, a target serine protease and its substrate are combined under assay conditions permitting reaction of the protease with its substrate. The
- 10 assay is performed in the absence of test compound, and in the presence of increasing concentrations of the test compound. The concentration of test compound at which 50% of the serine protease activity is inhibited by the test compound is the  $IC_{50}$  value (Inhibitory Concentration) or  $EC_{50}$  (Effective Concentration) value for that compound. Within a series or group of test
- 15 compounds, those having lower  $IC_{50}$  or  $EC_{50}$  values are considered more potent inhibitors of the serine protease than those compounds having higher  $IC_{50}$  or  $EC_{50}$  values. The  $IC_{50}$  measurement is often used for more simplistic assays, whereas the  $EC_{50}$  is often used for more complicated assays, such as those employing cells.
- 20 Typically candidate compounds have an  $IC_{50}$  value of 100 nM or less as measured in an *in vitro* assay for inhibition of MTSP7 protein activity. The test compounds also are evaluated for selectivity toward a serine protease. As described herein, and as generally known, a test compound is assayed for its potency toward a panel of serine proteases and other enzymes and an  $IC_{50}$  value
- 25 or  $EC_{50}$  value is determined for each test compound in each assay system. A compound that demonstrates a low  $IC_{50}$  value or  $EC_{50}$  value for the target enzyme, *e.g.*, MTSP7 protein, and a higher  $IC_{50}$  value or  $EC_{50}$  value for other enzymes within the test panel (*e.g.*, urokinase tissue plasminogen activator, thrombin, Factor Xa), is considered to be selective toward the target enzyme.
- 30 Generally, a compound is deemed selective if its  $IC_{50}$  value or  $EC_{50}$  value in the target enzyme assay is at least one order of magnitude less than the next smallest  $IC_{50}$  value or  $EC_{50}$  value measured in the selectivity panel of enzymes.

-77-

Compounds also are evaluated for their activity *in vivo*. The type of assay chosen for evaluation of test compounds will depend on the pathological condition to be treated or prevented by use of the compound, as well as the route of administration to be evaluated for the test compound.

5           For instance, to evaluate the activity of a compound to reduce tumor growth through inhibition of MTSP7 protein, the procedures described by Jankun et al., *Canc. Res.*, 57:559-563 (1997) to evaluate PAI-1 can be employed. Briefly, the ATCC cell lines DU145 and LnCaP are injected into SCID mice. After tumors are established, the mice are given test compound according to a dosing  
10       regime determined from the compound's *in vitro* characteristics. The Jankun et al. compound was administered in water. Tumor volume measurements are taken twice a week for about five weeks. A compound is deemed active if an animal to which the compound was administered exhibited decreased tumor volume, as compared to animals receiving appropriate control compounds.

15           Another *in vivo* experimental model designed to evaluate the effect of p-aminobenzamidine, a swine protease inhibitor, on reducing tumor volume is described by Billström et al., *Int. J. Cancer*, 61:542-547 (1995).

To evaluate the ability of a compound to reduce the occurrence of, or inhibit, metastasis, the procedures described by Kobayashi et al., *Int. J. Canc.*,  
20       57:727-733d (1994) can be employed. Briefly, a murein xenograft selected for high lung colonization potential is injected into C57B1/6 mice i.v. (experimental metastasis) or s.c. into the abdominal wall (spontaneous metastasis). Various concentrations of the compound to be tested can be admixed with the tumor cells in Matrigel prior to injection. Daily i.p. injections of the test compound are  
25       made either on days 1-6 or days 7-13 after tumor inoculation. The animals are sacrificed about three or four weeks after tumor inoculation, and the lung tumor colonies are counted. Evaluation of the resulting data permits a determination as to efficacy of the test compound, optimal dosing and route of administration.

The activity of the tested compounds toward decreasing tumor volume  
30       and metastasis can be evaluated in known models (see, e.g., Rabbani et al., *Int. J. Cancer* 63:840-845 (1995)). Evaluation of the resulting data permits a determination of the efficacy of a test compound, optimal dosing, and route of

-78-

administration (see, also, Xing et al., *Canc. Res.*, 57:3585-3593 (1997), which describes a related protocol).

To evaluate the anti-angiogenesis activity of a compound, a rabbit cornea neovascularization model can be employed (see, e.g., Avery et al. (1990) *Arch. Ophthalmol.*, 108:1474-147). Averu et al. describes anesthetizing New Zealand albino rabbits and then making a central corneal incision and forming a radial corneal pocket. A slow release prostaglandin pellet was placed in the pocket to induce neovascularization. Test compound was administered i.p. for five days, at which time the animals were sacrificed. The effect of the test compound is evaluated by review of periodic photographs taken of the limbus, which can be used to calculate the area of neovascular response and, therefore, limbal neovascularization. A decreased area of neovascularization as compared with appropriate controls indicates the test compound was effective at decreasing or inhibiting neovascularization.

An angiogenesis model used to evaluate the effect of a test compound in preventing angiogenesis is described by Min et al., *Canc. Res.*, 56:2428-2433 (1996). C57BL6 mice receive subcutaneous injections of a Matrigel mixture containing bFGF, as the angiogenesis-inducing agent, with and without the test compound. After five days, the animals are sacrificed and the Matrigel plugs, in which neovascularization can be visualized, are photographed. An experimental animal receiving Matrigel and an effective dose of test compound will exhibit less vascularization than a control animal or an experimental animal receiving a less- or non-effective dose of compound.

An *in vivo* system designed to test compounds for their ability to limit the spread of primary tumors is described by Crowley et al., *Proc. Natl. Acad. Sci.*, 90:5021-5025 (1993). Nude mice are injected with tumor cells (PC3) engineered to express CAT (chloramphenicol acetyltransferase). Compounds to be tested for their ability to decrease tumor size and/or metastases are administered to the animals, and subsequent measurements of tumor size and/or metastatic growths are made. In addition, the level of CAT detected in various organs provides an indication of the ability of the test compound to inhibit

-79-

metastasis; detection of less CAT in tissues of a treated animal versus a control animal indicates less CAT-expressing cells migrated to that tissue.

*In vivo* experimental modes designed to evaluate the inhibitory potential of a test serine protease inhibitors, using a tumor cell line F3II, the to be highly  
5 invasive, are described by Alonso et al., *Breast Canc. Res. Treat.*, 40:209-223 (1996). This group describes *in vivo* studies for toxicity determination, tumor growth, invasiveness, spontaneous metastasis, experimental lung metastasis, and an angiogenesis assay.

The CAM model (chick embryo chorioallantoic membrane model), first  
10 described by L. Ossowski in 1998 (*J. Cell Biol.*, 107:2437-2445 (1988)), provides another method for evaluating the inhibitory activity of a test compound. In the CAM model, tumor cells invade through the chorioallantoic membrane containing CAM with tumor cells in the presence of several serine protease inhibitors results in less or no invasion of the tumor cells through the  
15 membrane. Thus, the CAM assay is performed with CAM and tumor cells in the presence and absence of various concentrations of test compound. The invasiveness of tumor cells is measured under such conditions to provide an indication of the compound's inhibitory activity. A compound having inhibitory activity correlates with less tumor invasion.

20 The CAM model is also used in a standard assay of angiogenesis (*i.e.*, effect on formation of new blood vessels (Brooks et al., *Methods in Molecular Biology*, 129:257-269 (1999)). According to this model, a filter disc containing an angiogenesis inducer, such as basic fibroblast growth factor (bFDG) is placed onto the CAM. Diffusion of the cytokine into the CAM induces local  
25 angiogenesis, which can be measured in several ways such as by counting the number of blood vessel branch points within the CAM directly below the filter disc. The ability of identified compounds to inhibit cytokine-induced angiogenesis can be tested using this model. A test compound can either be added to the filter disc that contains the angiogenesis inducer, be placed directly  
30 on the membrane or be administered systemically. The extent of new blood vessel formation in the presence and/or absence of test compound can be compared using this model. The formation of fewer new blood vessels in the

-80-

presence of a test compound would be indicative of anti-angiogenesis activity. Demonstration of anti-angiogenesis activity for inhibitors of an MTSP7 protein indicates a role in angiogenesis for that MTSP protein.

**b. Known serine protease inhibitors**

5 Compounds for screening can be serine protease inhibitors, which can be tested for their ability to inhibit the activity of an MTSP7.

- Exemplary, but not limiting serine proteases, include the following known serine protease inhibitors are used in the screening assays: Serine Protease Inhibitor 3 (SPI-3) (Chen, M.C., et al., *Citokine*, 11(11):856-862 (1999));
- 10 Aprotinin (Iijima, R., et al., *J. Biochem. (Tokyo)*, 126(5):912-916 (1999)); Kazal-type serine protease inhibitor-like proteins (Niimi, T., et al., *Eur. J. Biochem.*, 266(1):282-292 (1999)); Kunitz-type serine protease inhibitor (Ravichandran, S., et al., *Acta Crystallogr. D. Biol. Crystallogr.*, 55(11):1814-1821 (1999)); Tissue factor pathway inhibitor-2/Matrix-associated serine protease inhibitor (TFPI-
- 15 2/MSPI), (Liu, Y., et al., *Arch. Biochem. Biophys.*, 370(1):112-8 (1999)); Bukunin, (Cui, C.Y., et al., *J. Invest. Dermatol.*, 113(2):182-8 (1999)); Nafmostat mesilate (Ryo, R., et al., *Vox Sang.*, 76(4):241-6 (1999)); TPCK (Huang, Y., et al., *Oncogene*, 18(23):3431-9 (1999)); A synthetic cotton-bound serine protease inhibitor (Edwards, J.V., et al., *Wound Repair Regen.*, 7(2):106-
- 20 18 (1999)); FUT-175 (Sawada, M., et al., *Stroke*, 30(3):644-50 (1999)); Combination of serine protease inhibitor FUT-0175 and thromboxane synthetase inhibitor OKY-046 (Kaminogo, M., et al., *Neurol. Med. Chir. (Tokyo)*, 38(11):704-8; discussion 708-9 (1998)); The rat serine protease inhibitor 2.1 gene (LeCam, A., et al., *Biochem. Biophys. Res. Commun.*, 253(2):311-4
- 25 (1998)); A new intracellular serine protease inhibitor expressed in the rat pituitary gland complexes with granzyme B (Hill, R.M., et al., *FEBS Lett.*, 440(3):361-4 (1998)); 3,4-Dichloroisocoumarin (Hammed, A., et al., *Proc. Soc. Exp. Biol. Med.*, 219(2):132-7 (1998)); LEX032 (Bains, A.S., et al., *Eur. J. Pharmacol.*, 356(1):67-72 (1998)); N-tosyl-L-phenylalanine chloromethyl ketone
- 30 (Dryjanski, M., et al., *Biochemistry*, 37(40):14151-6 (1998)); Mouse gene for the serine protease inhibitor neuroserpin (P112) (Berger, P., et al., *Gene*, 214(1-2):25-33 (1998)); Rat serine protease inhibitor 2.3 gene (Paul, C., et al., *Eur. J.*

- Biochem.*, 254(3):538-46 (1998)); Ecotin (Yang, S.Q., et al., *J. Mol. Biol.*, 279(4):945-57 (1998)); A 14 kDa plant-related serine protease inhibitor (Roch, P., et al., *Dev. Comp. Immunol.*, 22(1):1-12 (1998)); Matrix-associated serine protease inhibitor TFPI-2/33 kDa MSPI (Rao, C.N., et al., *Int. J. Cancer*, 76(5):749-56 (1998)); ONO-3403 (Hiwasa, T., et al., *Cancer Lett.*, 126(2):221-5 (1998)); Bdellastasin (Moser, M., et al., *Eur. J. Biochem.*, 253(1):212-20 (1998)); Bikunin (Xu, Y., et al., *J. Mol. Biol.*, 276(5):955-66 (1998)); Nafamostat mesilate (Mellgren, K., et al., *Thromb. Haemost.*, 79(2):342-7 (1998)); The growth hormone dependent serine protease inhibitor, Spi 2.1 (Maake, C., et al., *Endocrinology*, 138(12):5630-6 (1997)); Growth factor activator inhibitor type 2, a Kunitz-type serine protease inhibitor (Kawaguchi, T., et al., *J. Biol. Chem.*, 272(44):27558-64 (1997)); Heat-stable serine protease inhibitor protein from ovaries of the desert locust, *Schistocerca gregaria* (Hamdaoui, A., et al., *Biochem. Biophys. Res. Commun.*, 238(2):357-60 (1997)); Bikunin, (Delaria, K.A., et al., *J. Biol. Chem.*, 272(18):12209-14 (1997)); Human placental bikunin (Marlor, C.W., et al., *J. Biol. Chem.*, 272(10):12202-8 (1997)); Hepatocyte growth factor activator inhibitor, a novel Kunitz-type serine protease inhibitor (Shimomura, T., et al., *J. Biol. Chem.*, 272(10):6370-6 (1997)); FUT-187, oral serine protease inhibitor, (Shiozaki, H., et al., *Gan To Kagaku Ryoho*, 23(14): 1971-9 (1996)); Extracellular matrix-associated serine protease inhibitors (Mr 33,000, 31,000, and 27,000 (Rao, C.N., et al., *Arch. Biochem. Biophys.*, 335(1):82-92 (1996)); An irreversible isocoumarin serine protease inhibitor (Palencia, D.D., et al., *Biol. Reprod.*, 55(3):536-42 (1996)); 4-(2-aminoethyl)-benzenesulfonyl fluoride (AEBSF) (Nakabo, Y., et al., *J. Leukoc. Biol.*, 60(3):328-36 (1996)); Neuroserpin (Osterwalder, T., et al., *EMBO J.*, 15(12):2944-53 (1996)); Human serine protease inhibitor alpha-1-antitrypsin (Forney, J.R., et al., *J. Parasitol.*, 82(3):496-502 (1996)); Rat serine protease inhibitor 2.3 (Simar-Blanchet, A.E., et al., *Eur. J. Biochem.*, 236(2):638-48 (1996)); Gebaxate mesilate (parodi, F., et al., *J. Cardiothorac. Vasc. Anesth.*, 10(2):235-7 (1996)); Recombinant serine protease inhibitor, CPTI II (Stankiewicz, M., et al., *Acta Biochim. Pol.*, 43(3):525-9 (1996)); A cysteine-rich serine protease inhibitor (Guamerin II) (Kim,

-82-

- D.R., et al., *J. Enzym. Inhib.*, 10(2):81-91 (1996)); Diisopropylfluorophosphate (Lundqvist, H., et al., *Inflamm. Res.*, 44(12):510-7 (1995)); Nexin 1 (Yu, D.W., et al., *J. Cell Sci.*, 108(Pt 12):3867-74 (1995)); LEX032 (Scalia, R., et al., *Shock*, 4(4):251-6 (1995)); Protease nexin I (Houenou, L.J., et al., *Proc. Natl. Acad. Sci. U.S.A.*, 92(3):895-9 (1995)); Chymase-directed serine protease inhibitor (Woodard S.L., et al., *J. Immunol.*, 153(11):5016-25 (1994)); N-alpha-tosyl-L-lysyl-chloromethyl ketone (TLCK) (Bourinbaier, A.S., et al., *Cell Immunol.*, 155(1):230-6 (1994)); Smpi56 (Ghendler, Y., et al., *Exp. Parasitol.*, 78(2):121-31 (1994)); Schistosoma haematobium serine protease (Blanton, R.E., et al., *Mol. Biochem. Parasitol.*, 63(1):1-11 (1994)); Spi-1 (Warren, W.C., et al., *Mol. Cell Endocrinol.*, 98(1):27-32 (1993)); TAME (Jessop, J.J., et al., *Inflammation*, 17(5):613-31 (1993)); Antithrombin III (Kalaria, R.N., et al., *Am. J. Pathol.*, 143(3):886-93 (1993)); FOY-305 (Ohkoshi, M., et al., *Anticancer Res.*, 13(4):963-6 (1993)); Camostat mesilate (Senda, S., et al., *Intern. Med.*, 32(4):350-4 (1993)); Pigment epithelium-derived factor (Steele, F.R., et al., *Proc. Natl. Acad. Sci. U.S.A.*, 90(4):1526-30 (1993)); Antistasin (Holstein, T.W., et al., *FEBS Lett.*, 309(3):288-92 (1992)); The vaccinia virus K2L gene encodes a serine protease inhibitor (Zhou, J., et al., *Virology*, 189(2):678-86 (1992)); Bowman-Birk serine-protease inhibitor (Werner, M.H., et al., *J. Mol. Biol.*, 225(3):873-89 (1992)); FUT-175 (Yanamoto, H., et al., *Neurosurgery*, 30(3):358-63 (1992)); FUT-175; (Yanamoto, H., et al., *Neurosurgery*, 30(3):351-6, discussion 356-7 (1992)); PAI-I (Yreadwell, B.V., et al., *J. Orthop. Res.*, 9(3):309-16 (1991)); 3,4-Dichloroisocoumarin (Rusbridge, N.M., et al., *FEBS Lett.*, 268(1):133-6 (1990)); Alpha 1-antichymotrypsin (Lindmark, B.E., et al., *Am. Rev. Respir. Dis.*, 141(4 Pt 1):884-8 (1990)); P-toluenesulfonyl-L-arginine methyl ester (TAME) (Scuderi, P., *J. Immunol.*, 143(1):168-73 (1989)); Aprotinin (Seto, S., et al., *Adv. Exp. Med. Biol.*, 247B:49-54 (1989)); Alpha 1-antichymotrypsin (Abraham, C.R., et al., *Cell*, 52(4):487-501 (1988)); Contrapsin (Modha, J., et al., *Parasitology*, 96 (Pt 1):99-109 (1988)); (FOY-305) (Yamauchi, Y., et al., *Hiroshima J. Med. Sci.*, 36(1):81-7 No abstract available (1987)); Alpha 2-antiplasmin (Holmes, W.E., et al., *J. Biol. Chem.*, 262(4):1659-64 (1987)); 3,4-dichloroisocoumarin (Harper, J.W., et al., *Biochemistry*,

-83-

24(8):1831-41 (1985)); Diisopropylfluorophosphate (Tsutsui, K., et al., *Biochem. Biophys. Res. Commun.*, 123(1):271-7 (1984)); Gabexate mesilate (Hesse, B., et al., *Pharmacol. Res. Commun.*, 16(7):637-45 (1984)); Phenyl methyl sulfonyl fluoride (Dufer, J., et al., *Scand. J. Haematol.*, 32(1):25-32 (1984)); Aprotinin (Seto, S., et al., *Hypertension*, 5(6):893-9 (1983)); Protease inhibitor Cl-2 (McPhalen, C.A., et al., *J. Mol. Biol.*, 168(2):445-7 (1983)); Phenylmethanesulfonyl fluoride (Sekar V., et al., *Biochem. Biophys. Res. Commun.*, 89(2):474-8 (1979)); PGE1 (Feinstein, M.D., et al., *Prostaglandine*, 14(6):1075-93 (1977))

#### 10 c. Combinatorial libraries and other libraries

The source of compounds for the screening assays, can be libraries, including, but are not limited to, combinatorial libraries. Methods for synthesizing combinatorial libraries and characteristics of such combinatorial libraries are known in the art (See generally, *Combinatorial Libraries: Synthesis, Screening and Application Potential* (Cortese Ed.) Walter de Gruyter, Inc., 1995; Tietze and Lieb, *Curr. Opin. Chem. Biol.*, 2(3):363-71 (1998); Lam, *Anticancer Drug Des.*, 12(3):145-67 (1997); Blaney and Martin, *Curr. Opin. Chem. Biol.*, 1(1):54-9 (1997); and Schultz and Schultz, *Biotechnol. Prog.*, 12(6):729-43 (1996)).

20 Methods and strategies for generating diverse libraries, primarily peptide- and nucleotide-based oligomer libraries, have been developed using molecular biology methods and/or simultaneous chemical synthesis methodologies (see, e.g., Dower et al., *Annu. Rep. Med. Chem.*, 26:271-280 (1991); Fodor et al., *Science*, 251:767-773 (1991); Jung et al., *Angew. Chem. Int. Ed. Engl.*, 31:367-383 (1992); Zuckerman et al., *Proc. Natl. Acad. Sci. USA*, 89:4505-4509 (1992); Scott et al., *Science*, 249:386-390 (1990); Devlin et al., *Science*, 249:404-406 (1990); Cwirla et al., *Proc. Natl. Acad. Sci. USA*, 87:6378-6382 (1990); and Gallop et al., *J. Medicinal Chemistry*, 37:1233-1251 (1994)). The resulting combinatorial libraries potentially contain millions of compounds and that can be screened to identify compounds that exhibit a selected activity.

30 The libraries fall into roughly three categories: fusion-protein-displayed peptide libraries in which random peptides or proteins are presented on the



-84-

surface of phage particles or proteins expressed from plasmids; support-bound synthetic chemical libraries in which individual compounds or mixtures of compounds are presented on insoluble matrices, such as resin beads (*see, e.g.,* Lam et al., *Nature*, 354:82-84 (1991)) and cotton supports (*see, e.g.,* Eichler et al., *Biochemistry* 32:11035-11041 (1993)); and methods in which the compounds are used in solution (*see, e.g.,* Houghten et al., *Nature*, 354:84-86 (1991); Houghten et al., *BioTechniques*, 313:412-421 (1992); and Scott et al., *Curr. Opin. Biotechnol.*, 5:40-48 (1994)). There are numerous examples of synthetic peptide and oligonucleotide combinatorial libraries and there are many methods for producing libraries that contain non-peptidic small organic molecules. Such libraries can be based on basis set of monomers that are combined to form mixtures of diverse organic molecules or that can be combined to form a library based upon a selected pharmacophore monomer.

Either a random or a deterministic combinatorial library can be screened by the presently disclosed and/or claimed screening methods. In either of these two libraries, each unit of the library is isolated and/or immobilized on a solid support. In the deterministic library, one knows *a priori* a particular unit's location on each solid support. In a random library, the location of a particular unit is not known *a priori* although each site still contains a single unique unit. Many methods for preparing libraries are known to those of skill in this art (*see, e.g.,* Geysen et al., *Proc. Natl. Acad. Sci. USA*, 81:3998-4002 (1984), Houghten et al., *Proc. Natl. Acad. Sci. USA*, 81:5131-5135 (1985)). Combinatorial library generated by the any techniques known to those of skill in the art are contemplated (*see, e.g.,* Table 1 of Schultz and Schultz, *Biotechnol. Prog.*, 12(6):729-43 (1996)) for screening; Bartel *et al.*, *Science*, 261:1411-1418 (1993); Baumbach *et al.* *BioPharm*, (*Can*):24-35 (1992); Bock *et al.* *Nature*, 355:564-566 (1992); Borman, S., Combinatorial chemists focus on small molecules molecular recognition, and automation, *Chem. Eng. News*, 2(12):29 (1996); Boublik, et al., Eukaryotic Virus Display: Engineering the Major Surface Glycoproteins of the Autographa California Nuclear Polyhedrosis Virus (ACNPV) for the Presentation of Foreign Proteins on the Virus Surface, *Bio/Technology*, 13:1079-1084 (1995); Brenner, et al., Encoded Combinatorial

-85-

- Chemistry, *Proc. Natl. Acad. Sci. U.S.A.*, 89:5381-5383 (1992); Caflisch, et al., Computational Combinatorial Chemistry for *De Novo* Ligand Design: Review and Assessment, *Perspect. Drug Discovery Des.*, 3:51-84 (1995); Cheng, et al., Sequence-Selective Peptide Binding with a Peptido-A,B-*trans*-steroidal Receptor
- 5 Selected from an Encoded Combinatorial Library, *J. Am. Chem. Soc.*, 118:1813-1814 (1996); Chu, et al., Affinity Capillary Electrophoresis to Identify the Peptide in A Peptide Library that Binds Most Tightly to Vancomycin, *J. Org. Chem.*, 58:648-652 (1993); Clackson, et al., Making Antibody Fragments Using Phage Display Libraries, *Nature*, 352:624-628 (1991); Combs, et al., Protein
- 10 Structure-Based Combinatorial Chemistry: Discovery of Non-Peptide Binding Elements to Src SH3 Domain, *J. Am. Chem. Soc.*, 118:287-288 (1996); Cwirla, et al., Peptides On Phage: A Vast Library of Peptides for Identifying Ligands, *Proc. Natl. Acad. Sci. U.S.A.*, 87:6378-6382 (1990); Ecker, et al., Combinatorial Drug Discovery: Which Method will Produce the Greatest Value,
- 15 *Bio/Technology*, 13:351-360 (1995); Ellington, et al., *In Vitro* Selection of RNA Molecules That Bind Specific Ligands, *Nature*, 346:818-822 (1990); Ellman, J.A., Variants of Benzodiazepines, *J. Am. Chem. Soc.*, 114:10997 (1992); Erickson, et al., *The Proteins*; Neurath, H., Hill, R.L., Eds.: Academic: New York, 1976; pp. 255-257; Felici, et al., *J. Mol. Biol.*, 222:301-310 (1991); Fodor, et
- 20 al., Light-Directed, Spatially Addressable Parallel Chemical Synthesis, *Science*, 251:767-773 (1991); Francisco, et al., Transport and Anchoring of Beta-Lactamase to the External Surface of *E. Coli.*, *Proc. Natl. Acad. Sci. U.S.A.*, 89:2713-2717 (1992); Georgiou, et al., Practical Applications of Engineering Gram-Negative Bacterial Cell Surfaces, *TIBTECH*, 11:6-10 (1993); Geysen, et al.,
- 25 Use of peptide synthesis to probe viral antigens for epitopes to a resolution of a single amino acid, *Proc. Natl. Acad. Sci. U.S.A.*, 81:3998-4002 (1984); Glaser, et al., Antibody Engineering by Condon-Based Mutagenesis in a Filamentous Phage Vector System, *J. Immunol.*, 149:3903-3913 (1992); Gram, et al., In vitro selection and affinity maturation of antibodies from a naive combinatorial
- 30 immunoglobulin library, *Proc. Natl. Acad. Sci.*, 89:3576-3580 (1992); Han, et al., Liquid-Phase Combinatorial Synthesis, *Proc. Natl. Acad. Sci. U.S.A.*, 92:6419-6423 (1995); Hoogenboom, et al., Multi-Subunit Proteins on the

- Surface of Filamentous Phage: Methodologies for Displaying Antibody (Fab) Heavy and Light Chains, *Nucleic Acids Res.*, 19:4133-4137 (1991); Houghten, et al., General Method for the Rapid Solid-Phase Synthesis of Large Numbers of Peptides: Specificity of Antigen-Antibody Interaction at the Level of Individual
- 5 Amino Acids, *Proc. Natl. Acad. Sci. U.S.A.*, 82:5131-5135 (1985); Houghten, et al., The Use of Synthetic Peptide Combinatorial Libraries for the Determination of Peptide Ligands in Radio-Receptor Assays-Opioid-Peptides, *Bioorg. Med. Chem. Lett.*, 3:405-412 (1993); Houghten, et al., Generation and Use of Synthetic Peptide Combinatorial Libraries for Basic Research and Drug Discovery,
- 10 *Nature*, 354:84-86 (1991); Huang, et al., Discovery of New Ligand Binding Pathways in Myoglobin by Random Mutagenesis, *Nature Struct. Biol.*, 1:226-229 (1994); Huse, et al., Generation of a Large Combinatorial Library of the Immunoglobulin Repertoire In Phage Lambda, *Science*, 246:1275-1281 (1989); Janda, K.D., New Strategies for the Design of Catalytic Antibodies, *Biotechnol. Prog.*, 6:178-181 (1990); Jung, et al., Multiple Peptide Synthesis Methods and Their Applications, *Angew. Chem. Int. Ed. Engl.*, 31:367-486 (1992); Kang, et al., Linkage of Recognition and Replication Functions By Assembling Combinatorial Antibody Fab Libraries Along Phage Surfaces, *Proc. Natl. Acad. Sci. U.S.A.*, 88:4363-4366 (1991a); Kang, et al., Antibody Redesign by Chain
- 20 Shuffling from Random Combinatorial Immunoglobulin Libraries, *Proc. Natl. Acad. Sci. U.S.A.*, 88:11120-11123 (1991b); Kay, et al., An M13 Phage Library Displaying Random 38-Amino-Acid-Peptides as a Source of Novel Sequences with Affinity to Selected Targets Genes, *Gene*, 128:59-65 (1993); Lam, et al., A new type of synthetic peptide library for identifying ligand-binding activity,
- 25 *Nature*, 354:82-84 (1991) (published errata appear in *Nature*, 358:434 (1992) and *Nature*, 360:768 (1992); Lebl, et al., One Bead One Structure Combinatorial Libraries, *Biopolymers (Pept. Sci.)*, 37:177-198 (1995); Lerner, et al., Antibodies without Immunization, *Science*, 258:1313-1314 (1992); Li, et al., Minimization of a Polypeptide Hormone, *Science*, 270:1657-1660 (1995); Light, et al.,
- 30 Display of Dimeric Bacterial Alkaline Phosphatase on the Major Coat Protein of Filamentous Bacteriophage, *Bioorg. Med. Chem. Lett.*, 3:1073-1079 (1992); Little, et al., Bacterial Surface Presentation of Proteins and Peptides: An

-87-

- Alternative to Phage Technology, *Trends Biotechnol.*, 11:3-5 (1993); Marks, et al., By-Passing Immunization. Human Antibodies from V-Gene Libraries Displayed on Phage, *J. Mol. Biol.*, 222:581-597 (1991); Matthews, et al., Substrate Phage: Selection of Protease Substrates by Monovalent Phage Display, *Science*, 260:1113-1117 (1993); McCafferty, et al., Phage Enzymes: Expression and Affinity Chromatography of Functional Alkaline Phosphatase on the Surface of Bacteriophage, *Protein Eng.*, 4:955-961 (1991); Menger, et al., Phosphatase Catalysis Developed Via Combinatorial Organic Chemistry, *J. Org. Chem.*, 60:6666-6667 (1995); Nicolaou, et al., *Angew. Chem. Int. Ed. Engl.*, 34:2289-2291 (1995); Oldenburg, et al., Peptide Ligands for A Sugar-Binding Protein Isolated from a Random Peptide Library, *Proc. Natl. Acad. Sci. U.S.A.*, 89:5393-5397 (1992); Parmley, et al., Antibody-Selectable Filamentous fd Phage Vectors: Affinity Purification of Target Genes, *Genes*, 73:305-318 (1988); Pinilla, et al., Synthetic Peptide Combinatorial Libraries (SPCLS)--Identification of the Antigenic Determinant of Beta-Endorphin Recognized by Monoclonal Antibody-3E7, *Gene*, 128:71-76 (1993); Pinilla, et al., Review of the Utility of Soluble Combinatorial Libraries, *Biopolymers*, 37:221-240 (1995); Pistor, et al., Expression of Viral Hemagglutinin On the Surface of *E. Coli.*, *Klin. Wochenschr.*, 66:110-116 (1989); Pollack, et al., Selective Chemical Catalysis by an Antibody, *Science*, 234:1570-1572 (1986); Rigler, et al., Fluorescence Correlations, Single Molecule Detection and Large Number Screening: Applications in Biotechnology, *J. Biotechnol.*, 41:177-186 (1995); Sarvetnick, et al., Increasing the Chemical Potential of the Germ-Line Antibody Repertoire, *Proc. Natl. Acad. Sci. U.S.A.*, 90:4008-4011 (1993); Sastry, et al., Cloning of the Immunological Repertoire in *Escherichia Coli* for Generation of Monoclonal Catalytic Antibodies: Construction of a Heavy Chain Variable Region-Specific cDNA Library, *Proc. Natl. Acad. Sci. U.S.A.*, 86:5728-5732 (1989); Scott, et al., Searching for Peptide Ligands with an Epitope Library, *Science*, 249:386-390 (1990); Sears, et al., Engineering Enzymes for Bioorganic Synthesis: Peptide Bond Formation, *Biotechnol. Prog.*, 12:423-433 (1996); Simon, et. al., Peptides: A Modular Approach to Drug Discovery, *Proc. Natl. Acad. Sci. U.S.A.*, 89:9367-9371 (1992); Still, et al., Discovery of Sequence-Selective Peptide Binding by Synthetic Receptors Using

-88-

- Encoded Combinatorial Libraries, *Acc. Chem. Res.*, 29:155-163 (1996); Thompson, et al., Synthesis and Applications of Small Molecule Libraries, *Chem. Rev.*, 96:555-600 (1996); Tramontano, et al., Catalytic Antibodies, *Science*, 234:1566-1570 (1986); Wrighton, et al., Small Peptides as Potent Mimetics of the Protein Hormone Erythropoietin, *Science*, 273:458-464 (1996); York, et al., Combinatorial mutagenesis of the reactive site region in plasminogen activator inhibitor I, *J. Biol. Chem.*, 266:8595-8600 (1991); Zebedee, et al., Human Combinatorial Antibody Libraries to Hepatitis B Surface Antigen, *Proc. Natl. Acad. Sci. U.S.A.*, 89:3175-3179 (1992); Zuckerman, et al., Identification of Highest-Affinity Ligands by Affinity Selection from Equimolar Peptide Mixtures Generated by Robotic Synthesis, *Proc. Natl. Acad. Sci. U.S.A.*, 89:4505-4509 (1992).

For example, peptides that bind to an MTSP7 protein or a protease domain of an MTSP protein can be identified using phage display libraries. In an exemplary embodiment, this method can include a) contacting phage from a phage library with the MTSP7 protein or a protease domain thereof; (b) isolating phage that bind to the protein; and (c) determining the identity of at least one peptide coded by the isolated phage to identify a peptide that binds to an MTSP7 protein.

#### 20 H. Modulators of the activity of MTSP7 proteins

Provided herein are compounds, identified by screening or produced using the MTSP7 protein or protease domain in other screening methods, that modulate the activity of an MTSP7. These compounds act by directly interacting with the MTSP7 protein or by altering transcription or translation thereof. Such molecules include, but are not limited to, antibodies that specifically react with an MTSP7 protein, particularly with the protease domain thereof, antisense nucleic acids that alter expression of the MTSP7 protein or dsRNA, such as RNAi,, antibodies, peptide mimetics and other such compounds.

-89-

# **1. Antibodies**

Antibodies, including polyclonal and monoclonal antibodies, that specifically bind to the MTSP7 protein provided herein, particularly to the single chain protease domains thereof or the activated forms of the full-length or  
5 protease domain or the zymogen form, are provided. Typically, the antibody is a monoclonal antibody, and generally, the antibody specifically binds to the protease domain of the MTSP7 protein. In particular embodiments, antibodies to each of the single chain of the protease domain of MTSP7 are provided. Also provided are antibodies that specifically bind to any domain of MTSP7 and to  
10 two chain forms thereof.

The MTSP7 protein and domains, fragments, homologs and derivatives thereof can be used as immunogens to generate antibodies that specifically bind such immunogens. Such antibodies include but are not limited to polyclonal, monoclonal, chimeric, single chain, Fab fragments, and an Fab expression  
15 library. In a specific embodiment, antibodies to human MTSP7 protein are produced. In another embodiment, complexes formed from fragments of MTSP7 protein, which fragments contain the serine protease domain, are used as immunogens for antibody production.

Various procedures known in the art can be used for the production of  
20 polyclonal antibodies to MTSP7 protein, its domains, derivatives, fragments or analogs. For production of the antibody, various host animals can be immunized by injection with the native MTSP7 protein or a synthetic version, or a derivative of the foregoing, such as a cross-linked MTSP7 protein. Such host animals include but are not limited to rabbits, mice and rats. Various adjuvants can be  
25 used to increase the immunological response, depending on the host species, and include but are not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, dinitrophenol, and potentially useful human adjuvants such as bacille Calmette-Guerin (BCG)  
30 and corynebacterium parvum.

For preparation of monoclonal antibodies directed towards an MTSP7 protein or domains, derivatives, fragments or analogs thereof, any technique that

-90-

provides for the production of antibody molecules by continuous cell lines in culture can be used. Such techniques include but are not restricted to the hybridoma technique originally developed by Kohler and Milstein (*Nature* 256:495-497 (1975)), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., *Immunology Today* 4:72 (1983)), and the EBV hybridoma technique to produce human monoclonal antibodies (Cole et al., in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96 (1985)). In an additional embodiment, monoclonal antibodies can be produced in germ-free animals utilizing recent technology (PCT/US90/02545). Human antibodies can be used and can be obtained by using human hybridomas (Cote et al., *Proc. Natl. Acad. Sci. USA* 80:2026-2030 (1983)). Or by transforming human B cells with EBV virus *in vitro* (Cole et al., in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96 (1985)). Techniques developed for the production of "chimeric antibodies" (Morrison et al., *Proc. Natl. Acad. Sci. USA* 81:6851-6855 (1984); Neuberger et al., *Nature* 312:604-608 (1984); Takeda et al., *Nature* 314:452-454 (1985)) by splicing the genes from a mouse antibody molecule specific for the MTSP7 protein together with genes from a human antibody molecule of appropriate biological activity can be used.

Techniques described for the production of single chain antibodies (U.S. patent 4,946,778) can be adapted to produce MTSP7 protein-specific single chain antibodies. An additional embodiment uses the techniques described for the construction of Fab expression libraries (Huse et al., *Science* 246:1275-1281 (1989)) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity for MTSP7 protein or domains, derivatives, or analogs thereof. Non-human antibodies can be "humanized" by known methods (see, e.g., U.S. Patent No. 5,225,539).

Antibody fragments that contain the idiotypes of MTSP7 protein can be generated by techniques known in the art. For example, such fragments include but are not limited to: the F(ab')<sub>2</sub> fragment which can be produced by pepsin digestion of the antibody molecule; the Fab' fragments that can be generated by reducing the disulfide bridges of the F(ab')<sub>2</sub> fragment, the Fab fragments that

-91-

can be generated by treating the antibody molecular with papain and a reducing agent, and Fv fragments.

In the production of antibodies, screening for the desired antibody can be accomplished by techniques known in the art, *e.g.*, ELISA (enzyme-linked  
5 immunosorbent assay). To select antibodies specific to a particular domain of the MTSP7 protein one can assay generated hybridomas for a product that binds to the fragment of the MTSP7 protein that contains such a domain

The foregoing antibodies can be used in methods known in the art relating to the localization and/or quantitation of MTSP7 protein proteins, *e.g.*,  
10 for imaging these proteins, measuring levels thereof in appropriate physiological samples, in, for example, diagnostic methods. In another embodiment, anti-MTSP7 protein antibodies, or fragments thereof, containing the binding domain are used as therapeutic agents.

## 2. Peptides, Polypeptides and Peptide Mimetics

15 Provided herein are methods for identifying molecules that bind to and modulate the activity of MTSP proteins. Included among molecules that bind to MTSP7, particularly the single chain protease domain or catalytically active fragments thereof, are peptides, polypeptides and peptide mimetics, including cyclic peptides. Peptide mimetics are molecules or compounds that mimic the  
20 necessary molecular conformation of a ligand or polypeptide for specific binding to a target molecule such as an MTSP7 protein. In an exemplary embodiment, the peptides, polypeptides and peptide mimetics or peptide mimetics bind to the protease domain of the MTSP7 protein. Such peptides and peptide mimetics include those of antibodies that specifically bind an MTSP7 protein  
25 and, typically, bind to the protease domain of an MTSP7 protein. The peptides, polypeptides and peptide mimetics and peptide mimetics identified by methods provided herein can be agonists or antagonists of MTSP7 proteins.

Such peptides, polypeptides and peptide mimetics and peptide mimetics are useful for diagnosing, treating, preventing, and screening for a disease or  
30 disorder associated with MTSP7 protein activity in a mammal. In addition, the peptides, polypeptides and peptide mimetics are useful for identifying, isolating, and purifying molecules or compounds that modulate the activity of an MTSP7



-92-

protein, or specifically bind to an MTSP7 protein, generally, the protease domain of an MTSP7 protein. Low molecular weight peptides and peptide mimetics can have strong binding properties to a target molecule, *e.g.*, an MTSP7 protein or, generally, to the protease domain of an MTSP7 protein.

5           Peptides, polypeptides and peptide mimetics that bind to MTSP7 proteins as described herein can be administered to mammals, including humans, to modulate MTSP7 protein activity. Thus, methods for therapeutic treatment and prevention of neoplastic diseases comprise administering a peptide, polypeptides or peptide mimetic compound in an amount sufficient to modulate such activity  
10 are provided. Thus, also provided herein are methods for treating a subject having such a disease or disorder in which a peptide, polypeptides or peptide mimetic compound is administered to the subject in a therapeutically effective dose or amount.

          Compositions containing the peptides, polypeptides or peptide mimetics  
15 can be administered for prophylactic and/or therapeutic treatments. In therapeutic applications, compositions can be administered to a patient already suffering from a disease, as described above, in an amount sufficient to cure or at least partially arrest the symptoms of the disease and its complications. Amounts effective for this use will depend on the severity of the disease and the  
20 weight and general state of the patient.

          In prophylactic applications, compositions containing the peptides, polypeptides and peptide mimetics are administered to a patient susceptible to or otherwise at risk of a particular disease. Such an amount is defined to be a "prophylactically effective dose". In this use, the precise amounts again depend  
25 on the patient's state of health and weight.

          Accordingly, the peptides, polypeptides and peptide mimetics that bind to an MTSP7 protein can be used generating pharmaceutical compositions containing, as an active ingredient, at least one of the peptides, polypeptides or peptide mimetics in association with a pharmaceutical carrier or diluent. The  
30 compounds can be administered, for example, by oral, pulmonary, parental (intramuscular, intraperitoneal, intravenous (IV) or subcutaneous injection), inhalation (via a fine powder formulation), transdermal, nasal, vaginal, rectal, or

-93-

sublingual routes of administration and can be formulated in dosage forms appropriate for each route of administration (see, e.g., International PCT application Nos. WO 93/25221 and WO 94/17784; and European Patent Application 613,683).

- 5           Peptides, polypeptides and peptide mimetics that bind to MTSP7 proteins are useful *in vitro* as unique tools for understanding the biological role of MTSP7 proteins, including the evaluation of the many factors thought to influence, and be influenced by, the production of MTSP7 protein. Such peptides, polypeptides and peptide mimetics are also useful in the development of other compounds
- 10   that bind to and modulate the activity of an MTSP7 protein, because such compounds provide important information on the relationship between structure and activity that should facilitate such development.

- The peptides, polypeptides and peptide mimetics are also useful as competitive binders in assays to screen for new MTSP7 proteins or MTSP7
- 15   protein agonists. In such assay embodiments, the compounds can be used without modification or can be modified in a variety of ways; for example, by labeling, such as covalently or non-covalently joining a moiety which directly or indirectly provides a detectable signal. In any of these assays, the materials thereto can be labeled either directly or indirectly. Possibilities for direct labeling
- 20   include label groups such as: radiolabels such as <sup>125</sup>I enzymes (U.S. Pat. No. 3,645,090) such as peroxidase and alkaline phosphatase, and fluorescent labels (U.S. Pat. No. 3,940,475) capable of monitoring the change in fluorescence intensity, wavelength shift, or fluorescence polarization. Possibilities for indirect labeling include biotinylation of one constituent followed by binding to avidin
- 25   coupled to one of the above label groups. The compounds can also include spacers or linkers in cases where the compounds are to be attached to a solid support.

- Moreover, based on their ability to bind to an MTSP7 protein, the peptides, polypeptides and peptide mimetics can be used as reagents for
- 30   detecting MTSP7 proteins in living cells, fixed cells, in biological fluids, in tissue homogenates and in purified, natural biological materials. For example, by labelling such peptides, polypeptides and peptide mimetics, cells having MTSP7

-94-

proteins can be indentified. In addition, based on their ability to bind an MTSP7 protein, the peptides, polypeptides and peptide mimetics can be used in *in situ* staining, FACS (fluorescence-activated cell sorting), Western blotting, ELISA and other analytical protocols. Based on their ability to bind to an MTSP7 protein, the peptides, polypeptides and peptide mimetics can be used in purification of MTSP7 protein polypeptides or in purifying cells expressing the MTSP7 protein polypeptides, *e.g.*, a polypeptide encoding the protease domain of an MTSP7 protein.

The peptides, polypeptides and peptide mimetics can also be used as commercial reagents for various medical research and diagnostic uses. The activity of the peptides and peptide mimetics can be evaluated either *in vitro* or *in vivo* in one of the numerous models described in McDonald (1992) *Am. J. of Pediatric Hematology/Oncology*, 14:8-21.

### 3. Peptide, polypeptides and peptide mimetic therapy

Peptide analogs are commonly used in the pharmaceutical industry as non-peptide drugs with properties analogous to those of the template peptide. These types of non-peptide compounds are termed "peptide mimetics" or "peptidomimetics" (Luthman *et al.*, *A Textbook of Drug Design and Development*, 14:386-406, 2nd Ed., Harwood Academic Publishers (1996); Joachim Grante (1994) *Angew. Chem. Int. Ed. Engl.*, 33:1699-1720; Fauchere (1986) *J. Adv. Drug Res.*, 15:29; Veber and Freidinger (1985) *TINS*, p. 392; and Evans *et al.* (1987) *J. Med. Chem.* 30:1229). Peptide mimetics that are structurally similar to therapeutically useful peptides can be used to produce an equivalent or enhanced therapeutic or prophylactic effect. Preparation of peptidomimetics and structures thereof are known to those of skill in this art.

Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (*e.g.*, D-lysine in place of L-lysine) can be used to generate more stable peptides. In addition, constrained peptides containing a consensus sequence or a substantially identical consensus sequence variation can be generated by methods known in the art (Rizo *et al.* (1992) *An. Rev. Biochem.*, 61:387, incorporated herein by reference); for

-95-

example, by adding internal cysteine residues capable of forming intramolecular disulfide bridges which cyclize the peptide.

Those skilled in the art would appreciate that modifications can be made to the peptides and mimetics without deleteriously effecting the biological or functional activity of the peptide. Further, the skilled artisan would know how to design non-peptide structures in three dimensional terms, that mimic the peptides that bind to a target molecule, *e.g.*, an MTSP7 protein or, generally, the protease domain of MTSP7 proteins (see, *e.g.*, Eck and Sprang (1989) *J. Biol. Chem.*, 26: 17605-18795).

When used for diagnostic purposes, the peptides and peptide mimetics can be labeled with a detectable label and, accordingly, the peptides and peptide mimetics without such a label can serve as intermediates in the preparation of labeled peptides and peptide mimetics. Detectable labels can be molecules or compounds, which when covalently attached to the peptides and peptide mimetics, permit detection of the peptide and peptide mimetics *in vivo*, for example, in a patient to whom the peptide or peptide mimetic has been administered, or *in vitro*, *e.g.*, in a sample or cells. Suitable detectable labels are well known in the art and include, by way of example, radioisotopes, fluorescent labels (*e.g.*, fluorescein), and the like. The particular detectable label employed is not critical and is selected relative to the amount of label to be employed as well as the toxicity of the label at the amount of label employed. Selection of the label relative to such factors is well within the skill of the art.

Covalent attachment of a detectable label to the peptide or peptide mimetic is accomplished by conventional methods well known in the art. For example, when the  $^{125}\text{I}$  radioisotope is employed as the detectable label, covalent attachment of  $^{125}\text{I}$  to the peptide or the peptide mimetic can be achieved by incorporating the amino acid tyrosine into the peptide or peptide mimetic and then iodinating the peptide (see, *e.g.*, Weaner *et al.* (1994) *Synthesis and Applications of Isotopically Labelled Compounds*, pp. 137-140). If tyrosine is not present in the peptide or peptide mimetic, incorporation of tyrosine to the N or C terminus of the peptide or peptide mimetic can be achieved by well known chemistry. Likewise,  $^{32}\text{P}$  can be incorporated onto the peptide or peptide

-96-

mimetic as a phosphate moiety through, for example, a hydroxyl group on the peptide or peptide mimetic using conventional chemistry.

Labeling of peptidomimetics usually involves covalent attachment of one or more labels, directly or through a spacer (*e.g.*, an amide group), to  
5 non-interfering position(s) on the peptidomimetic that are predicted by quantitative structure-activity data and/or molecular modeling. Such non-interfering positions generally are positions that do not form direct contacts with the macromolecule(s) to which the peptidomimetic binds to produce the therapeutic effect. Derivatization (*e.g.*, labeling) of peptidomimetics should not  
10 substantially interfere with the desired biological or pharmacological activity of the peptidomimetic.

Peptides, polypeptides and peptide mimetics that can bind to MTSP7 proteins or the protease domain of MTSP7 proteins and modulate the activity thereof, or have MTSP7 protein activity, can be used for treatment of neoplastic  
15 diseases. The peptides, polypeptides and peptide mimetics can be delivered, *in vivo* or *ex vivo*, to the cells of a subject in need of treatment. Further, peptides which have MTSP7 protein activity can be delivered, *in vivo* or *ex vivo*, to cells which carry mutant or missing alleles encoding the MTSP7 protein gene. Any of the techniques described herein or known to the skilled artisan can be used for  
20 preparation and *in vivo* or *ex vivo* delivery of such peptides, polypeptides and peptide mimetics that are substantially free of other human proteins. For example, the peptides, polypeptides and peptide mimetics can be readily prepared by expression in a microorganism or synthesis *in vitro*.

The peptides or peptide mimetics can be introduced into cells, *in vivo* or  
25 *ex vivo*, by microinjection or by use of liposomes, for example. Alternatively, the peptides, polypeptides or peptide mimetics can be taken up by cells, *in vivo* or *ex vivo*, actively or by diffusion. In addition, extracellular application of the peptide, polypeptides or peptide mimetic can be sufficient to effect treatment of a neoplastic disease. Other molecules, such as drugs or organic compounds,  
30 that: 1) bind to an MTSP7 protein or protease domain thereof; or 2) have a similar function or activity to an MTSP7 protein or protease domain thereof, can be used in methods for treatment.

#### 4. Rational drug design

The goal of rational drug design is to produce structural analogs of biologically active polypeptides or peptides of interest or of small molecules or peptide mimetics with which they interact (e.g., agonists, antagonists, inhibitors) in order to fashion drugs which are, e.g., more active or stable forms thereof; or which, for example, enhance or interfere with the function of a polypeptide *in vivo* (e.g., an MTSP7 protein). In one approach, one first determines the three-dimensional structure of a protein of interest (e.g., an MTSP7 protein or polypeptide having a protease domain) or, for example, of a MTSP7 protein-ligand complex, by X-ray crystallography, by computer modeling or most typically, by a combination of approaches (see, e.g., Erickson *et al.* 1990). Also, useful information regarding the structure of a polypeptide can be gained by modeling based on the structure of homologous proteins. In addition, peptides can be analyzed by an alanine scan. In this technique, an amino acid residue is replaced by Ala, and its effect on the peptide's activity is determined. Each of the amino acid residues of the peptide is analyzed in this manner to determine the important regions of the peptide.

Also, a polypeptide or peptide that binds to an MTSP7 protein or, generally, the protease domain of an MTSP7 protein, can be selected by a functional assay, and then the crystal structure of this polypeptide or peptide can be determined. The polypeptide can be, for example, an antibody specific for an MTSP7 protein or the protein domain of an MTSP7 protein. This approach can yield a pharmacore upon which subsequent drug design can be based. Further, it is possible to bypass the crystallography altogether by generating anti-idiotypic polypeptides or peptides, (anti-ids) to a functional, pharmacologically active polypeptide or peptide that binds to an MTSP7 protein or protease domain of an MTSP7 protein. As a mirror image of a mirror image, the binding site of the anti-ids is expected to be an analog of the original target molecule, e.g., an MTSP7 protein or polypeptide having an MTSP7 protein. The anti-id could then be used to identify and isolate peptides from banks of chemically or biologically produced banks of peptides. Selected peptides would then act as the pharmacore.

-98-

Thus, one can design drugs which have, e.g., improved activity or stability or which act as modulators (e.g., inhibitors, agonists, antagonists) of MTSP7 protein activity, and are useful in the methods, particularly the methods for diagnosis, treatment, prevention, and screening of a neoplastic disease. By virtue of the availability of cloned MTSP7 protein sequences, sufficient amounts of the MTSP7 protein polypeptide can be made available to perform such analytical studies as X-ray crystallography. In addition, the knowledge of the amino acid sequence of an MTSP7 protein or the protease domain thereof, e.g., the protease domain encoded by the amino acid sequence of SEQ ID NO: 2, can provide guidance on computer modeling techniques in place of, or in addition to, X-ray crystallography.

**Methods of identifying peptides and peptide mimetics that bind to MTSP7 proteins**

Peptides having a binding affinity to the MTSP7 protein polypeptides provided herein (e.g., an MTSP7 protein or a polypeptide having a protease domain of an MTSP7 protein) can be readily identified, for example, by random peptide diversity generating systems coupled with an affinity enrichment process. Specifically, random peptide diversity generating systems include the "peptides on plasmids" system (see, e.g., U.S. Patent Nos. 5,270,170 and 5,338,665); the "peptides on phage" system (see, e.g., U.S. Patent No. 6,121,238 and Cwirla, *et al.* (1990) *Proc. Natl. Acad. Sci. U.S.A.* 87:6378-6382); the "polysome system;" the "encoded synthetic library (ESL)" system; and the "very large scale immobilized polymer synthesis" system (see, e.g., U.S. Patent No. 6,121,238; and Dower *et al.* (1991) *An. Rep. Med. Chem.* 26:271-280).

For example, using the procedures described above, random peptides can generally be designed to have a defined number of amino acid residues in length (e.g., 12). To generate the collection of oligonucleotides encoding the random peptides, the codon motif (NNK)<sub>x</sub>, where N is nucleotide A, C, G, or T (equimolar; depending on the methodology employed, other nucleotides can be employed), K is G or T (equimolar), and x is an integer corresponding to the number of amino acids in the peptide (e.g., 12) can be used to specify any one

-99-

of the 32 possible codons resulting from the NNK motif: 1 for each of 12 amino acids, 2 for each of 5 amino acids, 3 for each of 3 amino acids, and only one of the three stop codons. Thus, the NNK motif encodes all of the amino acids, encodes only one stop codon, and reduces codon bias.

- 5           The random peptides can be presented, for example, either on the surface of a phage particle, as part of a fusion protein containing either the pIII or the pVIII coat protein of a phage fd derivative (peptides on phage) or as a fusion protein with the LacI peptide fusion protein bound to a plasmid (peptides on plasmids). The phage or plasmids, including the DNA encoding the peptides, can
- 10 be identified and isolated by an affinity enrichment process using immobilized MTSP7 protein polypeptide having a protease domain. The affinity enrichment process, sometimes called "panning," typically involves multiple rounds of incubating the phage, plasmids, or polysomes with the immobilized MTSP7 protein polypeptide, collecting the phage, plasmids, or polysomes that bind to
- 15 the MTSP7 protein polypeptide (along with the accompanying DNA or mRNA), and producing more of the phage or plasmids (along with the accompanying LacI-peptide fusion protein) collected.

#### **Characteristics of peptides and peptide mimetics**

- Among the peptides, polypeptides and peptide mimetics for therapeutic
- 20 application are those of having molecular weights from about 250 to about 8,000 daltons. If such peptides are oligomerized, dimerized and/or derivatized with a hydrophilic polymer (*e.g.*, to increase the affinity and/or activity of the compounds), the molecular weights of such peptides can be substantially greater and can range anywhere from about 500 to about 120,000 daltons, generally
- 25 from about 8,000 to about 80,000 daltons. Such peptides can contain 9 or more amino acids that are naturally occurring or synthetic (non-naturally occurring) amino acids. One skilled in the art can determine the affinity and molecular weight of the peptides and peptide mimetics suitable for therapeutic and/or diagnostic purposes (*e.g.*, see Dower *et al.*, U.S. Patent No. 6,121,238).
- 30           The peptides can be covalently attached to one or more of a variety of hydrophilic polymers. Suitable hydrophilic polymers include, but are not limited to, polyalkylethers as exemplified by polyethylene glycol and polypropylene



-100-

glycol, polylactic acid, polyglycolic acid, polyoxyalkenes, polyvinylalcohol, polyvinylpyrrolidone, cellulose and cellulose derivatives, dextran and dextran derivatives. When the peptide compounds are derivatized with such polymers, their solubility and circulation half-lives can be increased with little, if any,

5   diminishment in their binding activity. The peptide compounds can be dimerized and each of the dimeric subunits can be covalently attached to a hydrophilic polymer. The peptide compounds can be PEGylated, i.e., covalently attached to polyethylene glycol (PEG).

#### 6.     Methods of preparing peptides and peptide mimetics

10       Peptides that bind to MTSP7 proteins can be prepared by classical methods known in the art, for example, by using standard solid phase techniques. The standard methods include exclusive solid phase synthesis, partial solid phase synthesis methods, fragment condensation, classical solution synthesis, and even by recombinant DNA technology (see, *e.g.*, Merrifield  
15   (1963) *J. Am. Chem. Soc.*, 85:2149, incorporated herein by reference.)

      Using the "encoded synthetic library" or "very large scale immobilized polymer synthesis" systems (see, *e.g.*, U.S. Patent No. 5,925,525, and 5,902,723); the minimum size of a peptide with the activity of interest can be determined. In addition all peptides that form the group of peptides that differ  
20   from the desired motif (or the minimum size of that motif) in one, two, or more residues can be prepared. This collection of peptides then can be screened ability to bind to the target molecule, *e.g.*, and MTSP7 protein or, generally, the protease domain of an MTSP7 protein. This immobilized polymer synthesis system or other peptide synthesis methods can also be used to synthesize  
25   truncation analogs and deletion analogs and combinations of truncation and deletion analogs of the peptide compounds.

      These procedures can also be used to synthesize peptides in which amino acids other than the 20 naturally occurring, genetically encoded amino acids are substituted at one, two, or more positions of the peptide. For instance,  
30   naphthylalanine can be substituted for tryptophan, facilitating synthesis. Other synthetic amino acids that can be substituted into the peptides include L-hydroxypropyl, L-3, 4-dihydroxy-phenylalanyl, D amino acids such as

-101-

L-d-hydroxylysyl and D-d-methylalanyl, L- $\alpha$ -methylalanyl,  $\beta$  amino acids, and isoquinolyl. D amino acids and non-naturally occurring synthetic amino acids can also be incorporated into the peptides (see, e.g., Roberts *et al.* (1983) *Unusual Amino/Acids in Peptide Synthesis*, 5(6):341-449).

5           The peptides can also be modified by phosphorylation (see, e.g., W. Bannwarth *et al.* (1996) *Biorganic and Medicinal Chemistry Letters*, 6(17):2141-2146), and other methods for making peptide derivatives (see, e.g., Hruby *et al.* (1990) *Biochem. J.*, 268(2):249-262). Thus, peptide compounds also serve as a basis to prepare peptide mimetics with similar biological activity.

10           Those of skill in the art recognize that a variety of techniques are available for constructing peptide mimetics with the same or similar desired biological activity as the corresponding peptide compound but with more favorable activity than the peptide with respect to solubility, stability, and susceptibility to hydrolysis and proteolysis (see, e.g., Morgan *et al.* (1989) *An. Rep. Med. Chem.*, 24:243-252). Methods for preparing peptide mimetics modified at the N-terminal amino group, the C-terminal carboxyl group, and/or changing one or more of the amido linkages in the peptide to a non-amido linkage are known to those of skill in the art.

20           Amino terminus modifications include, but are not limited to, alkylating, acetylating and adding a carbobenzoyl group, forming a succinimide group (see, e.g., Murray *et al.* (1995) *Burger's Medicinal Chemistry and Drug Discovery*, 5th ed., Vol. 1, Manfred E. Wolf, ed., John Wiley and Sons, Inc.). C-terminal modifications include mimetics wherein the C-terminal carboxyl group is replaced by an ester, an amide or modifications to form a cyclic peptide.

25           In addition to N-terminal and C-terminal modifications, the peptide compounds, including peptide mimetics, can advantageously be modified with or covalently coupled to one or more of a variety of hydrophilic polymers. It has been found that when peptide compounds are derivatized with a hydrophilic polymer, their solubility and circulation half-lives can be increased and their  
30 immunogenicity is masked, with little, if any, diminishment in their binding activity. Suitable nonproteinaceous polymers include, but are not limited to, polyalkylethers as exemplified by polyethylene glycol and polypropylene glycol,

-102-

polylactic acid, polyglycolic acid, polyoxyalkenes, polyvinylalcohol, polyvinylpyrrolidone, cellulose and cellulose derivatives, dextran and dextran derivatives. Generally, such hydrophilic polymers have an average molecular weight ranging from about 500 to about 100,000 daltons, including from about 2,000 to about 40,000 daltons and, from about 5,000 to about 20,000 daltons. The hydrophilic polymers also can have an average molecular weights of about 5,000 daltons, 10,000 daltons and 20,000 daltons.

Methods for derivatizing peptide compounds or for coupling peptides to such polymers have been described (see, *e.g.*, Zallipsky (1995) *Bioconjugate Chem.*, 6:150-165; Monfardini *et al.* (1995) *Bioconjugate Chem.*, 6:62-69; U.S. Pat. No. 4,640,835; U.S. Pat. No. 4,496,689; U.S. Pat. No. 4,301,144; U.S. Pat. No. 4,670,417; U.S. Pat. No. 4,791,192; U.S. Pat. No. 4,179,337 and WO 95/34326, all of which are incorporated by reference in their entirety herein).

Other methods for making peptide derivatives are described, for example, in Hruby *et al.* (1990), *Biochem J.*, 268(2):249-262, which is incorporated herein by reference. Thus, the peptide compounds also serve as structural models for non-peptidic compounds with similar biological activity. Those of skill in the art recognize that a variety of techniques are available for constructing compounds with the same or similar desired biological activity as a particular peptide compound but with more favorable activity with respect to solubility, stability, and susceptibility to hydrolysis and proteolysis (see, *e.g.*, Morgan *et al.* (1989) *An. Rep. Med. Chem.*, 24:243-252, incorporated herein by reference). These techniques include replacing the peptide backbone with a backbone composed of phosphonates, amidates, carbamates, sulfonamides, secondary amines, and N-methylamino acids.

Peptide compounds can exist in a cyclized form with an intramolecular disulfide bond between the thiol groups of the cysteines. Alternatively, an intermolecular disulfide bond between the thiol groups of the cysteines can be produced to yield a dimeric (or higher oligomeric) compound. One or more of the cysteine residues can also be substituted with a homocysteine.

-103-

**I. CONJUGATES**

A conjugate, containing: a) a single chain protease domain (or proteolytically active portion thereof) of an MTSP7 protein or a full length zymogen, activated form thereof, or two or single chain protease domain thereof; and b) a targeting agent linked to the MTSP7 protein directly or via a linker, wherein the agent facilitates: i) affinity isolation or purification of the conjugate; ii) attachment of the conjugate to a surface; iii) detection of the conjugate; or iv) targeted delivery to a selected tissue or cell, is provided herein. The conjugate can be a chemical conjugate or a fusion protein mixture thereof.

10       The targeting agent is generally a protein or peptide fragment, such as a tissue specific or tumor specific monoclonal antibody or growth factor or fragment thereof linked either directly or via a linker to an MTSP7 protein or a protease domain thereof. The targeting agent can also be a protein or peptide fragment that contains a protein binding sequence, a nucleic acid binding  
15       sequence, a lipid binding sequence, a polysaccharide binding sequence, or a metal binding sequence, or a linker for attachment to a solid support. In a particular embodiment, the conjugate contains a) the MTSP7 or portion thereof, as described herein; and b) a targeting agent linked to the MTSP7 protein directly or via a linker.

20       Conjugates, such as fusion proteins and chemical conjugates, of the MTSP7 protein with a protein or peptide fragment (or plurality thereof) that functions, for example, to facilitate affinity isolation or purification of the MTSP7 protein domain, attachment of the MTSP7 protein domain to a surface, or detection of the MTSP7 protein domain are provided. The conjugates can be  
25       produced by chemical conjugation, such as via thiol linkages, but are generally produced by recombinant means as fusion proteins. In the fusion protein, the peptide or fragment thereof is linked to either the N-terminus or C-terminus of the MTSP7 protein domain. In chemical conjugates the peptide or fragment thereof can be linked anywhere that conjugation can be effected, and there can  
30       be a plurality of such peptides or fragments linked to a single MTSP7 protein domain or to a plurality thereof.

-104-

The targeting agent is for *in vitro* or *in vivo* delivery to a cell or tissue, and includes agents such as cell or tissue-specific antibodies, growth factors and other factors that bind to moieties expressed on specific cells; and other cell or tissue specific agents that promote directed delivery of a linked protein.

- 5 Generally the targeting agent is one that specifically delivers the MTSP7 protein to selected cells by interaction with a cell surface protein and internalization of conjugate or MTSP7 protein portion thereof.

- These conjugates are used in a variety of methods and are particularly suited for use in methods of activation of prodrugs, such as prodrugs that, upon  
10 cleavage by the particular MTSP7 protein are cytotoxic. The prodrugs are administered prior to simultaneously with or subsequently to the conjugate. Upon delivery to the targeted cells, the protease activates the prodrug, which then exhibits is therapeutic effect, such as a cytotoxic effect.

1. Conjugation

- 15 Conjugates with linked MTSP7 protein domains can be prepared either by chemical conjugation, recombinant DNA technology, or combinations of recombinant expression and chemical conjugation. The MTSP7 protein domains and the targeting agent can be linked in any orientation and more than one targeting agents and/or MTSP7 protein domains can be present in a conjugate.

20 a. Fusion proteins

Fusion proteins are proved herein. A fusion protein contains: a) one or a plurality of domains of an MTSP7 proteins and b) a targeting agent. The fusion proteins are generally produced by recombinant expression of nucleic acids that encode the fusion protein.

25 b. Chemical conjugation

- To effect chemical conjugation herein, the MTSP7 protein domain is linked via one or more selected linkers or directly to the targeting agent. Chemical conjugation must be used if the targeted agent is other than a peptide or protein, such a nucleic acid or a non-peptide drug. Any means known to  
30 those of skill in the art for chemically conjugating selected moieties can be used.

-105-

## 2. Linkers

Linkers for two purposes are contemplated herein. The conjugates can include one or more linkers between the MTSP7 protein portion and the targeting agent. Additionally, linkers are used for facilitating or enhancing immobilization of an MTSP7 protein or portion thereof on a solid support, such as a microtiter plate, silicon or silicon-coated chip, glass or plastic support, such as for high throughput solid phase screening protocols.

Any linker known to those of skill in the art for preparation of conjugates can be used herein. These linkers are typically used in the preparation of chemical conjugates; peptide linkers can be incorporated into fusion proteins.

Linkers can be any moiety suitable to associate a domain of MTSP7 protein and a targeting agent. Such linkers and linkages include, but are not limited to, peptidic linkages, amino acid and peptide linkages, typically containing between one and about 60 amino acids, more generally between about 10 and 30 amino acids, chemical linkers, such as heterobifunctional cleavable cross-linkers, including but are not limited to, N-succinimidyl (4-iodoacetyl)-aminobenzoate, sulfosuccinimidyl (4-iodoacetyl)-aminobenzoate, 4-succinimidyl-oxycarbonyl- $\alpha$ -(2-pyridyldithio)toluene, sulfosuccinimidyl-6-[ $\alpha$ -methyl- $\alpha$ -(pyridyldithio)-toluamido] hexanoate, N-succinimidyl-3-(2-pyridyldithio) - propionate, succinimidyl 6[3-(2-pyridyldithio)-propionamido] hexanoate, sulfosuccinimidyl 6[3-(2-pyridyldithio)-propionamido] hexanoate, 3-(2-pyridyldithio)-propionyl hydrazide, Ellman's reagent, dichlorotriazinic acid, and S-(2-thiopyridyl)-L-cysteine. Other linkers include, but are not limited to peptides and other moieties that reduce steric hindrance between the domain of MTSP7 protein and the targeting agent, intracellular enzyme substrates, linkers that increase the flexibility of the conjugate, linkers that increase the solubility of the conjugate, linkers that increase the serum stability of the conjugate, photocleavable linkers and acid cleavable linkers.

-106-

Other exemplary linkers and linkages that are suitable for chemically linked conjugates include, but are not limited to, disulfide bonds, thioether bonds, hindered disulfide bonds, and covalent bonds between free reactive groups, such as amine and thiol groups. These bonds are produced using  
5 heterobifunctional reagents to produce reactive thiol groups on one or both of the polypeptides and then reacting the thiol groups on one polypeptide with reactive thiol groups or amine groups to which reactive maleimido groups or thiol groups can be attached on the other. Other linkers include, acid cleavable linkers, such as bismaleimideoxy propane, acid labile-transferrin conjugates  
10 and adipic acid dihydrazide, that would be cleaved in more acidic intracellular compartments; cross linkers that are cleaved upon exposure to UV or visible light and linkers, such as the various domains, such as C<sub>H</sub>1, C<sub>H</sub>2, and C<sub>H</sub>3, from the constant region of human IgG<sub>1</sub> (see, Batra *et al. Molecular Immunol.*, 30:379-386 (1993)). In some embodiments, several linkers can be included in  
15 order to take advantage of desired properties of each linker.

Chemical linkers and peptide linkers can be inserted by covalently coupling the linker to the domain of MTSP7 protein and the targeting agent. The heterobifunctional agents, described below, can be used to effect such covalent coupling. Peptide linkers can also be linked by expressing DNA encoding the  
20 linker and TA, linker and targeted agent, or linker, targeted agent and TA as a fusion protein. Flexible linkers and linkers that increase solubility of the conjugates are contemplated for use, either alone or with other linkers are also contemplated herein.

**a) Acid cleavable, photocleavable and heat sensitive linkers**

25 Acid cleavable linkers, photocleavable and heat sensitive linkers can also be used, particularly where it can be necessary to cleave the domain of MTSP7 protein to permit it to be more readily accessible to reaction. Acid cleavable linkers include, but are not limited to, bismaleimideoxy propane; and adipic acid dihydrazide linkers (see, *e.g.*, Fattom *et al.* (1992) *Infection & Immun.*  
30 60:584-589) and acid labile transferrin conjugates that contain a sufficient portion of transferrin to permit entry into the intracellular transferrin cycling pathway (see, *e.g.*, Welhöner *et al.* (1991) *J. Biol. Chem.* 266:4309-4314).

-107-

Photocleavable linkers are linkers that are cleaved upon exposure to light (see, e.g., Goldmacher *et al.* (1992) *Bioconj. Chem.* 3:104-107, which linkers are herein incorporated by reference), thereby releasing the targeted agent upon exposure to light. Photocleavable linkers that are cleaved upon exposure to light are known (see, e.g., Hazum *et al.* (1981) in *Pept., Proc. Eur. Pept. Symp.*, 16th, Brunfeldt, K (Ed), pp. 105-110, which describes the use of a nitrobenzyl group as a photocleavable protective group for cysteine; Yen *et al.* (1989) *Makromol. Chem* 190:69-82, which describes water soluble photocleavable copolymers, including hydroxypropylmethacrylamide copolymer, glycine copolymer, fluorescein copolymer and methylrhodamine copolymer; Goldmacher *et al.* (1992) *Bioconj. Chem.* 3:104-107, which describes a cross-linker and reagent that undergoes photolytic degradation upon exposure to near UV light (350 nm); and Senter *et al.* (1985) *Photochem. Photobiol* 42:231-237, which describes nitrobenzyloxycarbonyl chloride cross linking reagents that produce photocleavable linkages), thereby releasing the targeted agent upon exposure to light. Such linkers would have particular use in treating dermatological or ophthalmic conditions that can be exposed to light using fiber optics. After administration of the conjugate, the eye or skin or other body part can be exposed to light, resulting in release of the targeted moiety from the conjugate. Such photocleavable linkers are useful in connection with diagnostic protocols in which it is desirable to remove the targeting agent to permit rapid clearance from the body of the animal.

**b) Other linkers for chemical conjugation**

Other linkers, include trityl linkers, particularly, derivatized trityl groups to generate a genus of conjugates that provide for release of therapeutic agents at various degrees of acidity or alkalinity. The flexibility thus afforded by the ability to preselect the pH range at which the therapeutic agent will be released allows selection of a linker based on the known physiological differences between tissues in need of delivery of a therapeutic agent (see, e.g., U.S. Patent No. 5,612,474). For example, the acidity of tumor tissues appears to be lower than that of normal tissues.



-108-

**c) Peptide linkers**

The linker moieties can be peptides. Peptide linkers can be employed in fusion proteins and also in chemically linked conjugates. The peptide typically has from about 2 to about 60 amino acid residues, for example from about 5 to about 40, or from about 10 to about 30 amino acid residues. The length selected will depend upon factors, such as the use for which the linker is included.

Peptide linkers are advantageous when the targeting agent is proteinaceous. For example, the linker moiety can be a flexible spacer amino acid sequence, such as those known in single-chain antibody research. Examples of such known linker moieties include, but are not limited to, peptides, such as  $(\text{Gly}_m\text{Ser})_n$  and  $(\text{Ser}_m\text{Gly})_n$ , in which  $n$  is 1 to 6, generally 1 to 4 or 2 to 4, and  $m$  is 1 to 6, generally 1 to 4 or 2 to 4, enzyme cleavable linkers and others.

Additional linking moieties are known (see, *e.g.*, Huston *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 85:5879-5883, 1988; Whitlow, M., *et al.*, *Protein Engineering* 6:989-995, 1993; Newton *et al.*, *Biochemistry* 35:545-553, 1996; A. J. Cumber *et al.*, *Bioconj. Chem.* 3:397-401, 1992; Ladurner *et al.*, *J. Mol. Biol.* 273:330-337, 1997; and U.S. Patent. No. 4,894,443). In some embodiments, several linkers can be included in order to take advantage of desired properties of each linker.

**3. Targeting agents**

Any agent that facilitates detection, immobilization, or purification of the conjugate is contemplated for use herein. For chemical conjugates any moiety that has such properties is contemplated; for fusion proteins, the targeting agent is a protein, peptide or fragment thereof that sufficient to effects the targeting activity. Generally targeting agents are those that deliver the MTSP7 protein or portion thereof to selected cells and tissues. Such agents include tumor specific monoclonal antibodies and portions thereof, growth factors, such as FGF, EGF, PDGF, VEGF, cytokines, including chemokines, and other such agents.

-109-

#### 4. Nucleic acids, plasmids and cells

Isolated nucleic acid fragments encoding fusion proteins are provided.

The nucleic acid fragment that encodes the fusion protein includes: a) nucleic acid encoding a protease domain of an MTSP7 protein; and b) nucleic acid

5 encoding a protein, peptide or effective fragment thereof that facilitates: i) affinity isolation or purification of the fusion protein; ii) attachment of the fusion protein to a surface; or iii) detection of the fusion protein. Generally, the nucleic acid is DNA.

Plasmids for replication and vectors for expression that contain the above  
10 nucleic acid fragments are also provided. Cells containing the plasmids and vectors are also provided. The cells can be any suitable host including, but are not limited to, bacterial cells, yeast cells, fungal cells, plant cells, insect cell and animal cells. The nucleic acids, plasmids, and cells containing the plasmids can be prepared according to methods known in the art including any described  
15 herein.

Also provided are methods for producing the above fusion proteins. An exemplary method includes the steps of growing, i.e. culturing the cells so that the proliferate, cells containing a plasmid encoding the fusion protein under conditions whereby the fusion protein is expressed by the cell, and recovering  
20 the expressed fusion protein. Methods for expressing and recovering recombinant proteins are well known in the art (*See generally, Current Protocols in Molecular Biology* (1998) § 16, John Wiley & Sons, Inc.) and such methods can be used for expressing and recovering the expressed fusion proteins. Typically, the recombinant expression and recovery methods described herein can be used.

25 The recovered fusion proteins can be isolated or purified by methods known in the art such as centrifugation, filtration, chromatograph, electrophoresis, immunoprecipitation, or by a combination thereof (*See generally, Current Protocols in Molecular Biology* (1998) § 10, John Wiley & Sons, Inc.). For example, the recovered fusion protein is isolated or purified through affinity  
30 binding between the protein or peptide fragment of the fusion protein and an affinity binding moiety. As discussed in the above sections regarding the construction of the fusion proteins, any affinity binding pairs can be constructed

-110-

and used in the isolation or purification of the fusion proteins. For example, the affinity binding pairs can be protein binding sequences/protein, DNA binding sequences/DNA sequences, RNA binding sequences/RNA sequences, lipid binding sequences/lipid, polysaccharide binding sequences/polysaccharide, or  
5 metal binding sequences/metal.

#### **5. Immobilization and supports or substrates therefor**

In certain embodiments, where the targeting agents are designed for linkage to surfaces, the MTSP7 protein can be attached by linkage such as ionic or covalent, non-covalent or other chemical interaction, to a surface of a support  
10 or matrix material. Immobilization can be effected directly or via a linker. The MTSP7 protein can be immobilized on any suitable support, including, but are not limited to, silicon chips, and other supports described herein and known to those of skill in the art. A plurality of MTSP7 protein or protease domains thereof can be attached to a support, such as an array (*i.e.*, a pattern of two or  
15 more) of conjugates on the surface of a silicon chip or other chip for use in high throughput protocols and formats.

It is also noted that the domains of the MTSP7 protein can be linked directly to the surface or via a linker without a targeting agent linked thereto. Hence chips containing arrays of the domains of the MTSP7 protein.

20 The matrix material or solid supports contemplated herein are generally any of the insoluble materials known to those of skill in the art to immobilize ligands and other molecules, and are those that used in many chemical syntheses and separations. Such supports are used, for example, in affinity chromatography, in the immobilization of biologically active materials, and during  
25 chemical syntheses of biomolecules, including proteins, amino acids and other organic molecules and polymers. The preparation of and use of supports is well known to those of skill in this art; there are many such materials and preparations thereof known. For example, naturally-occurring support materials, such as agarose and cellulose, can be isolated from their respective sources, and  
30 processed according to known protocols, and synthetic materials can be prepared in accord with known protocols.

-111-

The supports are typically insoluble materials that are solid, porous, deformable, or hard, and have any required structure and geometry, including, but not limited to: beads, pellets, disks, capillaries, hollow fibers, needles, solid fibers, random shapes, thin films and membranes. Thus, the item can be  
5 fabricated from the matrix material or combined with it, such as by coating all or part of the surface or impregnating particles.

Typically, when the matrix is particulate, the particles are at least about 10-2000  $\mu$ M, but can be smaller or larger, depending upon the selected application. Selection of the matrices will be governed, at least in part, by their  
10 physical and chemical properties, such as solubility, functional groups, mechanical stability, surface area swelling propensity, hydrophobic or hydrophilic properties and intended use.

If necessary, the support matrix material can be treated to contain an appropriate reactive moiety. In some cases, the support matrix material already  
15 containing the reactive moiety can be obtained commercially. The support matrix material containing the reactive moiety can thereby serve as the matrix support upon which molecules are linked. Materials containing reactive surface moieties such as amino silane linkages, hydroxyl linkages or carboxysilane linkages can be produced by well established surface chemistry techniques  
20 involving silanization reactions, or the like. Examples of these materials are those having surface silicon oxide moieties, covalently linked to gamma-amino-propylsilane, and other organic moieties; N-[3-(triethoxysilyl)propyl]phthelamic acid; and bis-(2-hydroxyethyl)aminopropyltriethoxysilane. Exemplary of readily available materials containing amino group reactive functionalities, include, but  
25 are not limited to, para-aminophenyltriethoxysilane. Also derivatized polystyrenes and other such polymers are well known and readily available to those of skill in this art (*e.g.*, the Tentagel<sup>®</sup> Resins are available with a multitude of functional groups, and are sold by Rapp Polymere, Tubingen, Germany; see, U.S. Patent No. 4,908,405 and U.S. Patent No. 5,292,814; *see, also* Butz et al.,  
30 *Peptide Res.*, 7:20-23 (1994); and Kleine et al., *Immunobiol.*, 190:53-66 (1994)).

-112-

These matrix materials include any material that can act as a support matrix for attachment of the molecules of interest. Such materials are known to those of skill in this art, and include those that are used as a support matrix. These materials include, but are not limited to, inorganics, natural polymers, and  
5 synthetic polymers, including, but are not limited to: cellulose, cellulose derivatives, acrylic resins, glass, silica gels, polystyrene, gelatin, polyvinyl pyrrolidone, co-polymers of vinyl and acrylamide, polystyrene cross-linked with divinylbenzene and others (*see*, Merrifield, *Biochemistry*, 3:1385-1390 (1964)), polyacrylamides, latex gels, polystyrene, dextran, polyacrylamides, rubber,  
10 silicon, plastics, nitrocellulose, celluloses, natural sponges. Of particular interest herein, are highly porous glasses (*see, e.g.*, U.S. Patent No. 4,244,721) and others prepared by mixing a borosilicate, alcohol and water.

Synthetic supports include, but are not limited to: acrylamides, dextran-derivatives and dextran co-polymers, agarose-polyacrylamide blends, other  
15 polymers and co-polymers with various functional groups, methacrylate derivatives and co-polymers, polystyrene and polystyrene copolymers (*see, e.g.*, Merrifield, *Biochemistry*, 3:1385-1390 (1964); Berg et al., in *Innovation Perspect. Solid Phase Synth. Collect. Pap.*, Int. Symp., 1st, Epton, Roger (Ed), pp. 453-459 (1990); Berg et al., *Pept., Proc. Eur. Pept. Symp.*, 20th, Jung, G.  
20 *et al.* (Eds), pp. 196-198 (1989); Berg et al., *J. Am. Chem. Soc.*, 111:8024-8026 (1989); Kent et al., *Isr. J. Chem.*, 17:243-247 (1979); Kent et al., *J. Org. Chem.*, 43:2845-2852 (1978); Mitchell et al., *Tetrahedron Lett.*, 42:3795-3798 (1976); U.S. Patent No. 4,507,230; U.S. Patent No. 4,006,117; and U.S. Patent No. 5,389,449). Such materials include those made from  
25 polymers and co-polymers such as polyvinylalcohols, acrylates and acrylic acids such as polyethylene-co-acrylic acid, polyethylene-co-methacrylic acid, polyethylene-co-ethylacrylate, polyethylene-co-methyl acrylate, polypropylene-co-acrylic acid, polypropylene-co-methyl-acrylic acid, polypropylene-co-ethylacrylate, polypropylene-co-methyl acrylate, polyethylene-co-vinyl acetate, poly-  
30 propylene-co-vinyl acetate, and those containing acid anhydride groups such as polyethylene-co-maleic anhydride and polypropylene-co-maleic anhydride.

-113-

Liposomes have also been used as solid supports for affinity purifications (Powell et al. *Biotechnol. Bioeng.*, 33:173 (1989)).

Numerous methods have been developed for the immobilization of proteins and other biomolecules onto solid or liquid supports (*see, e.g.,*

- 5 Mosbach, *Methods in Enzymology*, 44 (1976); Weetall, *Immobilized Enzymes, Antigens, Antibodies, and Peptides*, (1975); Kennedy et al., *Solid Phase Biochemistry, Analytical and Synthetic Aspects*, Scouten, ed., pp. 253-391 (1983); *see, generally*, Affinity Techniques. Enzyme Purification: *Part B. Methods in Enzymology*, Vol. 34, ed. W. B. Jakoby, M. Wilchek, Acad. Press, N.Y. (1974); and Immobilized Biochemicals and Affinity Chromatography, *Advances in Experimental Medicine and Biology*, vol. 42, ed. R. Dunlap, Plenum Press, N.Y. (1974)).

- Among the most commonly used methods are absorption and adsorption or covalent binding to the support, either directly or via a linker, such as the
- 15 numerous disulfide linkages, thioether bonds, hindered disulfide bonds, and covalent bonds between free reactive groups, such as amine and thiol groups, known to those of skill in art (*see, e.g.,* the PIERCE CATALOG, ImmunoTechnology Catalog & Handbook, 1992-1993, which describes the preparation of and use of such reagents and provides a commercial source for
- 20 such reagents; Wong, *Chemistry of Protein Conjugation and Cross Linking*, CRC Press (1993); *see also* DeWitt et al., *Proc. Natl. Acad. Sci. U.S.A.*, 90:6909 (1993); Zuckermann et al., *J. Am. Chem. Soc.*, 114:10646 (1992); Kurth et al., *J. Am. Chem. Soc.*, 116:2661 (1994); Ellman et al., *Proc. Natl. Acad. Sci. U.S.A.*, 91:4708 (1994); Sucholeiki, *Tetrahedron Lttrs.*, 35:7307 (1994); Su-
- 25 Sun Wang, *J. Org. Chem.*, 41:3258 (1976); Padwa et al., *J. Org. Chem.*, 41:3550 (1971); and Vedejs et al., *J. Org. Chem.*, 49:575 (1984), which describe photosensitive linkers).

- To effect immobilization, a composition containing the protein or other biomolecule is contacted with a support material such as alumina, carbon, an
- 30 ion-exchange resin, cellulose, glass or a ceramic. Fluorocarbon polymers have been used as supports to which biomolecules have been attached by adsorption

-114-

(see, U.S. Patent No. 3,843,443; Published International PCT Application WO/86 03840).

**J. Prognosis and diagnosis**

MTSP7 protein proteins, domains, analogs, and derivatives thereof, and  
5 encoding nucleic acids (and sequences complementary thereto), and anti-MTSP7  
protein antibodies, can be used in diagnostics. Such molecules can be used in  
assays, such as immunoassays, to detect, prognose, diagnose, or monitor  
various conditions, diseases, and disorders affecting MTSP7 protein expression,  
or monitor the treatment thereof. For purposes herein, the presence of MTSP7s  
10 in body fluids or tumor tissues are of particular interest.

In particular, such an immunoassay is carried out by a method including  
contacting a sample derived from a patient with an anti-MTSP7 protein antibody  
under conditions such that specific binding can occur, and detecting or  
measuring the amount of any specific binding by the antibody. In a specific  
15 aspect, such binding of antibody, in tissue sections, can be used to detect  
aberrant MTSP7 protein localization or aberrant (*e.g.*, increased, decreased or  
absent) levels of MTSP7 protein. In a specific embodiment, antibody to MTSP7  
protein can be used to assay in a patient tissue or serum sample for the  
presence of MTSP7 protein where an aberrant level of MTSP7 protein is an  
20 indication of a diseased condition.

The immunoassays which can be used include but are not limited to  
competitive and non-competitive assay systems using techniques such as  
western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent  
assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin  
25 reactions, gel diffusion precipitin reactions, immunodiffusion assays,  
agglutination assays, complement-fixation assays, immunoradiometric assays,  
fluorescent immunoassays and protein A immunoassays.

MTSP7 protein genes and related nucleic acid sequences and  
subsequences, including complementary sequences, can also be used in  
30 hybridization assays. MTSP7 protein nucleic acid sequences, or subsequences  
thereof containing about at least 8 nucleotides, 14 or 16 or 30 or more  
continugous nucleotides, can be used as hybridization probes. Hybridization

-115-

assays can be used to detect, prognose, diagnose, or monitor conditions, disorders, or disease states associated with aberrant changes in MTSP7 protein expression and/or activity as described herein. In particular, such a hybridization assay is carried out by a method by contacting a sample containing nucleic acid  
5 with a nucleic acid probe capable of hybridizing to MTSP7 protein encoding DNA or RNA, under conditions such that hybridization can occur, and detecting or measuring any resulting hybridization.

In a specific embodiment, a method of diagnosing a disease or disorder characterized by detecting an aberrant level of an MTSP7 protein in a subject is  
10 provided herein by measuring the level of the DNA, RNA, protein or functional activity of the MTSP7 protein in a sample derived from the subject, wherein an increase or decrease in the level of the DNA, RNA, protein or functional activity of the MTSP7 protein, relative to the level of the DNA, RNA, protein or functional activity found in an analogous sample not having the disease or  
15 disorder indicates the presence of the disease or disorder in the subject.

Kits for diagnostic use are also provided, that contain in one or more containers an anti-MTSP7 protein antibody, and, optionally, a labeled binding partner to the antibody. Alternatively, the anti-MTSP7 protein antibody can be labeled (with a detectable marker, *e.g.*, a chemiluminescent, enzymatic,  
20 fluorescent, or radioactive moiety). A kit is also provided that includes in one or more containers a nucleic acid probe capable of hybridizing to MTSP protein-encoding RNA. In a specific embodiment, a kit can comprise in one or more containers a pair of primers (*e.g.*, each in the size range of 6-30 nucleotides) that are capable of priming amplification [*e.g.*, by polymerase chain reaction (see  
25 *e.g.*, Innis et al., 1990, PCR Protocols, Academic Press, Inc., San Diego, CA), ligase chain reaction (see EP 320,308) use of Q $\beta$  replicase, cyclic probe reaction, or other methods known in the art under appropriate reaction conditions of at least a portion of an MTSP protein-encoding nucleic acid. A kit can optionally further comprise in a container a predetermined amount of a purified MTSP7  
30 protein or nucleic acid, *e.g.*, for use as a standard or control.



-116-

**K. PHARMACEUTICAL COMPOSITIONS AND MODES OF ADMINISTRATION****1. Components of the compositions**

Pharmaceutical compositions containing the identified compounds that modulate the activity of an MTSP7 protein are provided herein. Also provided  
5 are combinations of a compound that modulates the activity of an MTSP7 protein and another treatment or compound for treatment of a neoplastic disorder, such as a chemotherapeutic compound.

The MTSP7 protein modulator and the anti-tumor agent can be packaged as separate compositions for administration together or sequentially or  
10 intermittently. Alternatively, they can be provided as a single composition for administration or as two compositions for administration as a single composition. The combinations can be packaged as kits.

**a. MTSP7 protein inhibitors**

Any MTSP7 protein inhibitors, including those described herein when  
15 used alone or in combination with other compounds, that can alleviate, reduce, ameliorate, prevent, or place or maintain in a state of remission of clinical symptoms or diagnostic markers associated with neoplastic diseases, including undesired and/or uncontrolled angiogenesis, can be used in the present combinations.

20 In one embodiment, the MTSP7 protein inhibitor is an antibody or fragment thereof that specifically reacts with an MTSP7 protein or the protease domain thereof, an inhibitor of the MTSP7 protein production, an inhibitor of the epithelial MTSP7 protein membrane-localization, or any inhibitor of the expression of or, especially, the activity of an MTSP7 protein.

**25 b. Anti-angiogenic agents and anti-tumor agents**

Any anti-angiogenic agents and anti-tumor agents, including those described herein, when used alone or in combination with other compounds, that can alleviate, reduce, ameliorate, prevent, or place or maintain in a state of remission of clinical symptoms or diagnostic markers associated with undesired  
30 and/or uncontrolled angiogenesis and/or tumor growth and metastasis, particularly solid neoplasms, vascular malformations and cardiovascular disorders, chronic inflammatory diseases and aberrant wound repairs, circulatory

-117-

disorders, crest syndromes, dermatological disorders, or ocular disorders, can be used in the combinations. Also contemplated are anti-tumor agents for use in combination with an inhibitor of an MTSP7 protein.

**c. Anti-tumor agents and anti-angiogenic agents**

5           The compounds identified by the methods provided herein or provided herein can be used in combination with anti-tumor agents and/or anti-angiogenesis agents.

**2. Formulations and route of administration**

10           The compounds herein and agents can be formulated as pharmaceutical compositions, generally for single dosage administration. The concentrations of the compounds in the formulations are effective for delivery of an amount, upon administration, that is effective for the intended treatment. Typically, the compositions are formulated for single dosage administration. To formulate a composition, the weight fraction of a compound or mixture thereof is dissolved, 15           suspended, dispersed or otherwise mixed in a selected vehicle at an effective concentration such that the treated condition is relieved or ameliorated. Pharmaceutical carriers or vehicles suitable for administration of the compounds provided herein include any such carriers known to those skilled in the art to be suitable for the particular mode of administration.

20           In addition, the compounds can be formulated as the sole pharmaceutically active ingredient in the composition or can be combined with other active ingredients. Liposomal suspensions, including tissue-targeted liposomes, can also be suitable as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art. For 25           example, liposome formulations can be prepared as described in U.S. Patent No. 4,522,811.

          The active compound is included in the pharmaceutically acceptable carrier in an amount sufficient to exert a therapeutically useful effect in the absence of undesirable side effects on the patient treated. The therapeutically 30           effective concentration can be determined empirically by testing the compounds in known in vitro and in vivo systems, such as the assays provided herein.

-118-

The concentration of active compound in the drug composition will depend on absorption, inactivation and excretion rates of the active compound, the physicochemical characteristics of the compound, the dosage schedule, and amount administered as well as other factors known to those of skill in the art.

5       Typically a therapeutically effective dosage is contemplated. The amounts administered can be on the order of 0.001 to 1 mg/ml, including 0.005-0.05 mg/ml or about 0.01 mg/ml, of blood volume. Pharmaceutical dosage unit forms are prepared to provide from about 1 mg to about 1000 mg, including 10 to about 500 mg, and generally about 25-75 mg of the essential  
10   active ingredient or a combination of essential ingredients per dosage unit form. The precise dosage can be empirically determined.

The active ingredient can be administered at once, or can be divided into a number of smaller doses to be administered at intervals of time. It is understood that the precise dosage and duration of treatment is a function of the  
15   disease being treated and can be determined empirically using known testing protocols or by extrapolation from in vivo or in vitro test data. It is to be noted that concentrations and dosage values can also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the  
20   individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that the concentration ranges set forth herein are exemplary only and are not intended to limit the scope or use of the claimed compositions and combinations containing them.

Exemplary pharmaceutically acceptable derivatives include acids, salts,  
25   esters, hydrates, solvates and prodrug forms. The derivative is typically selected such that its pharmacokinetic properties are superior to the corresponding neutral compound.

Thus, effective concentrations or amounts of one or more of the compounds provided herein or pharmaceutically acceptable derivatives thereof  
30   are mixed with a suitable pharmaceutical carrier or vehicle for systemic, topical or local administration to form pharmaceutical compositions. Compounds are included in an amount effective for ameliorating or treating the disorder for

-119-

which treatment is contemplated. The concentration of active compound in the composition will depend on absorption, inactivation, excretion rates of the active compound, the dosage schedule, amount administered, particular formulation as well as other factors known to those of skill in the art.

- 5           Solutions or suspensions used for parenteral, intradermal, subcutaneous, or topical application can include any of the following components: a sterile diluent, such as water for injection, saline solution, fixed oil, polyethylene glycol, glycerine, propylene glycol or other synthetic solvent; antimicrobial agents, such as benzyl alcohol and methyl parabens; antioxidants, such as ascorbic acid and
- 10       sodium bisulfite; chelating agents, such as ethylenediaminetetraacetic acid (EDTA); buffers, such as acetates, citrates and phosphates; and agents for the adjustment of tonicity such as sodium chloride or dextrose. Parenteral preparations can be enclosed in ampules, disposable syringes or single or multiple dose vials made of glass, plastic or other suitable material.
- 15           In instances in which the compounds exhibit insufficient solubility, methods for solubilizing compounds can be used. Such methods are known to those of skill in this art, and include, but are not limited to, using cosolvents, such as dimethylsulfoxide (DMSO), using surfactants, such as Tween®, or dissolution in aqueous sodium bicarbonate. Derivatives of the compounds, such
- 20       as prodrugs of the compounds can also be used in formulating effective pharmaceutical compositions. For ophthalmic indications, the compositions are formulated in an ophthalmically acceptable carrier. For the ophthalmic uses herein local administration typically is effected either by topical administration or by injection. Time release formulations are also desirable. Typically, the
- 25       compositions are formulated for single dosage administration, so that a single dose administers an effective amount.

- Upon mixing or addition of the compound with the vehicle, the resulting mixture can be a solution, suspension, emulsion or other composition. The form of the resulting mixture depends upon a number of factors, including the
- 30       intended mode of administration and the solubility of the compound in the selected carrier or vehicle. If necessary, pharmaceutically acceptable salts or other derivatives of the compounds are prepared.

-120-

The compound is included in the pharmaceutically acceptable carrier in an amount sufficient to exert a therapeutically useful effect in the absence of undesirable side effects on the patient treated. It is understood that number and degree of side effects depends upon the condition for which the compounds are administered. For example, certain toxic and undesirable side effects are tolerated when treating life-threatening illnesses that would not be tolerated when treating disorders of lesser consequence.

The compounds can also be mixed with other active materials, that do not impair the desired action, or with materials that supplement the desired action known to those of skill in the art. The formulations of the compounds and agents for use herein include those suitable for oral, rectal, topical, inhalational, buccal (*e.g.*, sublingual), parenteral (*e.g.*, subcutaneous, intramuscular, intradermal, or intravenous), transdermal administration or any route. The most suitable route in any given case will depend on the nature and severity of the condition being treated and on the nature of the particular active compound which is being used. The formulations are provided for administration to humans and animals in unit dosage forms, such as tablets, capsules, pills, powders, granules, sterile parenteral solutions or suspensions, and oral solutions or suspensions, and oil-water emulsions containing suitable quantities of the compounds or pharmaceutically acceptable derivatives thereof. The pharmaceutically therapeutically active compounds and derivatives thereof are typically formulated and administered in unit-dosage forms or multiple-dosage forms. Unit-dose forms as used herein refers to physically discrete units suitable for human and animal subjects and packaged individually as is known in the art. Each unit-dose contains a predetermined quantity of the therapeutically active compound sufficient to produce the desired therapeutic effect, in association with the required pharmaceutical carrier, vehicle or diluent. Examples of unit-dose forms include ampoules and syringes and individually packaged tablets or capsules. Unit-dose forms can be administered in fractions or multiples thereof. A multiple-dose form is a plurality of identical unit-dosage forms packaged in a single container to be administered in segregated unit-dose form. Examples of multiple-dose forms include vials, bottles of tablets or capsules or

-121-

bottles of pints or gallons. Hence, multiple dose form is a multiple of unit-doses which are not segregated in packaging.

The composition can contain along with the active ingredient: a diluent such as lactose, sucrose, dicalcium phosphate, or carboxymethylcellulose; a  
5 lubricant, such as magnesium stearate, calcium stearate and talc; and a binder such as starch, natural gums, such as gum acaciagelatin, glucose, molasses, polvinylpyrrolidone, celluloses and derivatives thereof, povidone, crospovidones and other such binders known to those of skill in the art. Liquid  
10 pharmaceutically administrable compositions can, for example, be prepared by dissolving, dispersing, or otherwise mixing an active compound as defined above and optional pharmaceutical adjuvants in a carrier, such as, for example, water, saline, aqueous dextrose, glycerol, glycols, ethanol, and the like, to thereby form a solution or suspension. If desired, the pharmaceutical composition to be  
15 administered can also contain minor amounts of nontoxic auxiliary substances such as wetting agents, emulsifying agents, or solubilizing agents, pH buffering agents and the like, for example, acetate, sodium citrate, cyclodextrine derivatives, sorbitan monolaurate, triethanolamine sodium acetate, triethanolamine oleate, and other such agents. Methods of preparing such  
20 dosage forms are known, or will be apparent, to those skilled in this art (see, *e.g.*, Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., 15th Edition, 1975). The composition or formulation to be administered will contain a quantity of the active compound in an amount sufficient to alleviate the symptoms of the treated subject.

Dosage forms or compositions containing active ingredient in the range of  
25 0.005% to 100% with the balance made up from non-toxic carrier can be prepared. For oral administration, the pharmaceutical compositions can take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (*e.g.*, pregelatinized maize starch, polyvinyl pyrrolidone or hydroxypropyl  
30 methylcellulose); fillers (*e.g.*, lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (*e.g.*, magnesium stearate, talc or silica); disintegrants (*e.g.*, potato starch or sodium starch glycolate); or wetting agents

-122-

(*e.g.*, sodium lauryl sulphate). The tablets can be coated by methods well-known in the art.

The pharmaceutical preparation can also be in liquid form, for example, solutions, syrups or suspensions, or can be presented as a drug product for reconstitution with water or other suitable vehicle before use. Such liquid preparations can be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (*e.g.*, sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (*e.g.*, lecithin or acacia); non-aqueous vehicles (*e.g.*, almond oil, oily esters, or fractionated vegetable oils); and preservatives (*e.g.*, methyl or propyl-p-hydroxybenzoates or sorbic acid).

Formulations suitable for rectal administration are generally presented as unit dose suppositories. These can be prepared by admixing the active compound with one or more conventional solid carriers, for example, cocoa butter, and then shaping the resulting mixture.

Formulations suitable for topical application to the skin or to the eye generally take the form of an ointment, cream, lotion, paste, gel, spray, aerosol and oil. Carriers which can be used include vaseline, lanoline, polyethylene glycols, alcohols, and combinations of two or more thereof. The topical formulations can further advantageously contain 0.05 to 15 percent by weight of thickeners selected from among hydroxypropyl methyl cellulose, methyl cellulose, polyvinylpyrrolidone, polyvinyl alcohol, poly (alkylene glycols), poly/hydroxyalkyl, (meth)acrylates or poly(meth)acrylamides. A topical formulation is often applied by instillation or as an ointment into the conjunctival sac. It can also be used for irrigation or lubrication of the eye, facial sinuses, and external auditory meatus. It can also be injected into the anterior eye chamber and other places. The topical formulations in the liquid state can be also present in a hydrophilic three-dimensional polymer matrix in the form of a strip, contact lens, and the like from which the active components are released.

For administration by inhalation, the compounds for use herein can be delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane,

-123-

trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, *e.g.*, gelatin, for use in an inhaler or insufflator can be formulated  
5 containing a powder mix of the compound and a suitable powder base such as lactose or starch.

Formulations suitable for buccal (sublingual) administration include, for example, lozenges containing the active compound in a flavored base, usually sucrose and acacia or tragacanth; and pastilles containing the compound in an  
10 inert base such as gelatin and glycerin or sucrose and acacia.

The compounds can be formulated for parenteral administration by injection, *e.g.*, by bolus injection or continuous infusion. Formulations for injection can be presented in unit dosage form, *e.g.*, in ampules or in multi-dose containers, with an added preservative. The compositions can be suspensions,  
15 solutions or emulsions in oily or aqueous vehicles, and can contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient can be in powder form for reconstitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water or other solvents, before use.

Formulations suitable for transdermal administration can be presented as  
20 discrete patches adapted to remain in intimate contact with the epidermis of the recipient for a prolonged period of time. Such patches suitably contain the active compound as an optionally buffered aqueous solution of, for example, 0.1 to 0.2 M concentration with respect to the active compound. Formulations suitable for transdermal administration can also be delivered by iontophoresis  
25 (*see, e.g.*, Pharmaceutical Research 3 (6), 318 (1986)) and typically take the form of an optionally buffered aqueous solution of the active compound.

The pharmaceutical compositions can also be administered by controlled release means and/or delivery devices (*see, e.g.*, in U.S. Patent Nos. 3,536,809; 3,598,123; 3,630,200; 3,845,770; 3,847,770; 3,916,899; 4,008,719;  
30 4,687,610; 4,769,027; 5,059,595; 5,073,543; 5,120,548; 5,354,566; 5,591,767; 5,639,476; 5,674,533 and 5,733,566).



-124-

Desirable blood levels can be maintained by a continuous infusion of the active agent as ascertained by plasma levels. It should be noted that the attending physician would know how to and when to terminate, interrupt or adjust therapy to lower dosage due to toxicity, or bone marrow, liver or kidney dysfunctions. Conversely, the attending physician would also know how to and when to adjust treatment to higher levels if the clinical response is not adequate (precluding toxic side effects).

The efficacy and/or toxicity of the MTSP7 protein inhibitor(s), alone or in combination with other agents can also be assessed by the methods known in the art (See generally, O'Reilly, *Investigational New Drugs*, 15:5-13 (1997)).

The active compounds or pharmaceutically acceptable derivatives can be prepared with carriers that protect the compound against rapid elimination from the body, such as time release formulations or coatings.

Kits containing the compositions and/or the combinations with instructions for administration thereof are provided. The kit can further include a needle or syringe, generally packaged in sterile form, for injecting the complex, and/or a packaged alcohol pad. Instructions are optionally included for administration of the active agent by a clinician or by the patient.

Finally, the compounds or MTSP7 proteins or protease domains thereof or compositions containing any of the preceding agents can be packaged as articles of manufacture containing packaging material, a compound or suitable derivative thereof provided herein, which is effective for treatment of a diseases or disorders contemplated herein, within the packaging material, and a label that indicates that the compound or a suitable derivative thereof is for treating the diseases or disorders contemplated herein. The label can optionally include the disorders for which the therapy is warranted.

#### **L. METHODS OF TREATMENT**

The compounds identified by the methods herein are used for treating or preventing neoplastic diseases in an animal, particularly a mammal, including a human, is provided herein. In one embodiment, the method includes administering to a mammal an effective amount of an inhibitor of an MTSP7 protein, whereby the disease or disorder is treated or prevented. In certain

-125-

embodiments, the MTSP7 protein inhibitor used in the treatment or prevention is administered with a pharmaceutically acceptable carrier or excipient. The mammal treated can be a human. The inhibitors provided herein are those identified by the screening assays. In addition, antibodies and antisense nucleic acids or or dsRNA, such as RNAi, are contemplated.

The treatment or prevention method can further include administering an anti-angiogenic treatment or agent or anti-tumor agent simultaneously with, prior to or subsequent to the MTSP7 protein inhibitor, which can be any compound identified that inhibits the activity of an MTSP7 protein, and includes an antibody or a fragment or derivative thereof containing the binding region thereof against the MTSP7 protein, an antisense nucleic acid encoding the MTSP7 protein or dsRNA, such as RNAi,, and a nucleic acid containing at least a portion of a gene encoding the MTSP7 protein into which a heterologous nucleotide sequence has been inserted such that the heterologous sequence inactivates the biological activity of at least a portion of the gene encoding the MTSP7 protein, in which the portion of the gene encoding the MTSP7 protein flanks the heterologous sequence so as to promote homologous recombination with a genomic gene encoding the MTSP7 protein.

#### 1. Antisense treatment

In a specific embodiment, as described hereinabove, MTSP7 protein function is reduced or inhibited by MTSP7 protein antisense nucleic acids, to treat or prevent neoplastic disease. The therapeutic or prophylactic use of nucleic acids of at least six nucleotides that are antisense to a gene or cDNA encoding MTSP7 protein or a portion thereof. An MTSP7 protein "antisense" nucleic acid as used herein refers to a nucleic acid capable of hybridizing to a portion of an MTSP7 protein RNA (generally mRNA) by virtue of some sequence complementarity, and usually under high stringency conditions. The antisense nucleic acid can be complementary to a coding and/or noncoding region of an MTSP7 protein mRNA. Such antisense nucleic acids have utility as therapeutics that reduce or inhibit MTSP7 protein function, and can be used in the treatment or prevention of disorders as described.

-126-

The MTSP7 protein antisense nucleic acids are of at least six nucleotides and are typically oligonucleotides (ranging from 6 to about 150 nucleotides, generally 6 to 50 nucleotides). In specific aspects, the oligonucleotide is at least 10 nucleotides, at least 15 nucleotides, at least 100 nucleotides, or at least 125  
 5 nucleotides depending upon the purpose and conditions of use, including desired specificity. The oligonucleotides can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone. The oligonucleotide can include other appending groups such as  
 10 peptides, or agents facilitating transport across the cell membrane (see, *e.g.*, Letsinger et al., *Proc. Natl. Acad. Sci. U.S.A.* 86:6553-6556 (1989); Lemaitre et al., *Proc. Natl. Acad. Sci. U.S.A.* 84:648-652 (1987); PCT Publication No. WO 88/09810, published December 15, 1988) or blood-brain barrier (see, *e.g.*, PCT Publication No. WO 89/10134, published April 25, 1988), hybridization-  
 15 triggered cleavage agents (see, *e.g.*, Krol et al., *BioTechniques* 6:958-976 (1988)) or intercalating agents (see, *e.g.*, Zon, *Pharm. Res.* 5:539-549 (1988)).

The MTSP7 protein antisense nucleic acid is generally an oligonucleotide, typically single-stranded DNA. In an embodiment, the oligonucleotide includes a sequence antisense to a portion of human MTSP7 protein. The oligonucleotide  
 20 can be modified at any position on its structure with substituents generally known in the art.

The MTSP7 protein antisense oligonucleotide can include at least one modified base moiety which is selected from the group including, but not limited to 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine,  
 25 xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine,  
 30 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v),

-127-

wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

- 5 In another embodiment, the oligonucleotide includes at least one modified sugar moiety selected from the group including but not limited to arabinose, 2-fluoroarabinose, xylulose, and hexose. The oligonucleotide can include at least one modified phosphate backbone selected from a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a  
 10 phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

- The oligonucleotide can be an  $\alpha$ -anomeric oligonucleotide. An  $\alpha$ -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which the strands run parallel to each other (Gautier et al., *Nucl. Acids Res.*  
 15 15:6625-6641 (1987)).

The oligonucleotide can be conjugated to another molecule, *e.g.*, a peptide, hybridization triggered cross-linking agent, transport agent and hybridization-triggered cleavage agent.

- The oligonucleotides can be synthesized by standard methods known in  
 20 the art, *e.g.* by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems). As examples, phosphorothioate oligonucleotides can be synthesized by the method of Stein et al. (*Nucl. Acids Res.* 16:3209 (1988)), methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., *Proc. Natl. Acad. Sci.*  
 25 *U.S.A.* 85:7448-7451 (1988)).

- In a specific embodiment, the MTSP7 protein antisense oligonucleotide includes catalytic RNA, or a ribozyme (see, *e.g.*, PCT International Publication WO 90/11364, published October 4, 1990; Sarver et al., *Science* 247:1222-1225 (1990)). In another embodiment, the oligonucleotide is a 2'-O-  
 30 methylribonucleotide (Inoue et al., *Nucl. Acids Res.* 15:6131-6148 (1987)), or a chimeric RNA-DNA analogue (Inoue et al., *FEBS Lett.* 215:327-330 (1987)).

-128-

In an alternative embodiment, the MTSP7 protein antisense nucleic acid is produced intracellularly by transcription from an exogenous sequence. For example, a vector can be introduced *in vivo* such that it is taken up by a cell, within which cell the vector or a portion thereof is transcribed, producing an antisense nucleic acid (RNA). Such a vector would contain a sequence encoding the MTSP7 protein antisense nucleic acid. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be constructed by recombinant DNA technology methods standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in mammalian cells. Expression of the sequence encoding the MTSP7 protein antisense RNA can be by any promoter known in the art to act in mammalian, including human, cells. Such promoters can be inducible or constitutive. Such promoters include but are not limited to: the SV40 early promoter region (Bernoist and Chambon, *Nature* 290:304-310 (1981)), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., *Cell* 22:787-797 (1980)), the herpes thymidine kinase promoter (Wagner et al., *Proc. Natl. Acad. Sci. U.S.A.* 78:1441-1445 (1981)), the regulatory sequences of the metallothionein gene (Brinster et al., *Nature* 296:39-42 (1982)).

The antisense nucleic acids include sequence complementary to at least a portion of an RNA transcript of an MTSP7 protein gene, including a human MTSP7 protein gene. Absolute complementarity, although desirable, is not required.

The amount of MTSP7 protein antisense nucleic acid that will be effective in the treatment or prevention of neoplastic disease will depend on the nature of the disease, and can be determined empirically by standard clinical techniques. Where possible, it is desirable to determine the antisense cytotoxicity in cells *in vitro*, and then in useful animal model systems prior to testing and use in humans.

-129-

## 2. RNA interference

For example, RNA interference (RNAi) (see, *e.g.* Chuang *et al.* (2000) *Proc. Natl. Acad. Sci. U.S.A.* 97:4985) can be employed to inhibit the expression of a gene encoding an MTSP7. Interfering RNA (RNAi) fragments, particularly double-stranded (ds) RNAi, can be used to generate loss-of- MTSP7 function. Methods relating to the use of RNAi to silence genes in organisms including, mammals, *C. elegans*, *Drosophila* and plants, and humans are known (see, *e.g.*, Fire *et al.* (1998) *Nature* 391:806-811 Fire (1999) *Trends Genet.* 15:358-363; Sharp (2001) *Genes Dev.* 15:485-490; Hammond, *et al.* (2001) *Nature Rev. Genet.* 2:110-1119; Tuschl (2001) *Chem. Biochem.* 2:239-245; Hamilton *et al.* (1999) *Science* 286:950-952; Hammond *et al.* (2000) *Nature* 404:293-296; Zamore *et al.* (2000) *Cell* 101:25-33; Bernstein *et al.* (2001) *Nature* 409: 363-366; Elbashir *et al.* (2001) *Genes Dev.* 15:188-200; Elbashir *et al.* (2001) *Nature* 411:494-498; International PCT application No. WO 01/29058; International PCT application No. WO 99/32619). Double-stranded RNA (dsRNA)-expressing constructs are introduced into a host, such as an animal or plant using, a replicable vector that remains episomal or integrates into the genome. By selecting appropriate sequences, expression of dsRNA can interfere with accumulation of endogenous mRNA encoding an MTSP7. RNAi can also be used to inhibit expression *in vitro*. Regions include at least about 21 (or 21) nucleotides that are selective (*i.e.* unique) for MTSP7 are used to prepare the RNAi. Smaller fragments of about 21 nucleotides can be transformed directly into cells; larger RNAi dsRNA molecules are generally introduced using vectors that encode them. dsRNA molecules are at least about 21 bp long or longer, such as 50, 100, 150, 200 and longer.

## 3. Gene Therapy

In an exemplary embodiment, nucleic acids that include a sequence of nucleotides encoding an MTSP7 protein or functional domains or derivative thereof, are administered to promote MTSP7 protein function, by way of gene therapy. Gene therapy refers to therapy performed by the administration of a nucleic acid to a subject. In this embodiment, the nucleic acid produces its encoded protein that mediates a therapeutic effect by promoting MTSP7 protein

-130-

function. Any of the methods for gene therapy available in the art can be used (see, Goldspiel et al., *Clinical Pharmacy* 12:488-505 (1993); Wu and Wu, *Biotherapy* 3:87-95 (1991); Tolstoshev, *An. Rev. Pharmacol. Toxicol.* 32:573-596 (1993); Mulligan, *Science* 260:926-932 (1993); and Morgan and Anderson, *An. Rev. Biochem.* 62:191-217 (1993); *TIBTECH* 11(5):155-215 (1993). For example, one therapeutic composition for gene therapy includes an MTSP7 protein-encoding nucleic acid that is part of an expression vector that expresses an MTSP7 protein or domain, fragment or chimeric protein thereof in a suitable host. In particular, such a nucleic acid has a promoter operably linked to the MTSP7 protein coding region, the promoter being inducible or constitutive, and, optionally, tissue-specific. In another particular embodiment, a nucleic acid molecule is used in which the MTSP7 protein coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the MTSP protein nucleic acid (Koller and Smithies, *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); Zijlstra et al., *Nature* 342:435-438 (1989)).

Delivery of the nucleic acid into a patient can be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid-carrying vector, or indirect, in which case, cells are first transformed with the nucleic acid *in vitro*, then transplanted into the patient. These two approaches are known, respectively, as *in vivo* or *ex vivo* gene therapy.

In a specific embodiment, the nucleic acid is directly administered *in vivo*, where it is expressed to produce the encoded product. This can be accomplished by any of numerous methods known in the art, *e.g.*, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, *e.g.*, by infection using a defective or attenuated retroviral or other viral vector (see U.S. Patent No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle bombardment (*e.g.*, a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering it in linkage to a peptide

-131-

which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see e.g., Wu and Wu, *J. Biol. Chem.* 262:4429-4432 (1987)) (which can be used to target cell types specifically expressing the receptors). In another embodiment, a nucleic acid-ligand complex  
5 can be formed in which the ligand is a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted *in vivo* for cell specific uptake and expression, by targeting a specific receptor (see, e.g., PCT Publications WO 92/06180 dated April 16, 1992 (Wu et al.); WO 92/22635  
10 dated December 23, 1992 (Wilson et al.); WO92/20316 dated November 26, 1992 (Findeis et al.); WO93/14188 dated July 22, 1993 (Clarke et al.), WO 93/20221 dated October 14, 1993 (Young)). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination (Koller and Smithies, *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); Zijlstra et al., *Nature* 342:435-438 (1989)).  
15

In a specific embodiment, a viral vector that contains the MTSP7 protein nucleic acid is used. For example, a retroviral vector can be used (see Miller et al., *Meth. Enzymol.* 217:581-599 (1993)). These retroviral vectors have been modified to delete retroviral sequences that are not necessary for packaging of  
20 the viral genome and integration into host cell DNA. The MTSP7 protein nucleic acid to be used in gene therapy is cloned into the vector, which facilitates delivery of the gene into a patient. More detail about retroviral vectors can be found in Boesen et al., *Biotherapy* 6:291-302 (1994), which describes the use of a retroviral vector to deliver the *mdr1* gene to hematopoietic stem cells in order  
25 to make the stem cells more resistant to chemotherapy. Other references illustrating the use of retroviral vectors in gene therapy are: Clowes et al., *J. Clin. Invest.* 93:644-651 (1994); Kiem et al., *Blood* 83:1467-1473 (1994); Salmons and Gunzberg, *Human Gene Therapy* 4:129-141 (1993); and Grossman and Wilson, *Curr. Opin. in Genetics and Devel.* 3:110-114 (1993).

30 Adenoviruses are other viral vectors that can be used in gene therapy. Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a



-132-

mild disease. Other targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Kozarsky and Wilson, *Current Opinion in Genetics and Development* 3:499-503 (1993) present a  
5 review of adenovirus-based gene therapy. Bout et al., *Human Gene Therapy* 5:3-10 (1994) demonstrated the use of adenovirus vectors to transfer genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of adenoviruses in gene therapy can be found in Rosenfeld et al., *Science* 252:431-434 (1991); Rosenfeld et al., *Cell* 68:143-155 (1992); and Mastrangeli et al., *J. Clin. Invest.* 91:225-234 (1993). Adeno-associated virus (AAV) has been used  
10 in gene therapy (Walsh et al., *Proc. Soc. Exp. Biol. Med.* 204:289-300 (1993).

Another approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of  
15 transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a patient.

In this embodiment, the nucleic acid is introduced into a cell prior to administration *in vivo* of the resulting recombinant cell. Such introduction can  
20 be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcell-mediated gene transfer and spheroplast fusion. Numerous techniques are known in the art for the  
25 introduction of foreign genes into cells (see *e.g.*, Loeffler and Behr, *Meth. Enzymol.* 217:599-618 (1993); Cohen et al., *Meth. Enzymol.* 217:618-644 (1993); Cline, *Pharmac. Ther.* 29:69-92 (1985)) and can be used, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the  
30 nucleic acid to the cell, so that the nucleic acid is expressible by the cell and also can be heritable and expressible by its cell progeny.

-133-

The resulting recombinant cells can be delivered to a patient by various methods known in the art. In certain embodiments, epithelial cells are injected, *e.g.*, subcutaneously. In another embodiment, recombinant skin cells can be applied as a skin graft onto the patient. Recombinant blood cells (*e.g.*,

5 hematopoietic stem or progenitor cells) are generally administered intravenously. The amount of cells envisioned for use depends on the desired effect, patient state and other parameter, and can be determined by one skilled in the art.

Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not  
10 limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as T lymphocytes, B lymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, *e.g.*, as obtained from bone marrow, hepatocytes, umbilical  
15 cord blood, peripheral blood and fetal liver and other sources of such cells.

In certain embodiments, the cell used for gene therapy is autologous to the patient. In an embodiment in which recombinant cells are used in gene therapy, an MTSP7 protein nucleic acid is introduced into the cells such that it is expressible by the cells or their progeny, and the recombinant cells are then  
20 administered *in vivo* for therapeutic effect. In a specific embodiment, stem or progenitor cells are used. Any stem and/or progenitor cells which can be isolated and maintained *in vitro* can potentially be used in accordance with this embodiment. Such stem cells include but are not limited to hematopoietic stem cells (HSC), stem cells of epithelial tissues such as the skin and the lining of the  
25 gut, embryonic heart muscle cells, liver stem cells (PCT Publication WO 94/08598, dated April 28, 1994), and neural stem cells (Stemple and Anderson, *Cell* 71:973-985 (1992)).

Epithelial stem cells (ESCs) or keratinocytes can be obtained from tissues such as the skin and the lining of the gut by known procedures (Rheinwald,  
30 *Meth. Cell Bio.* 21A:229 (1980)). In stratified epithelial tissue such as the skin, renewal occurs by mitosis of stem cells within the germinal layer, the layer closest to the basal lamina. Stem cells within the lining of the gut provide for a

-134-

rapid renewal rate of this tissue. ESCs or keratinocytes obtained from the skin or lining of the gut of a patient or donor can be grown in tissue culture (Rheinwald, *Meth. Cell Bio.* 21A:229 (1980); Pittelkow and Scott, *Cano Clinic Proc.* 61:771 (1986)). If the ESCs are provided by a donor, a method for  
5 suppression of host versus graft reactivity (e.g., irradiation, drug or antibody administration to promote moderate immunosuppression) can also be used.

With respect to hematopoietic stem cells (HSC), any technique which provides for the isolation, propagation, and maintenance *in vitro* of HSC can be used in this embodiment. Techniques by which this can be accomplished  
10 include (a) the isolation and establishment of HSC cultures from bone marrow cells isolated from the future host, or a donor, or (b) the use of previously established long-term HSC cultures, which can be allogeneic or xenogeneic. Non-autologous HSC are used generally in conjunction with a method of suppressing transplantation immune reactions of the future host/patient. In a  
15 particular embodiment, human bone marrow cells can be obtained from the posterior iliac crest by needle aspiration (see, e.g., Kodo et al., *J. Clin. Invest.* 73:1377-1384 (1984)). In certain embodiments, the HSCs can be made highly enriched or in substantially pure form. This enrichment can be accomplished before, during, or after long-term culturing, and can be done by any techniques  
20 known in the art. Long-term cultures of bone marrow cells can be established and maintained by using, for example, modified Dexter cell culture techniques (Dexter et al., *J. Cell Physiol.* 91:335 (1977) or Witlock-Witte culture techniques (Witlock and Witte, *Proc. Natl. Acad. Sci. USA* 79:3608-3612 (1982)).

In a specific embodiment, the nucleic acid to be introduced for purposes  
25 of gene therapy includes an inducible promoter operably linked to the coding region, such that expression of the nucleic acid is controllable by controlling the presence or absence of the appropriate inducer of transcription.

### 3. Prodrugs

A method for treating tumors is provided. The method is practiced by  
30 administering a prodrug that is specifically cleaved by an MTSP7 to release an active drug. Upon contact with a cell that expresses MTSP7 activity, the prodrug is converted into an active drug. The prodrug can be a conjugate that

-135-

contains the active agent, such as an anti-tumor drug, such as a cytotoxic agent, or other therapeutic agent, linked, linked to a substrate for the targeted MTSP7, such that the drug or agent is inactive or unable to enter a cell, in the conjugate, but is activated upon cleavage. The prodrug, for example, can  
5 contain an oligopeptide, typically a relatively short, less than about 10 amino acids peptide, that is selectively proteolytically cleaved by the targeted MTSP7. Cytotoxic agents, include, but are not limited to, alkylating agents, antiproliferative agents and tubulin binding agents. Others include, vinca drugs, mitomycins, bleomycins and taxanes.

#### 10 M. ANIMAL MODELS

Transgenic animal models are provided herein. Such an animal can be produced by promoting recombination between an exogenous MTSP7 gene that could be over-expressed or mis-expressed, such as by expression under a strong promoter, via homologous or other recombination event. For example,  
15 transgenic animals can be produced by introducing the nucleic acid using vectors or other modes of gene delivery into a germline cell, such as an embryonic stem cell. Typically the nucleic acid is introduced, such as an embryonic stem cell, which is then injected by transforming embryo-derived stem (ES) cells with a vector containing the MTSP7 protein-encoding nucleic acid followed by injecting  
20 the ES cells into a blastocyst, and implanting the blastocyst into a foster mother, followed by the birth of a transgenic animal. Generally introduction into a chromosome of the animal occurs by a recombination between the heterologous MTSP7-encoding nucleic acid and endogenous nucleic acid. The heterologous nucleic acid can be targeted to a specific chromosome.

25 In some instances, knockout animals can be produced. Such an animal can be initially produced by promoting homologous recombination between an MTSP7 protein gene in its chromosome and an exogenous MTSP7 protein gene that has been rendered biologically inactive (typically by insertion of a heterologous sequence, *e.g.*, an antibiotic resistance gene). In one embodiment,  
30 this homologous recombination is performed by transforming embryo-derived stem (ES) cells with a vector containing the insertionally inactivated MTSP7 protein gene, such that homologous recombination occurs, followed by injecting

-136-

the ES cells into a blastocyst, and implanting the blastocyst into a foster mother, followed by the birth of the chimeric animal ("knockout animal") in which an MTSP7 protein gene has been inactivated (see Capecchi, *Science* 244:1288-1292 (1989)). The chimeric animal can be bred to produce additional

5 knockout animals. Such animals include, but are not limited to, mice, hamsters, sheep, pigs, cattle, and other non-human mammals. For example, a knockout mouse is produced. Such knockout animals are expected to develop or be predisposed to developing neoplastic diseases and thus can have use as animal models of such diseases *e.g.*, to screen for or test molecules for the ability to

10 treat or prevent such diseases or disorders. Such an animal can be initially produced by promoting homologous recombination between an MTSP7 gene in its chromosome and an exogenous MTSP7 protein gene that would be over-expressed or mis-expressed (generally by expression under a strong promoter). In an embodiment, this homologous recombination is carried out by transforming

15 embryo-derived stem (ES) cells with a vector containing the over-expressed or mis-expressed MTSP7 protein gene, such that homologous recombination occurs, followed by injecting the ES cells into a blastocyst, and implanting the blastocyst into a foster mother, followed by the birth of the chimeric animal in which an MTSP7 gene has been over-expressed or mis-expressed (see Capecchi,

20 *Science* 244:1288-1292 (1989)). The chimeric animal can be bred to produce additional animals with over-expressed or mis-expressed MTSP7 protein. Such animals include, but are not limited to, mice, hamsters, sheep, pigs, cattle and other non-human mammals. In a specific embodiment, a mouse with over-expressed or mis-expressed MTSP7 protein is produced.

25

The following examples are included for illustrative purposes only and are not intended to limit the scope of the invention.

#### EXAMPLE 1

##### Identification of MTSP7

30 The protein sequence of the protease domain of matriptase (MTSP1; accession number AF118224) was used to search the human HTGS (high throughput genomic sequence) database using the tblastn algorithm. This

-137-

search and alignment algorithm compares a protein query sequence against a nucleotide sequence database dynamically translated in all six reading frames (both strands). Several potential new serine proteases were identified, among them was one designated herein as MTSP7. The translated sequence of MTSP7  
5 has 53% identity to matriptase and 48% identity with endotheliase 1 (accession number AF064819). A search of a human genome database revealed identity in clones designated AP002824 and AC012571; and searches of human EST databases revealed several matches as well.

#### **Identification of tissue source for cloning of MTSP7**

10 Using the nucleotide sequence of MTSP7 derived from the genomic sequence, two gene specific oligonucleotide primers were designed. The sequence for the 5' end primer is 5'-AATGGCCATGGCAGGCCAGCCTCC-3' SEQ ID No. 5 and that of the 3' end is  
5'-GTCCCCAACTTACTATACCTACAATGTACCAG-3' SEQ ID No. 6. These  
15 primers were used to screen a panel of 8 cDNA libraries derived normal human tissues (Human Multiple Tissue cDNA Panel I; Clontech, Palo Alto, CA; catalog no. K1420-1). A major band was detected in human placenta, and subsequent sequence analysis showed that the nucleotide sequence of this DNA fragment  
20 matched that of the genomic MTSP7 clone. The human placenta was then chosen as the tissue source for isolation of the full length cDNA.

#### **5'- and 3'- rapid amplification of cDNA ends (RACE)**

To obtain the full-length cDNA of MTSP7, 5'- and 3'-RACE reactions were performed. RACE-ready cDNA library from human placenta was prepared using the SMART RACE cDNA amplification kit (Clontech, Palo Alto, CA; catalog  
25 no. K1811-1). Two gene specific primers were used:  
5'-GTCCCCAACTTACTATACCTACAATGTACCAG -3' SEQ ID No. 7 for 5'-RACE reaction and 5'-AATGGCCATGGCAGGCCAGCCTCC -3' SEQ ID No. 8 for  
3'-RACE reaction. A ~1.5 kbp cDNA fragment was obtained from the 5'-RACE reaction. The 3'-RACE reaction also yielded a ~1.5 kbp fragment. The  
30 fragments were subcloned into pCR2.1-TOPO TA cloning vector (Invitrogen, Carlsbad, CA; catalog no. K-4500-01). The resulting clones were analyzed by

-138-

Southern analysis using the cDNA insert originally derived from human placenta as a probe and by DNA sequence analysis.

**Domain organization of MTSP7 and homology to endotheliase 1**

Sequence analysis of the translated MTSP7 coding sequence indicated  
5 that MTSP7 is a type-II membrane-type serine protease. It has a transmembrane domain at the N-terminus, followed by a SEA (sea urchin sperm protein-enterokinase-agrin) domain. Studies suggest that the SEA domain can function in the binding of carbohydrate moieties. The C-terminus contains a  
10 trypsin-like serine protease domain characterized by the presence of a protease activation cleavage site at the beginning of the domain and the catalytic triad residues (histidine, aspartate and serine) in 3 highly-conserved regions of the catalytic domain. Alignment of the protein sequence with that of endotheliase 1 showed 42% identity with the full-length protein and 58% identity with the protease domain.

**15 Amplification of cDNA encoding full-length protease domain of MTSP7**

To obtain the cDNA fragment encoding the protease domain of MTSP7, an end-to-end PCR amplification using gene-specific primers and the SMART RACE cDNA library from human placenta was used. The two primers used were: 5'- TGCCATTACCAGCATCCTCTTCTACTCAAAG -3' for the 5' end SEQ  
20 ID No. 9 and 5'- CCATGTGTATAACTCACGGACAATCCACACTAC -3' SEQ ID No. 10 for the 3' end. The 5' primer contained the sequence that encodes a region immediately upstream of the start of the MTSP7 protease domain (MPLPASSSTQ; SEQ ID No. 11). The 3' primer corresponds to the sequence flanking the stop codon (underlined) and the sequence immediately downstream  
25 of the stop codon. A 760-bp fragment was amplified from the human placenta cDNA library. The PCR product was isolated and purified using the QIAquick gel extraction kit (Qiagen, Valencia, CA; catalog no. 28704). The MTSP7 PCR product was used to amplify the cDNA fragment containing the appropriate restriction sites for subcloning into the *Pichia* vector, pPIC9KX. The  
30 gene-specific primers used were:  
5'- TCTCTCGAGAAAAGAATTGTCCAAGGAAGGGAAACAGCTATG-3' SEQ ID No. 12 at the 5' end and

-139-

5'-ATAGCGGCCGCGCACACTACATACCAGTCTTTGAGGCAATC-3' SEQ ID No. 13

at the 3' end. The 5' end primer contained an XhoI site (underlined) immediately upstream of the Pichia protease cleavage site and part of the MTSP7 protease domain (KRIVQGRETAM; SEQ ID No. 14), while the 3' end primer contained a

5 NotI site (underlined) downstream of the stop codon (in bold).

#### Gene expression profile of MTSP7 in normal tissues and tumor cell lines

To obtain information regarding the gene expression profile of the MTSP7 transcript, the MTSP7 cDNA fragment obtained from human placenta was used to probe an RNA dot blot composed of 76 different human tissues (Human

10 Multiple Tissue Expression (MTE) Array; Clontech, Palo Alto, CA; catalog no. 7775-1). Results show that MTSP7 is ubiquitously expressed, with highest levels (in decreasing intensity signal) found in kidney, spleen, placenta, lung, liver, bone marrow, pituitary gland, spinal cord, peripheral blood leukocyte, lymph node, ovary, mammary gland, adrenal gland, thyroid gland, bladder,

15 uterus and prostate. It is also highly expressed in lung carcinoma (A549 cell line), leukemia (K-562 cell line) and cervical carcinoma (HeLaS3 cell line).

#### Sequence analysis

MTSP7 DNA and protein sequences were analyzed using MacVector (version 6.5; Oxford Molecular Ltd., Madison, WI). The full length cDNA of

20 MTSP7 is 2,100 bp long with an ORF composed of 1,317 bp which translate to a 438-amino acid protein (SEQ ID Nos. 15 and 16). The cDNA encoding the protease domain in MTSP7 is composed of 702 base pairs, which translate to a 233-amino acid protein sequence and stop codon. The following are ORF cDNA sequence and the translated protein sequence of MTSP7 (see SEQ ID No. 15

25 and 16).

MTSP7/full length cDNA sequence Range: 1 to 2100

ATGATGTACACACCTGTTGAATTTTCAGAAGCTGAAT  
TCTCACGAGCTGAATATCAAAGAAAGCAGCAATTTTG  
GGACTCAGTACGGCTAGCTCTTTTCACATTAGCAATT  
30 GTAGCAATCATAGGAATTGCAATTGGTATTGTTACTC  
ATTTTGTTGTTGAGGATGATAAGTCTTTCTATTACCT  
TGCCTCTTTTAAAGTCACAAATATCAAATATAAAGAA  
AATTATGGCATAAGATCTTCAAGAGAGTTTATAGAAA  
GGAGTCATCAGATTGAAAGAATGATGTCTAGGATATT  
35 TCGACATTCTTCTGTAGGCGGTCGATTTATCAAATCT



-140-

CATGTTATCAAATTAAGTCCAGATGAACAAGGTGTGG  
ATATTCTTATAGTGCTCATATTTTCGATACCCATCTAC  
TGATAGTGCTGAACAAATCAAGAAAAAATTGAAAAG  
GCTTTATATCAAAGTTTGAAGACCAAACAATTGTCTT  
5 TGACCATAAACAAACCATCATTTAGACTCACACCTAT  
TGACAGCAAAAAGATGAGGAATCTTCTCAACAGTCGC  
TGTGGAATAAGGATGACATCTTCAAACATGCCATTAC  
CAGCATCCTCTTCTACTCAAAGAATTGTCCAAGGAAG  
GGAAACAGCTATGGAAGGGGAATGGCCATGGCAGGCC  
10 AGCCTCCAGCTCATAGGGTCAGGCCATCAGTGTGGAG  
CCAGCCTCATCAGTAACACATGGCTGCTCACAGCAGC  
TCACTGCTTTTGGAAAAATAAAGACCCAACTCAATGG  
ATTGCTACTTTTGGTGCAACTATAACACCACCCGCAG  
TGAAACGAAATGTGAGGAAAATTATTCTTCATGAGAA  
15 TTACCATAGAGAAACAAATGAAAATGACATTGCTTTG  
GTTTCTCAGCTCTCTACTGGAGTTGAGTTTTTCAAATATAG  
TCCAGAGAGTTTGCCTCCCAGACTCATCTATAAAGTT  
GCCACCTAAAACAAGTGTGTTTCGTCACAGGATTTGGA  
TCCATTGTAGATGATGGACCTATACAAAATACACTTC  
20 GGCAAGCCAGAGTGGAAACCATAAGCACTGATGTGTG  
TAACAGAAAGGATGTGTATGATGGCCTGATAACTCCA  
GGAATGTTATGTGCTGGATTTCATGGAAGGAAAAATAG  
ATGCATGTAAGGGAGATTCTGGTGGACCTCTGGTTTA  
TGATAATCATGACATCTGGTACATTGTAGGTATAGTA  
25 AGTTGGGGACAATCATGTGCACTTCCCAAAAAACCTG  
GAGTCTACACCAGAGTAACTAAGTATCGAGATTGGAT  
TGCCTCAAAGACTGGTATGTAG

30 MTSP7-coding region-cDNA and protein sequences Range: 45 to 1361

MMYTPVEFSEAEFSRAEYQRKQQFWDSVRLALFTLAIVA  
IIGIAIGIVTHFVVEDDKSFYYLASFKVTNIKYKENYGI  
RSSREFIERSHQIERMMSRIFRHSSVGGRFIKSHVIKLS  
35 PDEQGV DILIVLIFRYPSTDSAEQIKKKIEKALYQSLKT  
KQLSLTINKPSFRLTPIDSKKMRNLLNSRCGIRMTSSNM  
PLPASSSTQRIVQGRETAMEGEWPWQASLQLIGSGHQCG  
ASLISNTWLLTAAHCFWKNKDPTQWIATFGATITPPAVK  
RNVRKIILHENYHRETNENDIALVOLSTGVEFSNIVQRV  
40 CLPDSSIKLPPKTSVFVTGFGSIVDDGPIONTLRQARVE  
TISTDVCNRKDVYDGLITPGMLCAGFMEGKIDACKGDSG  
GPLVYDNHDIWYIVGIVSWGQSCALPKKPGVYTRVTKYR  
DWIASKTGM

-141-

**EXAMPLE 2****Cloning and expression of the protease domain of MTSP7 for expression  
C122S mutagenesis of the Protease domain of MTSP7**

The gene encoding the protease domain of MTSP7 (the N-terminus  
 5 starting at the I206) was mutagenized by PCR SOE (PCR-based splicing by  
 overlap extension) to replace the unpaired cysteine at position 122  
 (chymotrypsin numbering system; cysteine 313 in MTSP7 sequence) with a  
 serine. Two overlapping gene fragments, each replacing the TGC codon for  
 cysteine with an AGC codon for serine at position 122 were PCR amplified using  
 10 the following primers: for the 5' gene fragment,  
 TCTCTCGAGAAAAGAATTGTCCAAGGAAGGGAAACAGCTATG SEQ ID NO. 19)  
 and AGATGAGTCTGGGAGGCTAACTCTCTGGACTAT (SEQ ID No. 20); for the  
 3' gene fragment, ATTCGCGGCCGCCTACATACCAGTCTTTGAGGCAAT (SEQ  
 ID No. 21) and ATAGTCCAGAGAGTTAGCCTCCCAGACTCATCT (SEQ ID No.  
 15 22).

The amplified gene fragments were purified on a 1% agarose gel, mixed  
 and reamplified by PCR to produce the full length coding sequence for the  
 protease domain of MTSP7 C122S. This sequence was then cut with restriction  
 enzymes NotI and XhoI, and ligated into the XhoI/NotI sites of Pichia vector  
 20 pPic9KX.

**Expression of MTSP-7 by fermentation of Pichia**

Nucleic acid encoding each the MTSP7 protease domain thereof was  
 cloned (with a C313S mutation) into a derivative, modified by removal of a  
 restriction site as described below, of the *Pichia pastoris* vector pPIC9K  
 25 (available from Invitrogen; see SEQ ID NO. 45). Plasmid pPIC9k features include  
 the 5' AOX1 promoter fragment at 1-948; 5' AOX1 primer site at 855-875;  
 alpha-factor secretion signal(s) at 949-1218; alpha-factor primer site at  
 1152-1172; multiple cloning site at 1192-1241; 3' AOX1 primer site at  
 1327-1347; 3' AOX1 transcription termination region at 1253-1586; HIS4 ORF  
 30 at 4514-1980; kanamycin resistance gene at 5743-4928; 3' AOX1 fragment at  
 6122-6879; ColE1 origin at 7961-7288; and the ampicillin resistance gene at  
 8966-8106. The plasmid used herein is derived from pPIC9K by eliminating the

-142-

XhoI site in the kanamycin resistance gene and the resulting vector is herein designated pPIC9KX.

#### Fermentation

- P. pastoris* clone GS115/pPIC9K:MTSP7 C122S Sac SC2 that expresses
- 5 the C122S mutant form of MTSP7 was fermented at the 5 liter scale. An overnight culture of 200 ml (OD600 of approximately 12) was used to inoculate 3.2 liters of fermentation medium in each of two Bioflo vessels (New Brunswick Scientific, Edison, NJ). The batch phase complex medium contained 10 g/l yeast extract, 20 g/l peptone, 40 g/l glycerol, 5 g/l ammonium sulfate, 0.2 g/l
- 10 calcium sulfate(dihydrate), 2 g/l magnesium sulfate(heptahydrate), 2 g/l potassium sulfate, 25 g/l sodium hexametaphosphate, and 4.35 ml/l PTM1. The culture was grown at a pH of 6.0 and a temperature of 28 °C. Concentrated ammonium hydroxide was used to maintain the pH of the culture. KFO 880 (KABO Chemicals, Cheyenne,WY) was used as needed to control foaming.
- 15 The batch phase of the fermentation lasted about 26 hours at which time the culture had consumed all of the initial glycerol in the media. A substrate limited fed-batch of 50% (w/v) glycerol was initiated at 18 ml/l x hr at this point and continued for about 4 hours. The culture reached a density of about 200 g/l wet cell weight by this point.
- 20 Methanol induction was initiated following the end of the glycerol fed-batch phase. The culture was transitioned to methanol utilization (see, Zhang *et al.* (2000) Modeling *Pichia pastoris* Growth on Methanol and Optimizing the Production of a Recombinant Protein, the Heavy-Chain Fragment C of Botulinum Neurotoxin, Serotype A. Biotechnology and Bioengineering Vol. 70, No 1
- 25 October 5, 2000) by adding 1.5 ml of methanol per liter of culture and linearly decreasing the glycerol feed rate from 18 ml/l x hr to 0 ml/l x hr over a 3 hour period. The methanol addition served as an on-line calibration of the MeOH Sensor (Raven Biotech, Vancouver, BC, Canada) that was used to control the fermenter throughout induction. After the initial amount of methanol was
- 30 utilized, as indicated by the MeOH Sensor, another 1.5 ml/l was added to the culture and the MeOH sensor was used to control the methanol concentration in the fermenter at that level (i.e., 0.15%) throughout the induction phase. The

-143-

methanol fed to the fermenter was supplemented with 2 ml/l PTM4 solution.

The induction phase lasted about 42 hours.

**Initial Work-up of MTSP-7 containing conditioned media:**

The supernatant from each of the fermentations was harvested by  
5 centrifugation, pooled, and then was concentrated to about 0.5 liter using a 10  
kDa ultrafiltration cartridge (A/G Technologies Corp., Needham, MA) on a SRT5  
ultrafiltration system (North Carolina SRT, Cary, NC). The concentrate was  
drained from the system, and the system was rinsed with a volume of buffer (50  
mM Tris, pH=8.0, 50 mM NaCl, 0.005% Tween 80) equal to the initial volume  
10 of the concentrated material. The concentrate and the rinse material were  
combined to yield the final ultrafiltration product of about 1 liter. A final  
clarification of the supernatant was done with a SartoBran 300 0.45 + 0.2  $\mu$ m  
capsule filter (Sartorius Separations Div., Edgewood, NJ).

**Purification of MTSP7:**

15 After a 5L fermentation, cells were separated from the medium by  
centrifugation. The supernatant was brought to 40% saturation with ammonium  
sulfate and centrifuged. The pellet was dissolved in 250 mL 20 mM Hepes, 150  
mM NaCl pH 7.5 (HS buffer). The resulting protein solution was stored at  
-20°C in 50 mL aliquots. After thawing, 50 mL of protein solution was dialyzed  
20 against 4 L of HS buffer. To the dialysate, 4 mL of 10 % polyethylenimine was  
added, and the precipitate was removed by centrifugation at 20,000 rpm. The  
supernatant was diluted with HS buffer to a volume of 90 mL and 20 g of  
ammonium sulfate was added slowly. The resulting solution was applied onto  
an octyl sepharose column (2.5 cm x 10 cm) (Pharmacia) equilibrated with HS  
25 buffer containing 1.74 M ammonium sulfate at room temperature. After  
washing the column with HS buffer containing 1.74 M ammonium sulfate,  
MTSP7 was eluted in a gradient (1.74 M - 0 ammonium sulfate in HS buffer).  
Active fractions were pooled (60 mL) and dialyzed against HS buffer overnight  
with one buffer change. The dialysate was batch-absorbed onto 20 mL  
30 benzamidine sepharose (Pharmacia) and washed with 4 column volumes HS  
buffer. MTSP 7 was eluted by HS/4 mM benzamidine. The protein appeared

-144-

homogeneous by SDS-PAGE, and the identity of the protein was confirmed by amino acid sequencing.

### EXAMPLE 3

#### 5 Assays for identification of candidate compounds that modulate that activity of an MTSP

##### Assay for identifying inhibitors

The ability of test compounds to act as inhibitors of catalytic activity of an MTSP7 can be assessed in an amidolytic assay. The inhibitor-induced inhibition of amidolytic activity by a recombinant MTSP or the protease domain  
10 portions thereof, can be measured by IC<sub>50</sub> values in such an assay.

The protease domain of MTSP7 expressed in *Pichia pastoris* has been assayed in Costar 96 well tissue culture plates (Corning NY) for inhibition by various test compounds as follows. Approximately 1-10 nM protease is added without inhibitor, or with 100000 nM inhibitor and seven 1:6 dilutions into 1X  
15 direct buffer (29.2 mM Tris, pH 8.4, 29.2 mM Imidazole, 217 mM NaCl (100  $\mu$ L final volume)), and allowed to incubate at room temperature for 30 minutes. 400  $\mu$ M substrate S 2366 (DiaPharma, Westchester, OH) is added and the reaction is monitored in a SpectraMAX Plus microplate reader (Molecular Devices, Sunnyvale CA) by following change in absorbance at 405 nm for 20  
20 minutes at 37°C.

### EXAMPLE 4

#### Other Assays

These assays are described with reference to MTSP1, but such assays can be readily adapted for use with MTSP7.

#### 25 Amidolytic Assay for Determining Inhibition of Serine Protease Activity of Matriptase or MTSP1

The ability of test compounds to act as inhibitors of rMAP catalytic activity was assessed by determining the inhibitor-induced inhibition of amidolytic activity by the MAP, as measured by IC<sub>50</sub> values. The assay buffer  
30 was HBSA (10 mM Hepes, 150 mM sodium chloride, pH 7.4, 0.1% bovine serum albumin). All reagents were from Sigma Chemical Co. (St. Louis, MO), unless otherwise indicated.

-145-

Two  $IC_{50}$  assays (a) one at either 30-minutes or 60-minutes (a 30-minute or a 60-minute preincubation of test compound and enzyme) and (b) one at 0-minutes (no preincubation of test compound and enzyme) were conducted. For the  $IC_{50}$  assay at either 30-minutes or 60-minutes, the following reagents were combined in appropriate wells of a Corning microtiter plate: 50 microliters of HBSA, 50 microliters of the test compound, diluted (covering a broad concentration range) in HBSA (or HBSA alone for uninhibited velocity measurement), and 50 microliters of the rMAP (Corvas International) diluted in buffer, yielding a final enzyme concentration of 250 pM as determined by active site filtration. Following either a 30-minute or a 60-minute incubation at ambient temperature, the assay was initiated by the addition of 50 microliters of the substrate S-2765 (N- $\alpha$ -Benzyloxycarbonyl-D-arginyl-L-glycyl-L-arginine-p-nitroaniline dihydrochloride; DiaPharma Group, Inc.; Franklin, OH) to each well, yielding a final assay volume of 200 microliters and a final substrate concentration of 100  $\mu$ M (about 4-times  $K_m$ ). Before addition to the assay mixture, S-2765 was reconstituted in deionized water and diluted in HBSA. For the  $IC_{50}$  assay at 0 minutes; the same reagents were combined: 50 microliters of HBSA, 50 microliters of the test compound, diluted (covering the identical concentration range) in HBSA (or HBSA alone for uninhibited velocity measurement), and 50 microliters of the substrate S-2765. The assay was initiated by the addition of 50 microliters of rMAP. The final concentrations of all components were identical in both  $IC_{50}$  assays (at 30- or 60- and 0-minute).

The initial velocity of chromogenic substrate hydrolysis was measured in both assays by the change of absorbance at 405 nM using a Thermo Max<sup>®</sup> Kinetic Microplate Reader (Molecular Devices) over a 5 minute period, in which less than 5% of the added substrate was used. The concentration of added inhibitor, which caused a 50% decrease in the initial rate of hydrolysis was defined as the respective  $IC_{50}$  value in each of the two assays (30- or 60-minutes and 0-minute).

### ***In vitro* enzyme assays for specificity determination**

The ability of compounds to act as a selective inhibitor of matriptase activity was assessed by determining the concentration of test compound that

-146-

inhibits the activity of matriptase by 50%, ( $IC_{50}$ ) as described in the above Example, and comparing  $IC_{50}$  value for matriptase to that determined for all or some of the following serine proteases: thrombin, recombinant tissue plasminogen activator (rt-PA), plasmin, activated protein C, chymotrypsin, factor Xa and trypsin.

The buffer used for all assays was HBSA (10 mM HEPES, pH 7.5, 150 mM sodium chloride, 0.1% bovine serum albumin). The assay for  $IC_{50}$  determinations was conducted by combining in appropriate wells of a Corning microtiter plate, 50 microliters of HBSA, 50 microliters of the test compound at a specified concentration (covering a broad concentration range) diluted in HBSA (or HBSA alone for  $V_0$  (uninhibited velocity) measurement), and 50 microliters of the enzyme diluted in HBSA. Following a 30 minute incubation at ambient temperature, 50 microliters of the substrate at the concentrations specified below were added to the wells, yielding a final total volume of 200 microliters.

The initial velocity of chromogenic substrate hydrolysis was measured by the change in absorbance at 405 nm using a Thermo Max<sup>®</sup> Kinetic Microplate Reader over a 5 minute period in which less than 5% of the added substrate was used. The concentration of added inhibitor which caused a 50% decrease in the initial rate of hydrolysis was defined as the  $IC_{50}$  value.

#### Thrombin (fIIa) Assay

Enzyme activity was determined using the chromogenic substrate, Pefachrome t-PA ( $CH_3SO_2$ -D-hexahydrotyrosine-glycyl-L-Arginine-p-nitroaniline, obtained from Pentapharm Ltd.). The substrate was reconstituted in deionized water prior to use. Purified human  $\alpha$ -thrombin was obtained from Enzyme Research Laboratories, Inc. The buffer used for all assays was HBSA (10 mM HEPES, pH 7.5, 150 mM sodium chloride, 0.1% bovine serum albumin).

$IC_{50}$  determinations were conducted where HBSA (50  $\mu$ L),  $\alpha$ -thrombin (50  $\mu$ l) (the final enzyme concentration is 0.5 nM) and inhibitor (50  $\mu$ l) (covering a broad concentration range), were combined in appropriate wells and incubated for 30 minutes at room temperature prior to the addition of substrate Pefachrome-t-PA (50  $\mu$ l) (the final substrate concentration is 250  $\mu$ M, about 5 times  $K_m$ ). The initial velocity of Pefachrome t-PA hydrolysis was measured by

-147-

the change in absorbance at 405 nm using a Thermo Max<sup>®</sup> Kinetic Microplate Reader over a 5 minute period in which less than 5% of the added substrate was used. The concentration of added inhibitor which caused a 50% decrease in the initial rate of hydrolysis was defined as the IC<sub>50</sub> value.

**5 Factor Xa**

Factor Xa catalytic activity was determined using the chromogenic substrate S-2765 (N-benzyloxycarbonyl-D-arginine-L-glycine-L-arginine-p-nitro-aniline), obtained from DiaPharma Group (Franklin, OH). All substrates were reconstituted in deionized water prior to use. The final concentration of S-2765  
10 was 250  $\mu$ M (about 5-times K<sub>m</sub>). Purified human Factor X was obtained from Enzyme Research Laboratories, Inc. (South Bend, IN) and Factor Xa (FXa) was activated and prepared from it as described [Bock, P.E., Craig, P.A., Olson, S.T., and Singh, P. *Arch. Biochem. Biophys.* 273:375-388 (1989)]. The enzyme was diluted into HBSA prior to assay in which the final concentration was 0.25 nM.

**15 Recombinant tissue plasminogen activator (rt-PA) Assay**

rt-PA catalytic activity was determined using the substrate, Pefachrome t-PA (CH<sub>3</sub>SO<sub>2</sub>-D-hexahydrotyrosine-glycyl-L-arginine-p-nitroaniline, obtained from Pentapharm Ltd.). The substrate was made up in deionized water followed by dilution in HBSA prior to the assay in which the final concentration was 500  
20 micromolar (about 3-times K<sub>m</sub>). Human rt-PA (Activase<sup>®</sup>) was obtained from Genentech Inc. The enzyme was reconstituted in deionized water and diluted into HBSA prior to the assay in which the final concentration was 1.0 nM.

**Plasmin Assay**

Plasmin catalytic activity was determined using the chromogenic  
25 substrate, S-2366 [L-pyroglutamyl-L-prolyl-L-arginine-p-nitroaniline hydrochloride], which was obtained from DiaPharma group. The substrate was made up in deionized water followed by dilution in HBSA prior to the assay in which the final concentration was 300 micromolar (about 2.5-times K<sub>m</sub>). Purified human plasmin was obtained from Enzyme Research Laboratories, Inc.  
30 The enzyme was diluted into HBSA prior to assay in which the final concentration was 1.0 nM.



-148-

**Activated Protein C (aPC) Assay**

aPC catalytic activity was determined using the chromogenic substrate, Pefachrome PC (delta-carbobenzloxy-D-lysine-L-prolyl-L-arginine-p-nitroaniline dihydrochloride), obtained from Pentapharm Ltd.). The substrate was made up  
5 in deionized water followed by dilution in HBSA prior to the assay in which the final concentration was 400 micromolar (about 3-times  $K_m$ ). Purified human aPC was obtained from Hematologic Technologies, Inc. The enzyme was diluted into HBSA prior to assay in which the final concentration was 1.0 nM.

**Chymotrypsin Assay**

10 Chymotrypsin catalytic activity was determined using the chromogenic substrate, S-2586 (methoxy-succinyl-L-arginine-L-prolyl-L-tyrosyl-p-nitroanilide), which was obtained from DiaPharma Group. The substrate was made up in deionized water followed by dilution in HBSA prior to the assay in which the final concentration was 100 micromolar (about 9-times  $K_m$ ). Purified (3X-crystallized;  
15 CDI) bovine pancreatic alpha-chymotrypsin was obtained from Worthington Biochemical Corp. The enzyme was reconstituted in deionized water and diluted into HBSA prior to assay in which the final concentration was 0.5 nM.

**Trypsin Assay**

Trypsin catalytic activity was determined using the chromogenic  
20 substrate, S-2222 (benzoyl-L-isoleucine-L-glutamic acid-[gamma-methyl ester]-L-arginine-p-nitroanilide), which was obtained from DiaPharma Group. The substrate was made up in deionized water followed by dilution in HBSA prior to the assay in which the final concentration was 250 micromolar (about 4-times  $K_m$ ). Purified (3X-crystallized; TRL3) bovine pancreatic trypsin was obtained  
25 from Worthington Biochemical Corp. The enzyme was reconstituted in deionized water and diluted into HBSA prior to assay in which the final concentration was 0.5 nM.

Since modifications will be apparent to those of skill in this art, it is  
30 intended that this invention be limited only by the scope of the appended claims.

-149-

**WHAT IS CLAIMED IS:**

1. A substantially purified single or two chain polypeptide, comprising the protease domain of a type-II membrane-type serine protease 7 (MTSP7) or a catalytically active portion thereof.
- 5 2. The polypeptide of claim 1 that is an activated two chain protein.
3. A polypeptide of claim 1 selected from the group consisting of a polypeptide that comprises a sequence of amino acids encoded by the sequence of nucleotides set forth in SEQ ID No. 15; polypeptide that comprises a sequence of amino acids encoded by the sequence of nucleotides  
10 set forth in SEQ ID No. 17;  
a polypeptide that comprises a sequence of amino acids encoded by a sequence of nucleotides that hybridizes under conditions of high stringency to the sequence of nucleotides set forth in SEQ ID No. 15 or 17;  
a polypeptide that comprises the sequence of amino acids set  
15 forth as amino acids 206-438 of SEQ ID No. 16;  
a polypeptide that comprises a sequence of amino acids having at least about 90% sequence identity with the sequence of amino acids set forth in SEQ ID No. 16 or 18; and  
a polypeptide that is encoded by a sequence of nucleotides that is  
20 a splice variant of the sequence set forth in SEQ ID No. 15.
4. The polypeptide of claim 1, wherein:  
the MTSP7 portion of the polypeptide consists essentially of the protease domain of the MTSP7 or a catalytically active portion thereof.
5. The substantially purified polypeptide of claim 1, wherein the  
25 MTSP7 is a human polypeptide.
6. The substantially purified polypeptide of claim 1 that consists essentially of the protease domain of MTSP7 or a catalytically active portion of the protease domain of MTSP7.
7. The substantially purified polypeptide of claim 3 that consists  
30 essentially of the protease domain of MTSP7 or a catalytically active portion of the protease domain of MTSP7.

-150-

8. The substantially purified polypeptide of claim 1 that comprises the sequence of amino acids set forth in SEQ ID No. 16.

9. The substantially purified polypeptide of claim 1 that comprises the sequence of amino acids set forth in SEQ ID No. 18.

5 10. The substantially purified polypeptide of claim 1, wherein the protease domain comprises the sequence of amino acids set forth as amino acids 206-438 of SEQ ID No. 16.

11. The substantially purified polypeptide of claim 1 that has at least about 60%, 80%, 90% or 95% sequence identity with a polypeptide that  
10 comprises the sequence of amino acids set forth as SEQ ID No. 16 or as the sequence of amino acids set forth as SEQ ID No. 18, wherein the polypeptide is a protease.

12. A polypeptide of claim 1, wherein the protease domain portion is encoded by a nucleic acid molecule that hybridizes under conditions of high  
15 stringency along at least 70% of its full length to a nucleic acid molecule comprising a sequence of nucleotides set forth in SEQ ID No. 15 or at least one domain thereof or a catalytically active portion of the domain.

13. The polypeptide of claim 12, wherein the domain is the protease domain

20 14. The polypeptide of claim 1, wherein:  
the polypeptide does not comprise the complete sequence set forth in SEQ ID No. 16; and  
the MTSP7 portion of the polypeptide consists essentially of the protease domain of the MTSP7 or a catalytically active portion thereof.

25 15. A polypeptide of claim 3 that is a mutein, wherein:  
up to about 40% of the amino acids are replaced with another amino acid;

and the resulting polypeptide is a single chain or two chain polypeptide that has catalytic activity of at least 10% of the unmutated polypeptide.

30 16. The polypeptide of claim 15, wherein up to about 10% of the amino acids are replaced with another amino acid.

-151-

17. The polypeptide of claim 15, wherein the resulting polypeptide is a single chain or two chain polypeptide and has catalytic activity of at least 50% of the unmutated polypeptide.

18. The polypeptide of claim 15, wherein a free Cysteine in the  
5 protease domain is replaced with another amino acid.

19. The polypeptide of claim 18, wherein the replacing amino acid is a serine.

20. A nucleic acid molecule, comprising a sequence of nucleotides that encodes the polypeptide of claim 1.

10 21. A nucleic acid molecule, comprising a sequence of nucleotides that encodes the polypeptide of claim 3.

22. The nucleic acid molecule of claim 20 that comprises a sequence of nucleotides selected from the group consisting of:

- 15 (a) a sequence of nucleotides set forth in SEQ ID No. 15 or 17;  
(b) a sequence of nucleotides that hybridizes under high stringency along its length to the sequence of nucleotides set forth in SEQ ID No. 15 or 17;  
(c) degenerate codons of (a) or (b).

23. An isolated nucleic molecule that encodes a mutein of claim 15.

24. A vector comprising the nucleic acid molecule of claim 20.

20 25. The vector of claim 24 that is an expression vector.

26. The vector of claim 24 that is a eukaryotic vector.

27. The vector of claim 25 that includes a sequence of nucleotides that directs secretion of any polypeptide encoded by a sequence of nucleotides operatively linked thereto.

25 28. The vector of claim 24 that is a *Pichia* vector or an *E. coli* vector.

29. A cell, comprising the vector of claim 24.

30. The cell of claim 29 that is a prokaryotic cell.

31. The cells of claim 29 that is a eukaryotic cell.

32. The cell of claim 29 that is selected from among a bacterial cell, a  
30 yeast cell, a plant cell, an insect cell and an animal cell.

33. The cell of claim 29 that is a mammalian cell.

34. A nucleic acid molecule encoding a polypeptide of claim 6.

-152-

35. A vector, comprising nucleic acid molecule of claim 34.

36. A cell, comprising the vector of claim 35.

37. A recombinant non-human animal, wherein an endogenous gene that encodes a polypeptide of claim 1 has been deleted or inactivated by homologous recombination or insertional mutagenesis of the animal or an ancestor thereof.

38. A method for producing a polypeptide that contains a protease domain of an MTSP7 polypeptide, comprising:

culturing the cell of claim 29 under conditions whereby the encoded polypeptide is expressed by the cell; and recovering the expressed polypeptide.

39. The method of claim 38, wherein the polypeptide is secreted into the culture medium.

40. The method of claim 38, wherein the cell is a *Pichia* cell.

41. The method of claim 38, wherein the polypeptide is expressed in the cytoplasm of the host cell.

42. A method for producing a polypeptide that contains a protease domain of a polypeptide, comprising:

culturing the cell of claim 36 under conditions whereby the encoded polypeptide is expressed by the cell; and recovering the expressed polypeptide.

43. An antisense nucleic acid molecule that comprises at least 14 contiguous nucleotides or modified nucleotides that are complementary to a contiguous sequence of nucleotides encoding the protease domain of an MTSP7 of claim 1; or

comprises at least 16 contiguous nucleotides or modified nucleotides that are complementary to a contiguous sequence of nucleotides encoding the protease domain of an MTSP7 of claim 1; or

comprises at least 30 contiguous nucleotides or modified nucleotides that are complementary to a contiguous sequence of nucleotides encoding the protease domain of an MTSP7 of claim 1.

-153-

44. The antisense molecule of claim 43 that includes a contiguous sequence of nucleotides set forth in SEQ ID No. 15.

45. A double-stranded RNA (dsRNA) molecule that comprises at least about 21 contiguous nucleotides or modified nucleotides from the sequence of  
5 nucleotides encoding the MTSP7 of claim 1.

46. An antibody that specifically binds to the single chain form and/or two-chain form of a protease domain of the polypeptide of claim 1, or a fragment or derivative of the antibody containing a binding domain thereof, wherein the antibody is a polyclonal antibody or a monoclonal antibody.

10 47. The antibody of claim 46 that inhibits the enzymatic activity of the polypeptide.

48. An antibody that specifically binds to the single chain form and/or two-chain form of a protease domain of the polypeptide of claim 3, or a fragment or derivative of the antibody containing a binding domain thereof,  
15 wherein the antibody is a polyclonal antibody or a monoclonal antibody.

49. An antibody that specifically binds to the single chain form and/or two-chain form of a protease domain of the polypeptide of claim 6, or a fragment or derivative of the antibody containing a binding domain thereof, wherein the antibody is a polyclonal antibody or a monoclonal antibody.

20 50. A conjugate, comprising:  
a) a polypeptide of claim 1, and  
b) a targeting agent linked to the polypeptide directly or via a linker.

51. The conjugate of claim 50, wherein the targeting agent permits  
25 i) affinity isolation or purification of the conjugate;  
ii) attachment of the conjugate to a surface;  
iii) detection of the conjugate; or  
iv) targeted delivery to a selected tissue or cell.

-154-

52. A conjugate, comprising:
- a) a polypeptide of claim 3; and
  - b) a targeting agent linked to the polypeptide directly or via a linker.
- 5 53. The conjugate of claim 52, wherein the targeting agent permits
- i) affinity isolation or purification of the conjugate;
  - ii) attachment of the conjugate to a surface;
  - iii) detection of the conjugate; or
  - iv) targeted delivery to a selected tissue or cell.
- 10 54. A conjugate, comprising:
- a) a polypeptide of claim 6; and
  - b) a targeting agent linked to the polypeptide directly or via a linker.
- 15 55. The conjugate of claim 54, wherein the targeting agent permits
- i) affinity isolation or purification of the conjugate;
  - ii) attachment of the conjugate to a surface;
  - iii) detection of the conjugate; or
  - iv) targeted delivery to a selected tissue or cell.
- 20 56. A combination, comprising:
- a) an agent or treatment that inhibits the catalytic activity of the polypeptide of claim 1; and
  - b) another treatment or agent selected from anti-tumor and anti-angiogenic treatments and agents.
- 25 57. The combination of claim 56, wherein the inhibitor and the anti-tumor and/or anti-angiogenic agent are formulated in a single pharmaceutical composition or each is formulated in separate pharmaceutical compositions.
58. The combination of claim 56, wherein the inhibitor is selected from antibodies and antisense oligonucleotides and double-stranded RNA (dsRNA).
- 30 59. A solid support comprising two or more polypeptides of claim 1 linked thereto either directly or via a linker.
60. The support of claim 59, wherein the polypeptides comprise an array.

-155-

61. The support of claim 59, wherein the polypeptides comprise a plurality of different protease domains.

62. A solid support comprising two or more nucleic acid molecules of claim 20 or oligonucleotides portions thereof linked thereto either directly or via  
5 a linker, wherein the oligonucleotides contain at least 16 nucleotides.

63. The support of claim 62, wherein the nucleic acid molecules comprise an array.

64. The support of claim 62, wherein the nucleic acid molecules comprise a plurality of molecules that encode different protease domains.

10 65. A method for identifying compounds that modulate the protease activity of a polypeptide, comprising:

contacting a polypeptide of claim 1 with a substrate that is proteolytically cleaved by the polypeptide, and, either simultaneously, before or after, adding a test compound or plurality thereof;

15 measuring the amount of substrate cleaved in the presence of the test compound; and

selecting compounds that change the amount of substrate cleaved compared to a control, whereby compounds that modulate the activity of the polypeptide are identified.

20 66. The method of claim 65, wherein the test compounds are small molecules, peptides, peptidomimetics, natural products, antibodies or fragments thereof that modulate the activity of the polypeptide.

67. The method of claim 65, wherein a plurality of the test substances are screened simultaneously.

25 68. The method of claim 65, wherein the polypeptide consists essentially of a polypeptide encoded by a sequence of nucleotides selected from the group consisting of a sequence of nucleotides that:

(a) is set forth in SEQ ID No. 17;

(b) hybridizes under conditions of high stringency to nucleic  
30 acid complementary to an mRNA transcript present in a mammalian cell that encodes MTSP7 encoded by (a);

(c) encodes a splice variant of (a) or (b); and



-156-

(d) comprises degenerate codons of the sequences of nucleotides of (a), (b) or (c).

69. The method of claim 65, wherein the polypeptide consists essentially of a polypeptide selected from the group consisting of:

5 a polypeptide that comprises a sequence of amino acids encoded by the sequence of nucleotides set forth in SEQ ID No. 15; polypeptide that comprises a sequence of amino acids encoded by the sequence of nucleotides set forth in SEQ ID No. 17;

10 a polypeptide that comprises a sequence of amino acids encoded by a sequence of nucleotides that hybridizes under conditions of high stringency to the sequence of nucleotides set forth in SEQ ID No. 15 or 17;

a polypeptide that comprises the sequence of amino acids set forth as amino acids 206-438 of SEQ ID No. 16;

15 a polypeptide that comprises a sequence of amino acids having at least about 90% sequence identity with the sequence of amino acids set forth in SEQ ID No. 16 or 18; and

a polypeptide that is encoded by a sequence of nucleotides that is a splice variant of the sequence set forth in SEQ ID No. 15.

20 70. The method of claim 65, wherein the change in the amount of substrate cleaved is assessed by comparing the amount of substrate cleaved in the presence of the test compound with the amount of substrate cleaved in the absence of the test compound.

71. The method of claim 67, wherein a plurality of the polypeptides are linked to a solid support, either directly or via a linker.

25 72. The method of claim 71, wherein the polypeptides comprise an array.

73. A method of identifying a compound that specifically binds to a single-chain and/or two-chain protease domain and/or to single or two-chain full-length polypeptide, comprising:

30 contacting a polypeptide of claim 1 with a test compound or plurality thereof under conditions conducive to binding thereof; and

-157-

identifying compounds that specifically bind to the polypeptide single chain protease domain, or two chain form thereof, the full length or two chain form of the full length polypeptide or compounds that inhibit binding of a compound known to bind to the polypeptide single chain protease domain or two chain form thereof or the two chain form of the full length polypeptide, wherein the known compound is contacted with the polypeptide before, simultaneously with or after the test compound.

74. The method of claim 73, wherein the polypeptide is linked either directly or indirectly via a linker to a solid support.

10 75. The method of claim 73, wherein the test compounds are small molecules, peptides, peptidomimetics, natural products, antibodies or fragments thereof.

76. The method of claim 73, wherein a plurality of the test substances are screened simultaneously.

15 77. The method of claim 73, wherein a plurality of the polypeptides are linked to a solid support.

78. The method of claim 73, wherein the polypeptide consists essentially of a polypeptide encoded by a sequence of nucleotides that:

- (a) is set forth in SEQ ID No. 17;
- 20 (b) hybridizes under conditions of moderate or high stringency to nucleic acid complementary to an mRNA transcript present in a mammalian cell that encodes a MTSP7 encoded by (a);
- (c) encodes a splice variant of (a) or (b); and
- (d) comprises degenerate codons of the sequences of
- 25 nucleotides of (a), (b) or (c).

79. A method for identifying activators of the zymogen form of an MTSP7, comprising:

contacting a zymogen form of the polypeptide of claim 1 with a substrate of the activated form of the polypeptide;

30 add a test compound, wherein the test compound is added before, after or simultaneously with the addition of the substrate; and

-158-

detecting cleavage of the substrate, thereby identifying compounds that activate the zymogen.

80. The method of claim 77, wherein the substrate is a chromogenic substrate.

5 81. The method of claim 77, wherein the substrate is a L-pyroglutamyl-L-prolyl-L-arginine-p-nitroaniline hydrochloride.

82. The method of claim 77, wherein the test compound is a small molecule, a nucleic acid or a polypeptide.

83. A method for treating or preventing a neoplastic disease, in a  
10 mammal, comprising administering to a mammal an effective amount of an inhibitor of a polypeptide of claim 1.

84. The method of claim 83, wherein the inhibitor is an antibody that specifically binds to the polypeptide, or a fragment or derivative of the antibody containing a binding domain thereof, wherein the antibody is a polyclonal  
15 antibody or a monoclonal antibody.

85. A method for treating or preventing a neoplastic disease, in a mammal, comprising administering to a mammal an effective amount of an inhibitor of a polypeptide of claim 3.

86. A method for treating or preventing a neoplastic disease, in a  
20 mammal, comprising administering to a mammal an effective amount of an inhibitor of a polypeptide of claim 6.

87. A method of inhibiting tumor initiation, growth or progression or treating a malignant or pre-malignant condition, comprising administering an agent that inhibits activation of the zymogen form of a polypeptide of claim 1 or  
25 an activity of the activated form.

88. The method of claim 87, wherein the condition is a condition of the breast, cervix, prostate, lung, ovary or colon.

89. The method of claim 87, wherein the agent is an antisense oligonucleotide, double-stranded RNA (dsRNA) or an antibody.

30 90. The method of claim 87, further comprising administering another treatment or agent selected from anti-tumor and anti-angiogenic treatments or agents.

-159-

91. A method of inhibiting tumor initiation, growth or progression or treating a malignant or pre-malignant condition, comprising administering an agent that inhibits activation of the zymogen form of a polypeptide of claim 3 or an activity of the activated form.

5 92. The method of claim 91, wherein the condition is a condition of the breast, cervix, prostate, lung, ovary or colon.

93. The method of claim 91, wherein the agent is an antisense oligonucleotide, double-stranded RNA (dsRNA) or an antibody.

10 94. The method of claim 91, further comprising administering another treatment or agent selected from anti-tumor and anti-angiogenic treatments or agents.

95. A method of identifying a compound that binds to the single-chain or two-chain form of a polypeptide of claim 1, comprising:

- 15       contacting a test compound with both forms;  
      determining to which form the compound binds; and  
      if it binds to a form of polypeptide, further determining whether the compound has at least one of the following properties:
- (i) inhibits activation of the single-chain zymogen form of polypeptide;
  - 20       (ii) inhibits activity of the two-chain or single-chain form; and
  - (iii) inhibits dimerization of the polypeptide.

96. The method of claim 95, wherein both forms consist essentially of the protease domain produced by cleavage between the arginine and isoleucine in either the single- or two-chain form.

25 97. A method of identifying a compound that binds to the single-chain or two-chain form of a polypeptide of claim 3, comprising:

      contacting a test compound with both forms;  
      determining to which form the compound binds; and  
      if it binds to a form of polypeptide, further determining whether the

30 compound has at least one of the following properties:

- (i) inhibits activation of the single-chain zymogen form of polypeptide;

-160-

(ii) inhibits activity of the two-chain or single-chain form; and

(iii) inhibits dimerization of the polypeptide.

98. The method of claim 97, wherein both forms consist essentially of the protease domain produced by cleavage between the R and I in either the single-  
5 or two-chain form.

99. A method of detecting neoplastic disease, comprising: detecting a polypeptide that comprises a polypeptide of claim 1 in a biological sample, wherein the amount detected differs from the amount of polypeptide detected from a subject who does not have neoplastic disease.

100. The method of claim 99, wherein the biological sample is selected from the group consisting of blood, urine, saliva, tears, synovial fluid, sweat, interstitial fluid, cerebrospinal fluid, ascites fluid, tumor tissue biopsy and circulating tumor cells.

101. A method of detecting neoplastic disease, comprising: detecting a  
15 polypeptide that comprises a polypeptide of claim 3 in a biological sample, wherein the amount detected differs from the amount of polypeptide detected from a subject who does not have neoplastic disease.

102. The method of claim 101, wherein the biological sample is selected from the group consisting of blood, urine, saliva, tears, synovial fluid, sweat, interstitial fluid, cerebrospinal fluid, ascites fluid, tumor tissue biopsy and  
20 circulating tumor cells.

103. A method of detecting neoplastic disease, comprising: detecting a polypeptide that comprises a polypeptide of claim 6 in a biological sample, wherein the amount detected differs from the amount of polypeptide detected  
25 from a subject who does not have neoplastic disease.

104. The method of claim 103, wherein the biological sample is selected from the group consisting of blood, urine, saliva, tears, synovial fluid, sweat, interstitial fluid, cerebrospinal fluid, ascites fluid, tumor tissue biopsy and circulating tumor cells.

30 105. A method of diagnosing the presence of a pre-malignant lesion, a malignancy, or other pathologic condition in a subject, comprising:  
obtaining a biological sample from the subject; and

-161-

exposing it to a detectable agent that binds to a two-chain and/or single-chain form of a polypeptide of claim 1, wherein the pathological condition is characterized by the presence or absence of the two-chain or single-chain form.

106. A method of diagnosing the presence of a pre-malignant lesion, a malignancy, or other pathologic condition in a subject, comprising:

obtaining a biological sample from the subject; and

exposing it to a detectable agent that binds to a two-chain and/or single-chain form of a polypeptide of claim 3, wherein the pathological condition is characterized by the presence or absence of the two-chain or single-chain form.

107. A method of diagnosing the presence of a pre-malignant lesion, a malignancy, or other pathologic condition in a subject, comprising:

obtaining a biological sample from the subject; and

exposing it to a detectable agent that binds to a two-chain and/or single-chain form of a polypeptide of claim 6, wherein the pathological condition is characterized by the presence or absence of the two-chain or single-chain form.

108. A method of monitoring tumor progress and/or therapeutic effectiveness, comprising detecting and/or quantifying the level of a polypeptide of claim 1 in a body tissue or fluid sample.

109. The method of claim 108, wherein the tumor is a tumor of the breast, cervix, prostate, lung, ovary or colon.

110. The method of claim 108, wherein the body fluid is blood, urine, sweat, saliva, cerebrospinal fluid and synovial fluid.

111. A method of monitoring tumor progress and/or therapeutic effectiveness, comprising detecting and/or quantifying the level of a polypeptide of claim 3 in a body tissue or fluid sample.

112. The method of claim 111, wherein the tumor is a tumor of the breast, cervix, prostate, lung, ovary or colon.

113. The method of claim 111, wherein the body fluid is blood, urine, sweat, saliva, cerebrospinal fluid and synovial fluid.


114. A method of monitoring tumor progress and/or therapeutic effectiveness, comprising detecting and/or quantifying the level of a polypeptide of claim 6 in a body tissue or fluid sample.

-162-

115. The method of claim 114, wherein the tumor is a tumor of the breast, cervix, prostate, lung, ovary or colon.

116. The method of claim 114, wherein the body fluid is blood, urine, sweat, saliva, cerebrospinal fluid and synovial fluid.

Domain organization and amino acid sequence of MTSP7



10 20 30 40 50 60  
 MMYTPVEFSEAEFSRAEYQRKQFWDVRLAFTLAIVAIIIGIIVTHFWEDDKSFY  
 70 80 90 100 110 120  
 YLASFKVTNIKYKENGIRSSREFIERSHQIERMMSRIFRHSVGGRFKSHVIKLSPE  
 130 140 150 160 170 180  
 QGVDILIVLIFRYPSTDSAEQIKKIEKALYQSLTKQLSLTINKPSFRLTPIDSKMRN  
 190 200 210 220 230 240  
 LLNSRCGIRMTSSNMPLPASSSTQRIVQRETAMEGEWPQASLQIGSGHQCGASLISN  
 250 260 270 280 290 300  
 TWLLTAHCFWKNKDPQTQWIAITFGATITPPAVKRNVRKIILHENYHRETNENDIALVQLS  
 310 320 330 340 350 360  
 TGVEFSNIVQVRVCLPDSSI KLPKTSVFVTGFGSIVDDGPIONTLRQARVETISTDVCNR  
 370 380 390 400 410 420  
 KDVYDGLITPGMLCAGFMECKIDACKGDSGGPLVVDNHDWYIVGIVSWGQSCALPKKPG  
 430  
 VYTRVTKYRDWIASKTGM\*

↓ = protease cleavage site

FIG. 1



10 20 30 40 50 60  
AGATCAGATGGCGACTGAATAGAAAGCTGCCCCAGTCCTGGGTTTCATGATGTACACACCTG  
TCTAGTCTACCGCTGACTTATCTTCGACGGGGTCAGGACCCAAGTACTACATGTGTGGAC

70 80 90 100 110 120  
TTGAATTTTCAGAAGCTGAATTCTCAGGAGCTGAATATCAAAGAAAGCAGCAATTTTGGG  
AACTTAAAAGTCTTCGACTTAAGAGTGCTCGACTTATAGTTTCTTTCGTCGTTAAAACCC

130 140 150 160 170 180  
ACTCAGTACGGCTAGCTCTTTTCACATTAGCAATTGTAGCAATCATAGGAATTGCAATTG  
TGAGTCATGCCGATCGAGAAAAGTGAATCGTTAACATCGTTAGTATCCTTAACGTTAAC

190 200 210 220 230 240  
GTATTGTTACTCATTTTGTGTTGAGGATGATAAGTCTTTCTATTACCTTGCCTCTTTTA  
CATAACAATGAGTAAAACAACAACCTCCTACTATTAGAAAAGATAATGGAACGGAGAAAAT

250 260 270 280 290 300  
AAGTCACAAATATCAAATATAAAGAAAATTATGGCATAAGATCTTCAAGAGAGTTTATAG  
TTCAGTGTTTATAGTTTATATTTCTTTTAATACCGTATTCTAGAAGTTCTCTCAAATATC

310 320 330 340 350 360  
AAAGGAGTCATCAGATTGAAAGAATGATGTCTAGGATATTTGACATTCTTCTGTAGGCG  
TTTCTCAGTAGTCTAACTTTCTTACTACAGATCCTATAAAGCTGTAAGAAGACATCCGC

370 380 390 400 410 420  
GTCGATTTATCAAATCTCATGTTATCAAATTAAGTCCAGATGAACAAGGTGTGGATATTC  
CAGCTAAATAGTTTAGAGTACAATAGTTTAATTCAGGTCTACTTGTTCACACCTATAAG

430 440 450 460 470 480  
TTATAGTGCTCATATTTGATACCCATCTACTGATAGTGCTGAACAAATCAAGAAAAAAA  
AATATCAGGAGTATAAAGCTATGGGTAGATGACTATCAGACTTGTTTAGTTCTTTTTTT

490 500 510 520 530 540  
TTGAAAAGGCTTTATATCAAAGTTTGAAGACCAAAACAATTGTCTTTGACCATAAACAAC  
AACTTTTCCGAAATATAGTTTCAAACCTTCTGGTTTGTAAACAGAACTGGTATTTGTTTG

550 560 570 580 590 600  
CATCATTTAGACTCACACCTATTGACAGCAAAAAGATGAGGAATCTTCTCAACAGTCGCT  
GTAGTAAATCTGAGTGTGGATAACTGTGTTTTCTACTCCTTAGAAGAGTTGTGAGCGA

610 620 630 640 650 660  
GTGGAATAAGGATGACATCTTCAAACATGCCATTACCAGCATCCTCTTCTACTCAAAGAA  
CACCTTATTCCTACTGTAGAAGTTTGTACGGTAATGGTCGTAGGAGAAGATGAGTTTCTT

670 680 690 700 710 720  
TTGTCCAAGGAAGGGAACAGCTATGGAAGGGGAATGGCCATGGCAGGCCAGCCTCCAGC  
AACAGGTTCCCTCCCTTTGTGATACCTTCCCTTACCGGTACCGTCCGGTCGGAGGTCC

730 740 750 760 770 780  
TCATAGGGTCAGGCCATCAGTGTGGAGCCAGCCTCATCAGTAACACATGGCTGCTCACAG  
AGTATCCCAGTCCGGTAGTCACACCTCGGTCCGAGTAGTCATTGTGTACCGACGAGTGTG

790 800 810 820 830 840  
CAGCTCACTGCTTTTGGAAAAATAAAGACCCAACTCAATGGATTGCTACTTTTGGTGCAA  
GTCGAGTGACGAAAACCTTTTATTTCTGGGTTGAGTTACCTAACGATGAAAACCACGTT

850 860 870 880 890 900  
CTATAACACCACCCGAGTGAAACGAAATGTGAGGAAAATTATTCTTCATGAGAATTACC  
GATATTGTGGTGGGCGTCACCTTGTCTTACACTCCTTTTAATAAGAAGTACTCTTAATGG

910 920 930 940 950 960  
ATAGAGAAACAAATGAAAATGACATTGCTTTGGTTTCAGCTCTCTACTGGAGTTGAGTTT  
TATCTCTTTGTTTACTTTTACTGTAACGAAACCAAGTCGAGAGATGACCTCAACTCAAAA

970 980 990 1000 1010 1020  
CAAATATAGTCCAGAGAGTTTGCCTCCCAGACTCATCTATAAAGTTGCCACCTAAAACAA  
GTTTATATCAGGTCTCTCAAACGGAGGGTCTGAGTAGATATTTCAACGGTGGATTTTGT

1030 1040 1050 1060 1070 1080  
GTGTGTTTCGTACAGGATTTGGATCCATTGTAGATGATGGACCTATACAAAATACACTTC  
CACACAAGCAGTGTCTAAACCTAGGTAACATCTACTACCTGGATATGTTTATGTGAAG

1090 1100 1110 1120 1130 1140  
GGCAAGCCAGAGTGGAAACCATAAGCACTGATGTGTGTAAACAGAAAGGATGTGTATGATG  
CCGTTCCGGTCTCACCTTTGGTATTCGTGACTACACACATTGTCTTCTACACATACTAC

1150 1160 1170 1180 1190 1200  
GCCTGATAACTCCAGGAATGTTATGTGCTGGATTCATGGAAGGAAAAATAGATGCATGTA  
CGGACTATTGAGGTCTTACAATACACGACCTAAGTACCTTCTTTTATCTACGTACAT

1210 1220 1230 1240 1250 1260  
AGGCAGATTCTGGTGGACCTCTGGTTTATGATAATCATGACATCTGGTACATTGTAGGTA  
TCCCTCTAAGACCACCTGGAGACCAATACTATTAGTACTGTAGACCATGTAACATCCAT

1270 1280 1290 1300 1310 1320  
TAGTAAGTTGGGGACAATCATGTGCACTTCCCAAAAACCTGGAGTCTACACCAGAGTAA  
ATCATTCAACCCCTGTTAGTACACGTGAAGGGTTTTTGGACCTCAGATGTGGTCTCATT

1330 1340 1350 1360 1370 1380  
CTAAGTATCGAGATTGGATTGCCTCAAAGACTGGTATGTAGTGTGGATTGTCCATGAGTT  
GATTCATAGCTCTAACCTAACGGAGTTTCTGACCATACATCACACCTAACAGGTACTCAA

1390 1400 1410 1420 1430 1440  
ATACACATGGCACACAGAGCTGATACTCCTGCGTATTTTGTATTGTTTAAATTCAATTAC  
TATGTGTACCGTGTGTCTCGACTATGAGGACGCATAAAACATAACAAATTTAAGTAAATG

1450 1460 1470 1480 1490 1500  
TTTGGATTAGTGCTTTTGGCTAGATGTCAAGAAGCCCTTCAGACCCAGACAAATCTAATAT  
AAACCTAATCACGAAAACGATCTACAGTTCTTCGGGAAGTCTGGGTCTGTTTAGATTATA

1510 1520 1530 1540 1550 1560  
CCTGAGGTGGCCTTTACATACGTAGGACCAAACCTCTCTACCATGAGGGAAGAAGACAC  
GGACTCCACCGGAAATGTATGCATCCTGGTTTGGGAGAGATGGTACTCCCTTCTTCTGTG

1570 1580 1590 1600 1610 1620  
AGCAAATGACAGACAGCACCTATTCCTTACTCACAAGGGAACTGCTTGTGATACTTCCT  
TCGTTTACTGTCTGTCTGGATAAGGAATGAGTGTTCCTTTGACGAACACTATGAAGGA

1630 1640 1650 1660 1670 1680  
AATAAGATAAATAAGTGGTTTCCCTCAATTGAAGACAGGAACATCATTTTCCACAGGATA  
TTATTCTATTTATTCACCAAAGGGAGTTAACTTCTGTCTTGTAGTAAAAGGTGTCTTAT

1690 1700 1710 1720 1730 1740  
TGAAGAGCTGCCAGTAATGCCAAAATCTTACCTCATATAATACCTGGAGCATGTGAGATT  
ACTTCTCGACGGTCATTACGGTTTTTAGAATGGAGTATATTATGGACCTCGTACACTCTAA

1750 1760 1770 1780 1790 1800  
CTTCTAGTGAAAAAGAACAGTCTTCCCTGAAGACTCAGGGCTTCAACATTCTAGAACTGA  
GAAGATCACTTTTTCTTGTCTCAGAAGGGACTTCTGAGTCCCGAAGTTGTAAGATCTTGACT

1810 1820 1830 1840 1850 1860  
TAAGTGGACCTTCAGTGTGCAAGAATGGAGAAGCATGGGATTTGCATTATGACTTGAAC  
ATTACCTGGAAGTCACACGTTCTTACCTCTTCGTACCCTAAACGTAATACTGAACCTGA

1870 1880 1890 1900 1910 1920  
GGGCTTATATCTAATAATACAGAGCACTATCACTAACCTCAACAGTTGACATTTTAAAG  
CCCGAATATAGATTATTATGTCTCGTGATAGTGATTGGAGTTGTCAACTGTAAAATTTTC

1930	1940	1950	1960	1970	1980
TTTTTAAATGTATCTGAACTTGCTGTAAACACAGTGTATAACTCAAGCACTAGCTTCAG					
AAAAATTTACATAGACTTGAACGACAATTGTGTCACAATATTGAGTTCGTGATCGAAGTC					

1990	2000	2010	2020	2030	2040
GAAGCATGTTGTGTTGTTAAGAAGCTTTTCTGATTATTCTTTAACAGCATCTTGCCATC					
CTTCGTACAACACAACAATTCTTCGAAAAGACTAAATAAGAAATTGTCGTAGAACGGTAG					

2050	2060	2070	2080	2090	2100
TATATGTTAGTAGCAGTTGGCCCAGAAAGGACAAAAAAAAAAAAAAAAAAAAAAAAAAAA					
ATATACAATCATCGTCAACCGGGTCTTCTGTTTTTTTTTTTTTTTTTTTTTTTTTTTTT					

-1-

## SEQUENCE LISTING

<110> Edwin L. Madison  
Edgar O. Ong

<120> NUCLEIC ACID MOLECULES ENCODING TRANSMEMBRANE SERINE PROTEASE 7, THE  
ENCODED POLYPEPTIDES AND METHODS BASED THEREON

<130> 24745-1613PC

<140>

<141> Herewith

<150> 60/275,592

<151> 2001-03-13

<160> 22

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 3147

<212> DNA

<213> Homo Sapien

<220>

<221> CDS

<222> (23)...(2589)

<223> Nucleotide sequence encoding MTSP1

<300>

<301> O'Brien, T.J. and Tanimoto, H.

<308> GenBank #AR081724

<309> 2000-08-31

<310> 5,972,616

<311> 1998-02-20

<312> 1999-10-26

<400> 1

tcaagagcgg cctcggggta cc atg ggg agc gat cgg gcc cgc aag ggc gga	52
Met Gly Ser Asp Arg Ala Arg Lys Gly Gly	
1 5 10	
ggg ggc ccg aag gac ttc ggc gcg gga ctc aag tac aac tcc cgg cac	100
Gly Gly Pro Lys Asp Phe Gly Ala Gly Leu Lys Tyr Asn Ser Arg His	
15 20 25	
gag aaa gtg aat ggc ttg gag gaa ggc gtg gag ttc ctg cca gtc aac	148
Glu Lys Val Asn Gly Leu Glu Glu Gly Val Glu Phe Leu Pro Val Asn	
30 35 40	
aac gtc aag aag gtg gaa aag cat ggc ccg ggg cgc tgg gtg gtg ctg	196
Asn Val Lys Lys Val Glu Lys His Gly Pro Gly Arg Trp Val Val Leu	
45 50 55	
gca gcc gtg ctg atc ggc ctc ctc ttg gtc ttg ctg ggg atc ggc ttc	244
Ala Ala Val Leu Ile Gly Leu Leu Val Leu Leu Gly Ile Gly Phe	
60 65 70	

-2-

ctg	gtg	tgg	cat	ttg	cag	tac	cgg	gac	gtg	cgt	gtc	cag	aag	gtc	ttc	292
Leu	Val	Trp	His	Leu	Gln	Tyr	Arg	Asp	Val	Arg	Val	Gln	Lys	Val	Phe	
75					80					85					90	
aat	ggc	tac	atg	agg	atc	aca	aat	gag	aat	ttt	gtg	gat	gcc	tac	gag	340
Asn	Gly	Tyr	Met	Arg	Ile	Thr	Asn	Glu	Asn	Phe	Val	Asp	Ala	Tyr	Glu	
				95				100						105		
aac	tcc	aac	tcc	act	gag	ttt	gta	agc	ctg	gcc	agc	aag	gtg	aag	gac	388
Asn	Ser	Asn	Ser	Thr	Glu	Phe	Val	Ser	Leu	Ala	Ser	Lys	Val	Lys	Asp	
			110					115					120			
gcg	ctg	aag	ctg	ctg	tac	agc	gga	gtc	cca	ttc	ctg	ggc	ccc	tac	cac	436
Ala	Leu	Lys	Leu	Leu	Tyr	Ser	Gly	Val	Pro	Phe	Leu	Gly	Pro	Tyr	His	
		125					130					135				
aag	gag	tcg	gct	gtg	acg	gcc	ttc	agc	gag	ggc	agc	gtc	atc	gcc	tac	484
Lys	Glu	Ser	Ala	Val	Thr	Ala	Phe	Ser	Glu	Gly	Ser	Val	Ile	Ala	Tyr	
	140					145					150					
tac	tgg	tct	gag	ttc	agc	atc	ccg	cag	cac	ctg	gtg	gag	gag	gcc	gag	532
Tyr	Trp	Ser	Glu	Phe	Ser	Ile	Pro	Gln	His	Leu	Val	Glu	Glu	Ala	Glu	
155					160					165					170	
cgc	gtc	atg	gcc	gag	gag	cgc	gta	gtc	atg	ctg	ccc	ccg	cgg	gcg	cgc	580
Arg	Val	Met	Ala	Glu	Glu	Arg	Val	Val	Met	Leu	Pro	Pro	Arg	Ala	Arg	
				175					180					185		
tcc	ctg	aag	tcc	ttt	gtg	gtc	acc	tca	gtg	gtg	gct	ttc	ccc	acg	gac	628
Ser	Leu	Lys	Ser	Phe	Val	Val	Thr	Ser	Val	Val	Ala	Phe	Pro	Thr	Asp	
			190					195					200			
tcc	aaa	aca	gta	cag	agg	acc	cag	gac	aac	agc	tgc	agc	ttt	ggc	ctg	676
Ser	Lys	Thr	Val	Gln	Arg	Thr	Gln	Asp	Asn	Ser	Cys	Ser	Phe	Gly	Leu	
		205					210					215				
cac	gcc	cgc	ggt	gtg	gag	ctg	atg	cgc	ttc	acc	acg	ccc	ggc	ttc	cct	724
His	Ala	Arg	Gly	Val	Glu	Leu	Met	Arg	Phe	Thr	Thr	Pro	Gly	Phe	Pro	
	220					225					230					
gac	agc	ccc	tac	ccc	gct	cat	gcc	cgc	tgc	cag	tgg	gcc	ctg	cgg	ggg	772
Asp	Ser	Pro	Tyr	Pro	Ala	His	Ala	Arg	Cys	Gln	Trp	Ala	Leu	Arg	Gly	
235					240				245						250	
gac	gcc	gac	tca	gtg	ctg	agc	ctc	acc	ttc	cgc	agc	ttt	gac	ctt	gcg	820
Asp	Ala	Asp	Ser	Val	Leu	Ser	Leu	Thr	Phe	Arg	Ser	Phe	Asp	Leu	Ala	
				255				260						265		
tcc	tgc	gac	gag	cgc	ggc	agc	gac	ctg	gtg	acg	gtg	tac	aac	acc	ctg	868
Ser	Cys	Asp	Glu	Arg	Gly	Ser	Asp	Leu	Val	Thr	Val	Tyr	Asn	Thr	Leu	
			270					275					280			
agc	ccc	atg	gag	ccc	cac	gcc	ctg	gtg	cag	ttg	tgt	ggc	acc	tac	cct	916
Ser	Pro	Met	Glu	Pro	His	Ala	Leu	Val	Gln	Leu	Cys	Gly	Thr	Tyr	Pro	
		285					290					295				
ccc	tcc	tac	aac	ctg	acc	ttc	cac	tcc	tcc	cag	aac	gtc	ctg	ctc	atc	964
Pro	Ser	Tyr	Asn	Leu	Thr	Phe	His	Ser	Ser	Gln	Asn	Val	Leu	Leu	Ile	
	300					305					310					

-3-

aca	ctg	ata	acc	aac	act	gag	cgg	cgg	cat	ccc	ggc	ttt	gag	gcc	acc	1012
Thr	Leu	Ile	Thr	Asn	Thr	Glu	Arg	Arg	His	Pro	Gly	Phe	Glu	Ala	Thr	
315					320					325					330	
ttc	ttc	cag	ctg	cct	agg	atg	agc	agc	tgt	gga	ggc	cgc	tta	cgt	aaa	1060
Phe	Phe	Gln	Leu	Pro	Arg	Met	Ser	Ser	Cys	Gly	Gly	Arg	Leu	Arg	Lys	
				335					340					345		
gcc	cag	ggg	aca	ttc	aac	agc	ccc	tac	tac	cca	ggc	cac	tac	cca	ccc	1108
Ala	Gln	Gly	Thr	Phe	Asn	Ser	Pro	Tyr	Tyr	Pro	Gly	His	Tyr	Pro	Pro	
			350					355					360			
aac	att	gac	tgc	aca	tgg	aac	att	gag	gtg	ccc	aac	aac	cag	cat	gtg	1156
Asn	Ile	Asp	Cys	Thr	Trp	Asn	Ile	Glu	Val	Pro	Asn	Asn	Gln	His	Val	
		365					370					375				
aag	gtg	agc	ttc	aaa	ttc	ttc	tac	ctg	ctg	gag	ccc	ggc	gtg	cct	gcg	1204
Lys	Val	Ser	Phe	Lys	Phe	Phe	Tyr	Leu	Leu	Glu	Pro	Gly	Val	Pro	Ala	
	380					385					390					
ggc	acc	tgc	ccc	aag	gac	tac	gtg	gag	atc	aat	ggg	gag	aaa	tac	tgc	1252
Gly	Thr	Cys	Pro	Lys	Asp	Tyr	Val	Glu	Ile	Asn	Gly	Glu	Lys	Tyr	Cys	
395					400					405					410	
gga	gag	agg	tcc	cag	ttc	gtc	gtc	acc	agc	aac	agc	aac	aag	atc	aca	1300
Gly	Glu	Arg	Ser	Gln	Phe	Val	Val	Thr	Ser	Asn	Ser	Asn	Lys	Ile	Thr	
				415				420						425		
gtt	cgc	ttc	cac	tca	gat	cag	tcc	tac	acc	gac	acc	ggc	ttc	tta	gct	1348
Val	Arg	Phe	His	Ser	Asp	Gln	Ser	Tyr	Thr	Asp	Thr	Gly	Phe	Leu	Ala	
			430					435					440			
gaa	tac	ctc	tcc	tac	gac	tcc	agt	gac	cca	tgc	ccg	ggg	cag	ttc	acg	1396
Glu	Tyr	Leu	Ser	Tyr	Asp	Ser	Ser	Asp	Pro	Cys	Pro	Gly	Gln	Phe	Thr	
		445					450					455				
tgc	cgc	acg	ggg	cgg	tgt	atc	cgg	aag	gag	ctg	cgc	tgt	gat	ggc	tgg	1444
Cys	Arg	Thr	Gly	Arg	Cys	Ile	Arg	Lys	Glu	Leu	Arg	Cys	Asp	Gly	Trp	
	460					465					470					
gcc	gac	tgc	acc	gac	cac	agc	gat	gag	ctc	aac	tgc	agt	tgc	gac	gcc	1492
Ala	Asp	Cys	Thr	Asp	His	Ser	Asp	Glu	Leu	Asn	Cys	Ser	Cys	Asp	Ala	
475					480					485					490	
ggc	cac	cag	ttc	acg	tgc	aag	aac	aag	ttc	tgc	aag	ccc	ctc	ttc	tgg	1540
Gly	His	Gln	Phe	Thr	Cys	Lys	Asn	Lys	Phe	Cys	Lys	Pro	Leu	Phe	Trp	
				495				500						505		
gtc	tgc	gac	agt	gtg	aac	gac	tgc	gga	gac	aac	agc	gac	gag	cag	ggg	1588
Val	Cys	Asp	Ser	Val	Asn	Asp	Cys	Gly	Asp	Asn	Ser	Asp	Glu	Gln	Gly	
			510					515					520			
tgc	agt	tgt	ccg	gcc	cag	acc	ttc	agg	tgt	tcc	aat	ggg	aag	tgc	ctc	1636
Cys	Ser	Cys	Pro	Ala	Gln	Thr	Phe	Arg	Cys	Ser	Asn	Gly	Lys	Cys	Leu	
		525					530					535				
tcg	aaa	agc	cag	cag	tgc	aat	ggg	aag	gac	gac	tgt	ggg	gac	ggg	tcc	1684
Ser	Lys	Ser	Gln	Gln	Cys	Asn	Gly	Lys	Asp	Asp	Cys	Gly	Asp	Gly	Ser	

-4-

540	545	550	
gac gag gcc tcc tgc ccc aag gtg aac gtc gtc act tgt acc aaa cac Asp Glu Ala Ser Cys Pro Lys Val Asn Val Val Thr Cys Thr Lys His 555 560 565 570			1732
acc tac cgc tgc ctc aat ggg ctc tgc ttg agc aag ggc aac cct gag Thr Tyr Arg Cys Leu Asn Gly Leu Cys Leu Ser Lys Gly Asn Pro Glu 575 580 585			1780
tgt gac ggg aag gag gac tgt agc gac ggc tca gat gag aag gac tgc Cys Asp Gly Lys Glu Asp Cys Ser Asp Gly Ser Asp Glu Lys Asp Cys 590 595 600			1828
gac tgt ggg ctg cgg tca ttc acg aga cag gct cgt gtt gtt ggg ggc Asp Cys Gly Leu Arg Ser Phe Thr Arg Gln Ala Arg Val Val Gly Gly 605 610 615			1876
acg gat gcg gat gag ggc gag tgg ccc tgg cag gta agc ctg cat gct Thr Asp Ala Asp Glu Gly Glu Trp Pro Trp Gln Val Ser Leu His Ala 620 625 630			1924
ctg ggc cag ggc cac atc tgc ggt gct tcc ctc atc tct ccc aac tgg Leu Gly Gln Gly His Ile Cys Gly Ala Ser Leu Ile Ser Pro Asn Trp 635 640 645 650			1972
ctg gtc tct gcc gca cac tgc tac atc gat gac aga gga ttc agg tac Leu Val Ser Ala Ala His Cys Tyr Ile Asp Asp Arg Gly Phe Arg Tyr 655 660 665			2020
tca gac ccc acg cag tgg acg gcc ttc ctg ggc ttg cac gac cag agc Ser Asp Pro Thr Gln Trp Thr Ala Phe Leu Gly Leu His Asp Gln Ser 670 675 680			2068
cag cgc agc gcc cct ggg gtg cag gag cgc agg ctc aag cgc atc atc Gln Arg Ser Ala Pro Gly Val Gln Glu Arg Arg Leu Lys Arg Ile Ile 685 690 695			2116
tcc cac ccc ttc ttc aat gac ttc acc ttc gac tat gac atc gcg ctg Ser His Pro Phe Phe Asn Asp Phe Thr Phe Asp Tyr Asp Ile Ala Leu 700 705 710			2164
ctg gag ctg gag aaa ccg gca gag tac agc tcc atg gtg cgg ccc atc Leu Glu Leu Glu Lys Pro Ala Glu Tyr Ser Ser Met Val Arg Pro Ile 715 720 725 730			2212
tgc ctg ccg gac gcc tcc cat gtc ttc cct gcc ggc aag gcc atc tgg Cys Leu Pro Asp Ala Ser His Val Phe Pro Ala Gly Lys Ala Ile Trp 735 740 745			2260
gtc acg ggc tgg gga cac acc cag tat gga ggc act ggc gcg ctg atc Val Thr Gly Trp Gly His Thr Gln Tyr Gly Gly Thr Gly Ala Leu Ile 750 755 760			2308
ctg caa aag ggt gag atc cgc gtc atc aac cag acc acc tgc gag aac Leu Gln Lys Gly Glu Ile Arg Val Ile Asn Gln Thr Thr Cys Glu Asn 765 770 775			2356
ctc ctg ccg cag cag atc acg ccg cgc atg atg tgc gtg ggc ttc ctc			2404

-5-

Leu	Leu	Pro	Gln	Gln	Ile	Thr	Pro	Arg	Met	Met	Cys	Val	Gly	Phe	Leu		
780						785					790						
agc	ggc	ggc	gtg	gac	tcc	tgc	cag	ggg	gat	tcc	ggg	gga	ccc	ctg	tcc	2452	
Ser	Gly	Gly	Val	Asp	Ser	Cys	Gln	Gly	Asp	Ser	Gly	Gly	Pro	Leu	Ser	810	
795					800					805							
agc	gtg	gag	gag	gat	ggg	cgg	atc	ttc	cag	gcc	ggg	gtg	gtg	agc	tgg	2500	
Ser	Val	Glu	Ala	Asp	Gly	Arg	Ile	Phe	Gln	Ala	Gly	Val	Val	Ser	Trp	825	
				815					820								
gga	gac	ggc	tgc	gct	cag	agg	aac	aag	cca	ggc	gtg	tac	aca	agg	ctc	2548	
Gly	Asp	Gly	Cys	Ala	Gln	Arg	Asn	Lys	Pro	Gly	Val	Tyr	Thr	Arg	Leu	840	
			830					835									
cct	ctg	ttt	cgg	gac	tgg	atc	aaa	gag	aac	act	ggg	gta	ta	ggggccgggg	2599		
Pro	Leu	Phe	Arg	Asp	Trp	Ile	Lys	Glu	Asn	Thr	Gly	Val					
		845					850					855					
ccacccaaat	gtgtacacct	gcgggggccac	ccatcggtcca	ccccagtggtg	cacgcctgca	2659											
ggctggagac	tggaccgctg	actgcaccag	cgccccccaga	acatacactg	tgaactcaat	2719											
ctccagggct	ccaaatctgc	ctagaaaacc	tctcgcttcc	tcagcctcca	aagtggagct	2779											
gggaggtaga	aggggaggac	actggtggtt	ctactgaccc	aactgggggc	aaaggtttga	2839											
agacacagcc	tcccccgcca	gcccccaagct	gggcccaggc	gcgtttgtgt	atatctgcct	2899											
cccctgtctg	taaggagcag	cgggaacgga	gcttcggagc	ctcctcagtg	aaggtggtgg	2959											
ggctgccgga	tctgggctgt	ggggcccttg	ggccacgctc	ttgaggaagc	ccaggctcgg	3019											
aggaccctgg	aaaacagacg	ggtctgagac	tgaaattgtt	ttaccagctc	ccagggtgga	3079											
cttcagtggtg	tgtatttgtg	taaatgggta	aaacaattta	tttcttttta	aaaaaaaaaa	3139											
aaaaaaaaaa						3147											

&lt;210&gt; 2

&lt;211&gt; 855

&lt;212&gt; PRT

&lt;213&gt; Homo Sapien

&lt;400&gt; 2

Met	Gly	Ser	Asp	Arg	Ala	Arg	Lys	Gly	Gly	Gly	Gly	Pro	Lys	Asp	Phe		
1				5				10						15			
Gly	Ala	Gly	Leu	Lys	Tyr	Asn	Ser	Arg	His	Glu	Lys	Val	Asn	Gly	Leu		
			20					25					30				
Glu	Glu	Gly	Val	Glu	Phe	Leu	Pro	Val	Asn	Asn	Val	Lys	Lys	Val	Glu		
			35				40						45				
Lys	His	Gly	Pro	Gly	Arg	Trp	Val	Val	Leu	Ala	Ala	Val	Leu	Ile	Gly		
			50			55				60							
Leu	Leu	Leu	Val	Leu	Leu	Gly	Ile	Gly	Phe	Leu	Val	Trp	His	Leu	Gln		
65				70					75						80		
Tyr	Arg	Asp	Val	Arg	Val	Gln	Lys	Val	Phe	Asn	Gly	Tyr	Met	Arg	Ile		
				85					90					95			
Thr	Asn	Glu	Asn	Phe	Val	Asp	Ala	Tyr	Glu	Asn	Ser	Asn	Ser	Thr	Glu		
			100					105						110			
Phe	Val	Ser	Leu	Ala	Ser	Lys	Val	Lys	Asp	Ala	Leu	Lys	Leu	Leu	Tyr		
			115				120						125				
Ser	Gly	Val	Pro	Phe	Leu	Gly	Pro	Tyr	His	Lys	Glu	Ser	Ala	Val	Thr		
			130			135					140						
Ala	Phe	Ser	Glu	Gly	Ser	Val	Ile	Ala	Tyr	Tyr	Trp	Ser	Glu	Phe	Ser		
145				150					155					160			
Ile	Pro	Gln	His	Leu	Val	Glu	Glu	Ala	Glu	Arg	Val	Met	Ala	Glu	Glu		
			165					170						175			
Arg	Val	Val	Met	Leu	Pro	Pro	Arg	Ala	Arg	Ser	Leu	Lys	Ser	Phe	Val		



-6-

Val	Thr	Ser	180	Val	Val	Ala	Phe	Pro	185	Thr	Asp	Ser	Lys	Thr	190	Val	Gln	Arg
Thr	Gln	Asp	195	Asn	Ser	Cys	Ser	Phe	200	Gly	Leu	His	Ala	Arg	205	Gly	Val	Glu
Leu	Met	Arg	210	Phe	Thr	Thr	Pro	Gly	215	Phe	Pro	Asp	Ser	Pro	220	Tyr	Pro	Ala
225	Ala	Arg	230	Cys	Gln	Trp	Ala	Leu	235	Arg	Gly	Asp	Ala	Asp	240	Ser	Val	Leu
His	Ala	Arg	245	Arg	Ser	Phe	Asp	Leu	250	Ala	Ser	Cys	Asp	Glu	255	Arg	Gly	
Ser	Leu	Thr	260	Val	Thr	Val	Tyr	Asn	265	Thr	Leu	Ser	Pro	Met	270	Glu	Pro	His
Ser	Asp	Leu	275	Gln	Leu	Cys	Gly	Thr	280	Tyr	Pro	Pro	Ser	Tyr	285	Asn	Leu	Thr
Ala	Leu	Val	290	Ser	Ser	Gln	Asn	Val	295	Leu	Leu	Ile	Thr	Leu	300	Ile	Thr	Asn
305	His	Ser	310	His	Pro	Gly	Phe	Glu	315	Ala	Thr	Phe	Phe	Gln	320	Leu	Pro	Arg
Glu	Arg	Arg	325	Cys	Gly	Gly	Arg	Leu	330	Arg	Lys	Ala	Gln	Gly	335	Thr	Phe	Asn
Met	Ser	Ser	340	Pro	Gly	His	Tyr	Pro	345	Pro	Pro	Asn	Ile	Asp	350	Cys	Thr	Trp
Ser	Pro	Tyr	355	Val	Pro	Asn	Asn	Gln	360	His	Val	Lys	Val	Ser	365	Phe	Lys	Phe
Asn	Ile	Glu	370	Leu	Leu	Glu	Pro	Gly	375	Val	Pro	Ala	Gly	Thr	380	Cys	Pro	Lys
385	Tyr	Val	390	Ile	Asn	Gly	Glu	Lys	395	Tyr	Cys	Gly	Glu	Arg	400	Ser	Gln	Phe
Tyr	Val	Glu	405	Asn	Ser	Asn	Lys	Ile	410	Thr	Val	Arg	Phe	His	415	Ser	Asp	
Val	Val	Thr	420	Thr	Asp	Thr	Gly	Phe	425	Leu	Ala	Glu	Tyr	Leu	430	Ser	Tyr	Asp
Gln	Ser	Tyr	435	Pro	Cys	Pro	Gly	Gln	440	Phe	Thr	Cys	Arg	Thr	445	Gly	Arg	Cys
Ser	Ser	Asp	450	Glu	Leu	Arg	Cys	Asp	455	Gly	Trp	Ala	Asp	Cys	460	Thr	Asp	His
Ile	Arg	Lys	465	Leu	Asn	Cys	Ser	Cys	470	Ala	Gly	His	Gln	Phe	475	Thr	Cys	
Ser	Asp	Glu	485	Cys	Lys	Pro	Leu	Phe	490	Trp	Val	Cys	Asp	Ser	495	Val	Asn	
Lys	Asn	Lys	500	Asn	Ser	Asp	Glu	Gln	505	Gly	Cys	Ser	Cys	Pro	510	Ala	Gln	
Asp	Cys	Gly	515	Ser	Asn	Gly	Lys	Cys	520	Leu	Ser	Lys	Ser	Gln	525	Gln	Cys	
Thr	Phe	Arg	530	Asp	Asp	Cys	Gly	Asp	535	Gly	Ser	Asp	Glu	Ala	540	Ser	Cys	Pro
Asn	Gly	Lys	545	Val	Val	Thr	Cys	Thr	550	Lys	His	Thr	Tyr	Arg	555	Cys	Leu	Asn
Lys	Val	Asn	565	Ser	Lys	Gly	Asn	Pro	570	Glu	Cys	Asp	Gly	Lys	575	Glu	Asp	
Gly	Leu	Cys	580	Ser	Asp	Glu	Lys	Asp	585	Cys	Asp	Cys	Gly	Leu	590	Arg	Ser	
Cys	Ser	Asp	595	Gln	Ala	Arg	Val	Val	600	Gly	Gly	Thr	Asp	Ala	605	Glu	Gly	
Phe	Thr	Arg	610	Trp	Gln	Val	Ser	Leu	615	His	Ala	Leu	Gly	Gln	620	Gly	His	Ile
Glu	Trp	Pro	625	Leu	Ile	Ser	Pro	Asn	630	Trp	Leu	Val	Ser	Ala	635	Ala	His	
Cys	Gly	Ala	645	Ser	Leu	Ile	Ser	Pro	650	Trp	Leu	Val	Ser	Ala	655	Ala	His	

-7-

Cys	Tyr	Ile	Asp	Asp	Arg	Gly	Phe	Arg	Tyr	Ser	Asp	Pro	Thr	Gln	Trp
			660					665					670		
Thr	Ala	Phe	Leu	Gly	Leu	His	Asp	Gln	Ser	Gln	Arg	Ser	Ala	Pro	Gly
		675					680					685			
Val	Gln	Glu	Arg	Arg	Leu	Lys	Arg	Ile	Ile	Ser	His	Pro	Phe	Phe	Asn
	690					695					700				
Asp	Phe	Thr	Phe	Asp	Tyr	Asp	Ile	Ala	Leu	Leu	Glu	Leu	Glu	Lys	Pro
705				710					715						720
Ala	Glu	Tyr	Ser	Ser	Met	Val	Arg	Pro	Ile	Cys	Leu	Pro	Asp	Ala	Ser
			725					730						735	
His	Val	Phe	Pro	Ala	Gly	Lys	Ala	Ile	Trp	Val	Thr	Gly	Trp	Gly	His
		740					745						750		
Thr	Gln	Tyr	Gly	Gly	Thr	Gly	Ala	Leu	Ile	Leu	Gln	Lys	Gly	Glu	Ile
	755					760					765				
Arg	Val	Ile	Asn	Gln	Thr	Thr	Cys	Glu	Asn	Leu	Leu	Pro	Gln	Gln	Ile
	770				775						780				
Thr	Pro	Arg	Met	Met	Cys	Val	Gly	Phe	Leu	Ser	Gly	Gly	Val	Asp	Ser
785				790					795						800
Cys	Gln	Gly	Asp	Ser	Gly	Gly	Pro	Leu	Ser	Ser	Val	Glu	Ala	Asp	Gly
			805					810						815	
Arg	Ile	Phe	Gln	Ala	Gly	Val	Val	Ser	Trp	Gly	Asp	Gly	Cys	Ala	Gln
		820					825						830		
Arg	Asn	Lys	Pro	Gly	Val	Tyr	Thr	Arg	Leu	Pro	Leu	Phe	Arg	Asp	Trp
	835					840						845			
Ile	Lys	Glu	Asn	Thr	Gly	Val									
	850					855									

&lt;210&gt; 3

&lt;211&gt; 3147

&lt;212&gt; DNA

&lt;213&gt; Homo Sapien

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1865)...(2590)

&lt;223&gt; Nucleic acid sequence of protease domain of MTSP1

&lt;400&gt; 3

tcaagagcgg	cctcggggta	ccatggggag	cgatcggggc	cgcaagggcg	gagggggccc	60
gaaggacttc	ggcgcgggac	tcaagtacaa	ctcccggcac	gagaaagtga	atggcttgga	120
ggaaggcgtg	gagttcctgc	cagtcaacaa	cgtcaagaag	gtggaaaagc	atggcccggg	180
gcgctgggtg	gtgctggcag	ccgtgctgat	cggcctcctc	ttggtcttgc	tggggatcgg	240
cttcctgggtg	tggcatttgc	agtaccggga	cgtgcgtgtc	cagaaggtct	tcaatggcta	300
catgaggatc	acaaatgaga	attttgtgga	tgcctacgag	aactccaact	ccactgagtt	360
tgtaaagcctg	gccagcaagg	tgaaggacgc	gctgaagctg	ctgtacagcg	gagtcccatt	420
cctggggccc	taccacaagg	agtcggctgt	gacggccttc	agcgagggca	gcgtcatcgc	480
ctactactgg	tctgagttca	gcatcccgca	gcacctgggtg	gaggaggccg	agcgcgatcat	540
ggccgaggag	cgcgtagtca	tgctgccccc	gcgggcgcgc	tccctgaagt	cctttgtggt	600
cacctcagtg	gtggctttcc	ccacggactc	caaaacagta	cagaggaccc	aggacaacag	660
ctgcagcttt	ggcctgcacg	cccgcgggtg	ggagctgatg	cgcttcacca	cgcccggctt	720
ccctgacagc	ccctaccccg	ctcatgcccc	ctgccagtgg	gccctgcggg	gggacgccga	780
ctcagtgctg	agcctcacct	tccgcagctt	tgaccttgcg	tcctgcgacg	agcgcggcag	840
cgacctgggtg	acggtgtaca	acaccttgag	ccccatggag	ccccacgccc	tggtgcagtt	900
gtgtggcacc	tacctcctc	cctacaacct	gaccttccac	tcctcccaga	acgtcctgct	960
catcacactg	ataaccaaca	ctgagcggcg	gcatcccggc	tttgaggcca	ccttcttcca	1020
gctgcctagg	atgagcagct	gtggaggccg	cttacgtaaa	gcccagggga	cattcaacag	1080
cccctactac	ccaggccact	accaccccaa	cattgactgc	acatggaaca	ttgaggtgcc	1140
caacaaccag	catgtgaagg	tgagcttcaa	attcttctac	ctgctggagc	ccggcgtgcc	1200
tgcgggcacc	tgccccaagg	actacgtgga	gatcaatggg	gagaaatact	gcggagagag	1260

-8-

gtcccagttc	gtcgtcacca	gcaacagcaa	caagatcaca	gttcgcttcc	actcagatca	1320
gtcctacacc	gacaccggt	tcttagctga	atacctctcc	tacgactcca	gtgacccatg	1380
cccggggcag	ttcacgtgcc	gcacggggcg	gtgtatccgg	aaggagctgc	gctgtgatgg	1440
ctgggccgac	tgcaccgacc	acagcgatga	gctcaactgc	agttgcgacg	ccggccacca	1500
gttcacgtgc	aagaacaagt	tctgcaagcc	cctcttctgg	gtctgcgaca	gtgtgaacga	1560
ctgcggagac	aacagcgacg	agcaggggtg	cagttgtccg	gcccagacct	tcaggtgttc	1620
caatgggaag	tgcctctcga	aaagccagca	gtgcaatggg	aaggacgact	gtggggacgg	1680
gtccgacgag	gcctcctgcc	ccaaggtgaa	cgctcgtcact	tgtaccaaac	acacctaccg	1740
ctgcctcaat	gggctctgct	tgagcaaggg	caaccctgag	tgtgacggga	aggaggactg	1800
tagcgacggc	tcagatgaga	aggactgcga	ctgtgggctg	cggtcattca	cgagacaggc	1860
tcgt gtt gtt ggg ggc acg gat gcg gat gag ggc gag tgg ccc tgg cag	Val Val Gly Gly Thr Asp Ala Asp Glu Gly Glu Trp Pro Trp Gln	1909				
1	5	10	15			
gta agc ctg cat gct ctg ggc cag ggc cac atc tgc ggt gct tcc ctc	Val Ser Leu His Ala Leu Gly Gln Gly His Ile Cys Gly Ala Ser Leu	1957				
	20	25	30			
atc tct ccc aac tgg ctg gtc tct gcc gca cac tgc tac atc gat gac	Ile Ser Pro Asn Trp Leu Val Ser Ala Ala His Cys Tyr Ile Asp Asp	2005				
	35	40	45			
aga gga ttc agg tac tca gac ccc acg cag tgg acg gcc ttc ctg ggc	Arg Gly Phe Arg Tyr Ser Asp Pro Thr Gln Trp Thr Ala Phe Leu Gly	2053				
	50	55	60			
ttg cac gac cag agc cag cgc agc gcc cct ggg gtg cag gag cgc agg	Leu His Asp Gln Ser Gln Arg Ser Ala Pro Gly Val Gln Glu Arg Arg	2101				
	65	70	75			
ctc aag cgc atc atc tcc cac ccc ttc ttc aat gac ttc acc ttc gac	Leu Lys Arg Ile Ile Ser His Pro Phe Phe Asn Asp Phe Thr Phe Asp	2149				
	80	85	90			
tat gac atc gcg ctg ctg gag ctg gag aaa ccg gca gag tac agc tcc	Tyr Asp Ile Ala Leu Leu Glu Leu Glu Lys Pro Ala Glu Tyr Ser Ser	2197				
	100	105	110			
atg gtg cgg ccc atc tgc ctg ccg gac gcc tcc cat gtc ttc cct gcc	Met Val Arg Pro Ile Cys Leu Pro Asp Ala Ser His Val Phe Pro Ala	2245				
	115	120	125			
ggc aag gcc atc tgg gtc acg ggc tgg gga cac acc cag tat gga ggc	Gly Lys Ala Ile Trp Val Thr Gly Trp Gly His Thr Gln Tyr Gly Gly	2293				
	130	135	140			
act ggc gcg ctg atc ctg caa aag ggt gag atc cgc gtc atc aac cag	Thr Gly Ala Leu Ile Leu Gln Lys Gly Glu Ile Arg Val Ile Asn Gln	2341				
	145	150	155			
acc acc tgc gag aac ctc ctg ccg cag cag atc acg ccg cgc atg atg	Thr Thr Cys Glu Asn Leu Leu Pro Gln Gln Ile Thr Pro Arg Met Met	2389				
	160	165	170			
tgc gtg ggc ttc ctc agc ggc ggc gtg gac tcc tgc cag ggt gat tcc	Cys Val Gly Phe Leu Ser Gly Gly Val Asp Ser Cys Gln Gly Asp Ser	2437				
	180	185	190			
ggg gga ccc ctg tcc agc gtg gag gcg gat ggg cgg atc ttc cag gcc		2485				

-9-

Gly Gly Pro Leu Ser Ser Val Glu Ala Asp Gly Arg Ile Phe Gln Ala  
 195 200 205  
 ggt gtg gtg agc tgg gga gac ggc tgc gct cag agg aac aag cca ggc 2533  
 Gly Val Val Ser Trp Gly Asp Gly Cys Ala Gln Arg Asn Lys Pro Gly  
 210 215 220  
 gtg tac aca agg ctc cct ctg ttt cgg gac tgg atc aaa gag aac act 2581  
 Val Tyr Thr Arg Leu Pro Leu Phe Arg Asp Trp Ile Lys Glu Asn Thr  
 225 230 235  
 ggg gta tag gggccggggc cacccaaattg tgtacacctg cggggccacc 2630  
 Gly Val \*  
 240  
 catcgtccac cccagtgtgc acgectgcag gctggagact ggaccgctga ctgcaccagc 2690  
 gccccagaa catacactgt gaactcaatc tccagggtc caaatctgcc tagaaaacct 2750  
 ctcgcttct cagcctccaa agtggagctg ggaggtagaa ggggaggaca ctggtggttc 2810  
 tactgaccca actggggggca aagggtttgaa gacacagcct cccccgccag cccaagctg 2870  
 ggccgagggc cgtttgtgta tatctgcctc ccctgtctgt aaggagcagc gggaacggag 2930  
 cttcggagcc tcctcagtga aggtggtggg gctgccggat ctgggctgtg gggcccttgg 2990  
 gccacgtctt tgaggaagcc caggctcggg ggaccctgga aaacagacgg gtctgagact 3050  
 gaaattgttt taccagctcc cagggtggac ttcagtgtgt gtatttgtgt aaatgggtaa 3110  
 aacaatttat ttctttttaa aaaaaaaaaa aaaaaaa 3147  
 <210> 4  
 <211> 241  
 <212> PRT  
 <213> Homo Sapien  
 <400> 4  
 Val Val Gly Gly Thr Asp Ala Asp Glu Gly Glu Trp Pro Trp Gln Val  
 1 5 10 15  
 Ser Leu His Ala Leu Gly Gln Gly His Ile Cys Gly Ala Ser Leu Ile  
 20 25 30  
 Ser Pro Asn Trp Leu Val Ser Ala Ala His Cys Tyr Ile Asp Asp Arg  
 35 40 45  
 Gly Phe Arg Tyr Ser Asp Pro Thr Gln Trp Thr Ala Phe Leu Gly Leu  
 50 55 60  
 His Asp Gln Ser Gln Arg Ser Ala Pro Gly Val Gln Glu Arg Arg Leu  
 65 70 75 80  
 Lys Arg Ile Ile Ser His Pro Phe Phe Asn Asp Phe Thr Phe Asp Tyr  
 85 90 95  
 Asp Ile Ala Leu Leu Glu Leu Glu Lys Pro Ala Glu Tyr Ser Ser Met  
 100 105 110  
 Val Arg Pro Ile Cys Leu Pro Asp Ala Ser His Val Phe Pro Ala Gly  
 115 120 125  
 Lys Ala Ile Trp Val Thr Gly Trp Gly His Thr Gln Tyr Gly Gly Thr  
 130 135 140  
 Gly Ala Leu Ile Leu Gln Lys Gly Glu Ile Arg Val Ile Asn Gln Thr  
 145 150 155 160  
 Thr Cys Glu Asn Leu Leu Pro Gln Gln Ile Thr Pro Arg Met Met Cys  
 165 170 175  
 Val Gly Phe Leu Ser Gly Gly Val Asp Ser Cys Gln Gly Asp Ser Gly  
 180 185 190  
 Gly Pro Leu Ser Ser Val Glu Ala Asp Gly Arg Ile Phe Gln Ala Gly  
 195 200 205  
 Val Val Ser Trp Gly Asp Gly Cys Ala Gln Arg Asn Lys Pro Gly Val

-10-

210		215		220
Tyr	Thr	Arg	Leu	Pro
225			Leu	Phe
Val			230	Arg
				Asp
				Trp
				Ile
				235
				Lys
				Glu
				Asn
				Thr
				Gly
				240

<210> 5  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Primer

<400> 5  
 aatggccatg gcaggccagc ctcc

24

<210> 6  
 <211> 32  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Primer

<400> 6  
 gtcccaaact tactatacct acaatgtacc ag

32

<210> 7  
 <211> 32  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Primer

<400> 7  
 gtcccaaact tactatacct acaatgtacc ag

32

<210> 8  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Primer

<400> 8  
 aatggccatg gcaggccagc ctcc

24

<210> 9  
 <211> 31  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Primer

<400> 9

-11-

tgccattacc agcatcctct tctactcaaa g 31

<210> 10  
 <211> 33  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Primer

<400> 10  
 ccatgtgtat aactcacgga caatccacac tac 33

<210> 11  
 <211> 10  
 <212> PRT  
 <213> Homo sapien

<400> 11  
 Met Pro Leu Pro Ala Ser Ser Ser Thr Gln  
 1 5 10

<210> 12  
 <211> 42  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Primer

<400> 12  
 tctctcgaga aaagaattgt ccaaggaagg gaaacagcta tg 42

<210> 13  
 <211> 39  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Primer

<400> 13  
 atagcggccg cacactacat accagtcttt gaggcaatc 39

<210> 14  
 <211> 11  
 <212> PRT  
 <213> Homo sapien

<400> 14  
 Lys Arg Ile Val Gln Gly Arg Glu Thr Ala Met  
 1 5 10

<210> 15  
 <211> 2100  
 <212> DNA  
 <213> Homo sapien

<220>

-12-

```
<221> CDS
<222> (45)...(1361)
<223> MTSP7: full length cDNA
```

```
<220>
<221> CDS
<222> (45)...(1361)
<223> MTSP7: full length cDNA
```

<400> 15																
agatcagatg gcgactgaat agaagctgcc ccagtcctgg gttc atg atg tac aca																56
Met Met Tyr Thr																
1																
cct	ggt	gaa	ttt	tca	gaa	gct	gaa	ttc	tca	cga	gct	gaa	tat	caa	aga	104
Pro	Val	Glu	Phe	Ser	Glu	Ala	Glu	Phe	Ser	Arg	Ala	Glu	Tyr	Gln	Arg	
5					10					15					20	
aag	cag	caa	ttt	tgg	gac	tca	gta	cgg	cta	gct	ctt	ttc	aca	tta	gca	152
Lys	Gln	Gln	Phe	Trp	Asp	Ser	Val	Arg	Leu	Ala	Leu	Phe	Thr	Leu	Ala	
				25					30						35	
att	gta	gca	atc	ata	gga	att	gca	att	ggg	att	gtt	act	cat	ttt	gtt	200
Ile	Val	Ala	Ile	Ile	Gly	Ile	Ala	Ile	Gly	Ile	Val	Thr	His	Phe	Val	
			40					45					50			
gtt	gag	gat	gat	aag	tct	ttc	tat	tac	ctt	gcc	tct	ttt	aaa	gtc	aca	248
Val	Glu	Asp	Asp	Lys	Ser	Phe	Tyr	Tyr	Leu	Ala	Ser	Phe	Lys	Val	Thr	
		55					60					65				
aat	atc	aaa	tat	aaa	gaa	aat	tat	ggc	ata	aga	tct	tca	aga	gag	ttt	296
Asn	Ile	Lys	Tyr	Lys	Glu	Asn	Tyr	Gly	Ile	Arg	Ser	Ser	Arg	Glu	Phe	
	70					75					80					
ata	gaa	agg	agt	cat	cag	att	gaa	aga	atg	atg	tct	agg	ata	ttt	cga	344
Ile	Glu	Arg	Ser	His	Gln	Ile	Glu	Arg	Met	Met	Ser	Arg	Ile	Phe	Arg	
85					90					95					100	
cat	tct	tct	gta	ggc	ggg	cga	ttt	atc	aaa	tct	cat	gtt	atc	aaa	tta	392
His	Ser	Ser	Val	Gly	Gly	Arg	Phe	Ile	Lys	Ser	His	Val	Ile	Lys	Leu	
				105					110					115		
agt	cca	gat	gaa	caa	ggg	gtg	gat	att	ctt	ata	gtg	ctc	ata	ttt	cga	440
Ser	Pro	Asp	Glu	Gln	Gly	Val	Asp	Ile	Leu	Ile	Val	Leu	Ile	Phe	Arg	
			120					125					130			
tac	cca	tct	act	gat	agt	gct	gaa	caa	atc	aag	aaa	aaa	att	gaa	aag	488
Tyr	Pro	Ser	Thr	Asp	Ser	Ala	Glu	Gln	Ile	Lys	Lys	Lys	Ile	Glu	Lys	
		135					140					145				
gct	tta	tat	caa	agt	ttg	aag	acc	aaa	caa	ttg	tct	ttg	acc	ata	aac	536
Ala	Leu	Tyr	Gln	Ser	Leu	Lys	Thr	Lys	Gln	Leu	Ser	Leu	Thr	Ile	Asn	
	150					155					160					
aaa	cca	tca	ttt	aga	ctc	aca	cct	att	gac	agc	aaa	aag	atg	agg	aat	584
Lys	Pro	Ser	Phe	Arg	Leu	Thr	Pro	Ile	Asp	Ser	Lys	Lys	Met	Arg	Asn	
165					170					175					180	
ctt	ctc	aac	agt	cgc	tgt	gga	ata	agg	atg	aca	tct	tca	aac	atg	cca	632

-13-

Leu	Leu	Asn	Ser	Arg 185	Cys	Gly	Ile	Arg	Met 190	Thr	Ser	Ser	Asn	Met 195	Pro		
tta	cca	gca	tcc	tct	tct	act	caa	aga	att	gtc	caa	gga	agg	gaa	aca	680	
Leu	Pro	Ala	Ser 200	Ser	Ser	Thr	Gln	Arg 205	Ile	Val	Gln	Gly	Arg 210	Glu	Thr		
gct	atg	gaa	ggg	gaa	tgg	cca	tgg	cag	gcc	agc	ctc	cag	ctc	ata	ggg	728	
Ala	Met	Glu	Gly 215	Glu	Trp	Pro	Trp 220	Gln	Ala	Ser	Leu	Gln 225	Leu	Ile	Gly		
tca	ggc	cat	cag	tgt	gga	gcc	agc	ctc	atc	agt	aac	aca	tgg	ctg	ctc	776	
Ser	Gly 230	His	Gln	Cys	Gly	Ala 235	Ser	Leu	Ile	Ser	Asn 240	Thr	Trp	Leu	Leu		
aca	gca	gct	cac	tgc	ttt	tgg	aaa	aat	aaa	gac	cca	act	caa	tgg	att	824	
Thr	Ala	Ala	His	Cys	Phe 250	Trp	Lys	Asn	Lys	Asp 255	Pro	Thr	Gln	Trp	Ile 260		
gct	act	ttt	ggg	gca	act	ata	aca	cca	ccc	gca	gtg	aaa	cga	aat	gtg	872	
Ala	Thr	Phe	Gly 265	Ala	Thr	Ile	Thr	Pro	Pro 270	Ala	Val	Lys	Arg	Asn	Val 275		
agg	aaa	att	att	ctt	cat	gag	aat	tac	cat	aga	gaa	aca	aat	gaa	aat	920	
Arg	Lys	Ile	Ile 280	Leu	His	Glu	Asn 285	Tyr	His	Arg	Glu	Thr	Asn 290	Glu	Asn		
gac	att	gct	ttg	gtt	cag	ctc	tct	act	gga	gtt	gag	ttt	tca	aat	ata	968	
Asp	Ile	Ala 295	Leu	Val	Gln	Leu	Ser 300	Thr	Gly	Val	Glu	Phe 305	Ser	Asn	Ile		
gtc	cag	aga	gtt	tgc	ctc	cca	gac	tca	tct	ata	aag	ttg	cca	cct	aaa	1016	
Val	Gln 310	Arg	Val	Cys	Leu	Pro 315	Asp	Ser	Ser	Ile	Lys 320	Leu	Pro	Pro	Lys		
aca	agt	gtg	ttc	gtc	aca	gga	ttt	gga	tcc	att	gta	gat	gat	gga	cct	1064	
Thr	Ser	Val	Phe	Val	Thr 330	Gly	Phe	Gly	Ser	Ile 335	Val	Asp	Asp	Gly	Pro 340		
ata	caa	aat	aca	ctt	cgg	caa	gcc	aga	gtg	gaa	acc	ata	agc	act	gat	1112	
Ile	Gln	Asn	Thr 345	Leu	Arg	Gln	Ala	Arg 350	Val	Glu	Thr	Ile	Ser	Thr 355	Asp		
gtg	tgt	aac	aga	aag	gat	gtg	tat	gat	ggc	ctg	ata	act	cca	gga	atg	1160	
Val	Cys	Asn 360	Arg	Lys	Asp	Val	Tyr 365	Asp	Gly	Leu	Ile	Thr	Pro 370	Gly	Met		
tta	tgt	gct	gga	ttc	atg	gaa	gga	aaa	ata	gat	gca	tgt	aag	gga	gat	1208	
Leu	Cys	Ala 375	Gly	Phe	Met	Glu	Gly 380	Lys	Ile	Asp	Ala	Cys 385	Lys	Gly	Asp		
tct	ggg	gga	cct	ctg	gtt	tat	gat	aat	cat	gac	atc	tgg	tac	att	gta	1256	
Ser	Gly 390	Gly	Pro	Leu	Val	Tyr 395	Asp	Asn	His	Asp 400	Ile	Trp	Tyr	Ile	Val		
ggg	ata	gta	agt	tgg	gga	caa	tca	tgt	gca	ctt	ccc	aaa	aaa	cct	gga	1304	
Gly	Ile	Val	Ser	Trp	Gly 410	Gln	Ser	Cys	Ala	Leu 415	Pro	Lys	Lys	Pro	Gly 420		



-14-

gtc tac acc aga gta act aag tat cga gat tgg att gcc tca aag act 1352  
 Val Tyr Thr Arg Val Thr Lys Tyr Arg Asp Trp Ile Ala Ser Lys Thr  
                   425                                  430                                  435

ggg atg tag tgtggattgt ccatgagtta tacacatggc acacagagct 1401  
 Gly Met \*

gatactcctg cgtattttgt attgttttaa ttcattttact ttggattagt gcttttgcta 1461  
 gatgtcaaga agcccttcag acccagacaa atctaatact ctgaggtggc ctttacatac 1521  
 gtaggaccaa accctctcta ccatgagggg agaagacaca gcaaatgaca gacagcacct 1581  
 attccttact cacaagggaa actgcttgtg atacttccta ataagataaa taagtgggtt 1641  
 ccctcaattg aagacaggaa catcattttc cacaggatat gaagagctgc cagtaatgcc 1701  
 aaaatcctac ctcatataat acctggagca tgtgagattc ttctagttaa aaagaacagt 1761  
 cttccctgaa gactcagggc ttcaacattc tagaactgat aagtggacct tcagtgtgca 1821  
 agaatggaga agcatgggat ttgcattatg acttgaactg ggcttatata taataatata 1881  
 gagcactatc actaacctca acagttgaca ttttaaaagt ttttaaatgt atctgaactt 1941  
 gctgttaaca cagtgttata actcaagcac tagcttcagg aagcatgttg tgttgttaag 2001  
 aagcttttct gatttattct ttaacagcat cttgccatct atatgttagt agcagttggc 2061  
 ccagaaagga caaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2100

<210> 16  
 <211> 438  
 <212> PRT  
 <213> Homo sapien

<400> 16  
 Met Met Tyr Thr Pro Val Glu Phe Ser Glu Ala Glu Phe Ser Arg Ala  
   1                  5                  10                  15  
 Glu Tyr Gln Arg Lys Gln Gln Phe Trp Asp Ser Val Arg Leu Ala Leu  
                   20                  25                  30  
 Phe Thr Leu Ala Ile Val Ala Ile Ile Gly Ile Ala Ile Gly Ile Val  
                   35                  40                  45  
 Thr His Phe Val Val Glu Asp Lys Ser Phe Tyr Tyr Leu Ala Ser  
                   50                  55                  60  
 Phe Lys Val Thr Asn Ile Lys Tyr Lys Glu Asn Tyr Gly Ile Arg Ser  
   65                  70                  75                  80  
 Ser Arg Glu Phe Ile Glu Arg Ser His Gln Ile Glu Arg Met Met Ser  
                   85                  90                  95  
 Arg Ile Phe Arg His Ser Ser Val Gly Gly Arg Phe Ile Lys Ser His  
                   100                  105                  110  
 Val Ile Lys Leu Ser Pro Asp Glu Gln Gly Val Asp Ile Leu Ile Val  
                   115                  120                  125  
 Leu Ile Phe Arg Tyr Pro Ser Thr Asp Ser Ala Glu Gln Ile Lys Lys  
                   130                  135                  140  
 Lys Ile Glu Lys Ala Leu Tyr Gln Ser Leu Lys Thr Lys Gln Leu Ser  
   145                  150                  155                  160  
 Leu Thr Ile Asn Lys Pro Ser Phe Arg Leu Thr Pro Ile Asp Ser Lys  
                   165                  170                  175  
 Lys Met Arg Asn Leu Leu Asn Ser Arg Cys Gly Ile Arg Met Thr Ser  
                   180                  185                  190  
 Ser Asn Met Pro Leu Pro Ala Ser Ser Ser Thr Gln Arg Ile Val Gln  
                   195                  200                  205  
 Gly Arg Glu Thr Ala Met Glu Gly Glu Trp Pro Trp Gln Ala Ser Leu  
                   210                  215                  220  
 Gln Leu Ile Gly Ser Gly His Gln Cys Gly Ala Ser Leu Ile Ser Asn  
   225                  230                  235                  240  
 Thr Trp Leu Leu Thr Ala Ala His Cys Phe Trp Lys Asn Lys Asp Pro  
                   245                  250                  255

-15-

```

Thr Gln Trp Ile Ala Thr Phe Gly Ala Thr Ile Thr Pro Pro Ala Val
      260      265      270
Lys Arg Asn Val Arg Lys Ile Ile Leu His Glu Asn Tyr His Arg Glu
      275      280      285
Thr Asn Glu Asn Asp Ile Ala Leu Val Gln Leu Ser Thr Gly Val Glu
      290      295      300
Phe Ser Asn Ile Val Gln Arg Val Cys Leu Pro Asp Ser Ser Ile Lys
305      310      315      320
Leu Pro Pro Lys Thr Ser Val Phe Val Thr Gly Phe Gly Ser Ile Val
      325      330      335
Asp Asp Gly Pro Ile Gln Asn Thr Leu Arg Gln Ala Arg Val Glu Thr
      340      345      350
Ile Ser Thr Asp Val Cys Asn Arg Lys Asp Val Tyr Asp Gly Leu Ile
      355      360      365
Thr Pro Gly Met Leu Cys Ala Gly Phe Met Glu Gly Lys Ile Asp Ala
      370      375      380
Cys Lys Gly Asp Ser Gly Gly Pro Leu Val Tyr Asp Asn His Asp Ile
385      390      395      400
Trp Tyr Ile Val Gly Ile Val Ser Trp Gly Gln Ser Cys Ala Leu Pro
      405      410      415
Lys Lys Pro Gly Val Tyr Thr Arg Val Thr Lys Tyr Arg Asp Trp Ile
      420      425      430
Ala Ser Lys Thr Gly Met
      435

```

&lt;210&gt; 17

&lt;211&gt; 702

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)...(702)

&lt;223&gt; Nucleotide sequence encoding Protease Domain

&lt;400&gt; 17

```

att gtc caa gga agg gaa aca gct atg gaa ggg gaa tgg cca tgg cag      48
Ile Val Gln Gly Arg Glu Thr Ala Met Glu Gly Glu Trp Pro Trp Gln
  1          5          10          15

gcc agc ctc cag ctc ata ggg tca ggc cat cag tgt gga gcc agc ctc      96
Ala Ser Leu Gln Leu Ile Gly Ser Gly His Gln Cys Gly Ala Ser Leu
          20          25          30

atc agt aac aca tgg ctg ctc aca gca gct cac tgc ttt tgg aaa aat      144
Ile Ser Asn Thr Trp Leu Leu Thr Ala Ala His Cys Phe Trp Lys Asn
          35          40          45

aaa gac cca act caa tgg att gct act ttt ggt gca act ata aca cca      192
Lys Asp Pro Thr Gln Trp Ile Ala Thr Phe Gly Ala Thr Ile Thr Pro
          50          55          60

ccc gca gtg aaa cga aat gtg agg aaa att att ctt cat gag aat tac      240
Pro Ala Val Lys Arg Asn Val Arg Lys Ile Ile Leu His Glu Asn Tyr
          65          70          75          80

cat aga gaa aca aat gaa aat gac att gct ttg gtt cag ctc tct act      288
His Arg Glu Thr Asn Glu Asn Asp Ile Ala Leu Val Gln Leu Ser Thr
          85          90          95

```

-16-

gga gtt gag ttt tca aat ata gtc cag aga gtt tgc ctc cca gac tca	336
Gly Val Glu Phe Ser Asn Ile Val Gln Arg Val Cys Leu Pro Asp Ser	
100 105 110	
tct ata aag ttg cca cct aaa aca agt gtg ttc gtc aca gga ttt gga	384
Ser Ile Lys Leu Pro Pro Lys Thr Ser Val Phe Val Thr Gly Phe Gly	
115 120 125	
tcc att gta gat gat gga cct ata caa aat aca ctt cgg caa gcc aga	432
Ser Ile Val Asp Asp Gly Pro Ile Gln Asn Thr Leu Arg Gln Ala Arg	
130 135 140	
gtg gaa acc ata agc act gat gtg tgt aac aga aag gat gtg tat gat	480
Val Glu Thr Ile Ser Thr Asp Val Cys Asn Arg Lys Asp Val Tyr Asp	
145 150 155 160	
ggc ctg ata act cca gga atg tta tgt gct gga ttc atg gaa gga aaa	528
Gly Leu Ile Thr Pro Gly Met Leu Cys Ala Gly Phe Met Glu Gly Lys	
165 170 175	
ata gat gca tgt aag gga gat tct ggt gga cct ctg gtt tat gat aat	576
Ile Asp Ala Cys Lys Gly Asp Ser Gly Gly Pro Leu Val Tyr Asp Asn	
180 185 190	
cat gac atc tgg tac att gta ggt ata gta agt tgg gga caa tca tgt	624
His Asp Ile Trp Tyr Ile Val Gly Ile Val Ser Trp Gly Gln Ser Cys	
195 200 205	
gca ctt ccc aaa aaa cct gga gtc tac acc aga gta act aag tat cga	672
Ala Leu Pro Lys Lys Pro Gly Val Tyr Thr Arg Val Thr Lys Tyr Arg	
210 215 220	
gat tgg att gcc tca aag act ggt atg tag	702
Asp Trp Ile Ala Ser Lys Thr Gly Met *	
225 230	

<210> 18  
 <211> 233  
 <212> PRT  
 <213> Homo sapien

<400> 18  
 Ile Val Gln Gly Arg Glu Thr Ala Met Glu Gly Glu Trp Pro Trp Gln  
 1 5 10 15  
 Ala Ser Leu Gln Leu Ile Gly Ser Gly His Gln Cys Gly Ala Ser Leu  
 20 25 30  
 Ile Ser Asn Thr Trp Leu Leu Thr Ala Ala His Cys Phe Trp Lys Asn  
 35 40 45  
 Lys Asp Pro Thr Gln Trp Ile Ala Thr Phe Gly Ala Thr Ile Thr Pro  
 50 55 60  
 Pro Ala Val Lys Arg Asn Val Arg Lys Ile Ile Leu His Glu Asn Tyr  
 65 70 75 80  
 His Arg Glu Thr Asn Glu Asn Asp Ile Ala Leu Val Gln Leu Ser Thr  
 85 90 95  
 Gly Val Glu Phe Ser Asn Ile Val Gln Arg Val Cys Leu Pro Asp Ser  
 100 105 110  
 Ser Ile Lys Leu Pro Pro Lys Thr Ser Val Phe Val Thr Gly Phe Gly

